



Full Length Article

High Grain Diet Triggers Inflammation in the Goat Uterus: a Comprehensive Regulation Diet Modulates the Immune Response

Muhammad Shahid Bilal^{1†}, Juma Ahamed Abaker^{1†}, Zain ul Aabdin¹, Usman Waheed², Tianle Xu¹, Hongyu Dai¹, Guo Junfei¹ and Xiangzhen Shen^{1*}

¹College of Veterinary Medicine, Nanjing Agricultural University, Nanjing 210095, P.R. China

²Department of Pathobiology, University of Veterinary and Animal Sciences, Sub-campus Jhang, Pakistan

*For correspondence: xzshen@njau.edu.cn

†These authors contributed equally to this work

Abstract

The aim of the present study was to explore the causal association between the increase in circulating subacute ruminal acidosis (SARA)-induced lipopolysaccharide (LPS) levels and uterine inflammation; moreover this study sought to evaluate the role of a comprehensive regulation diet in combating the inflammatory effects of a high grain diet. Twelve dairy goats were randomly assigned to two groups. One group was provided a high-concentrate buffered diet as the comprehensive regulation group (RG) control while, only a high concentrate powdered diet was fed to the treatment group (TG). The LPS concentrations in the rumen fluid were 428702 ± 34505.6 and 255027 ± 39756.2 EU/mL, in the TG and RG goats, respectively, while, the LPS levels in the peripheral plasma were 0.032 ± 0.003 and 0.012 ± 0.003 EU/mL in the TG and RG goats, respectively. The concentration of TNF- α , IL-1 β and IL-6 was significantly higher, and there was enhanced expression of the inflammatory genes *TLR4*, *MyD88*, *TRAF-6*, *NF- κ B*, *IL-1 β* and *IL-6* in the uteri of TG goats. SARA increased the NF- κ B expression and level of its phosphorylated form in TG goats. The uteri of TG goats showed higher mRNA levels of pro-inflammatory genes than the RG goats, indicating the presence of inflammation. Thus, SARA-induced LPS derived from a high concentrate diet triggers inflammation in the goat uterus by reinforcing *TLR4* signaling. © 2016 Friends Science Publishers

Keywords: Comprehensive regulation diet; High concentrate diet; Lipopolysaccharide; Sub-acute ruminal acidosis; Uterus

Introduction

In the dairy sector, a high concentrate diet is mostly fed to dairy animals to increase milk production (Jing *et al.*, 2014). The prolonged feeding of a high grain diet leads to subacute ruminal acidosis (SARA) (Dong *et al.*, 2011), which is described as moderately decreased ruminal pH that varies from 5.2 to 5.6 (Gozho *et al.*, 2005). Moreover, the high concentrate diet, due to a reduction in the ruminal pH, results in change in the microbial population from cellulose-digesting to starch-digesting (Metzler-Zebeli *et al.*, 2013). Many of the starch digesting microorganisms are gram-negative bacteria (Bryant, 1970). The concentration of lipopolysaccharide (LPS) in the rumen is altered due to this change in population (Dehority, 2003). Previous experiments with grain-based SARA revealed that LPS is liberated in large quantities in the gastrointestinal tract (GIT) of the affected animals due to the collapse of gram-negative bacteria (Khafipour *et al.*, 2009; Dong *et al.*, 2011). The ruminal epithelium works as a defensive barrier between the ruminal environment and portal circulation (Graham and Simmons, 2005). The high acidity of the ruminal

environment causes damage and sloughing of the epithelial cells in the main gut (Plaizier *et al.*, 2008) and the permeability of the gut might increase as a result of the rise in LPS levels (Khafipour *et al.*, 2009). Moreover, it was observed that high concentrate feeding increased the free LPS level in the rumen; LPS subsequently translocated into the blood stream with the possible major site of LPS translocation into the blood stream being the intestines (Khafipour *et al.*, 2009). Recently, Chang *et al.* (2015) reported that a harmful feature of high grain diet-induced SARA is the induction of liver inflammation. The high rise in circulating LPS triggers inflammation that in turn stimulates extensive *TLR4* signaling and ultimately elevates the expressions of innate immune genes and enhances the secretion of pro-inflammatory cytokines.

The uterus, especially the endometrial lining, has a major role in normal reproductive cycles, the implantation of embryos and the support of a healthy foetus (Turner *et al.*, 2012). Gram-negative bacteria are the primary initiators of bovine endometrial infection, causing uterine disease (Cronin *et al.*, 2012) that results in tissue damage to the endometrium (Sheldon *et al.*, 2009). Endometrial epithelial

cells (EECs) exclusively participate in clearing the attacking microorganisms after the identification of pathogen-associated molecular patterns (PAMPs) by pattern recognition receptors (PRR). The LPS from gram-negative bacteria is recognized by toll-like receptor 4 (*TLR4*), which initiates innate immune responses (Fu *et al.*, 2013). *TLR4* recognizes LPS with the help of LPS-binding protein (*LBP*) and cluster of differentiation antigen 14 (*CD14*) (Bannerman *et al.*, 2004; Sohn *et al.*, 2008). LPS triggers the activation of myeloid differentiating factor 88 (*MyD88*) after binding to *TLR4* on the cell surface of the host (Ju *et al.*, 2014). *MyD88* activates TNF receptor-associated factor 6 (*TRAF6*), which stimulates the I κ B kinase (*I κ K*) complex; consequently, phosphorylated nuclear transcription factor (*NF- κ B*) translocates from the cytoplasm to the nucleus (Turner *et al.*, 2012). The NF- κ B-dependent signaling pathway leads to the production of pro-inflammatory cytokines, including interleukins *IL-1 β* , *IL-6*, chemokines (*IL-8*) and other inflammatory mediators (Cronin *et al.*, 2012).

Sodium bicarbonate is normally supplemented in the diets of the ruminants to neutralize ruminal pH (Phy and Provenza, 1998). Magnesium oxide, sodium sesquicarbonate, and sodium bicarbonate together increase the pH of the ruminal fluid and increase its buffering ability, stabilizing the acid-base status of the rumen (Le Ruyet and Tucker, 1992). Phy and Provenza (1998) illustrated that small ruminants preferred pelletized fermentable concentrate diets supplemented with sodium bicarbonate. Dietary sodium butyrate decreased digestive tract permeability and facilitated considerable alterations in intestinal microbial population (Huang *et al.*, 2015) in addition to reducing colonic inflammation (Wirtz and Neurath, 2007).

Therefore, the present study focused on the following objectives: (I) to evaluate changes in the expression of inflammation associated genes in the goat uterus due to LPS derived from the digestive tract after extended of a high concentrate diet and (II) to study whether the comprehensive regulation buffered diet can reduce the negative effect of a high concentrate diet in ruminants.

Materials and Methods

Ethical Approval

The sampling procedures and experimental design were approved by the Animal Ethics Committee of Nanjing Agricultural University, China before the start of this experiment.

Animals, Diets and Experimental Design

Twelve multiparous mid-lactation dairy goats exhibiting an average body weight of 39 ± 6 kg were selected for the current study. They were randomly divided into two groups, the comprehensive regulation group (RG) and the treatment

group (TG), with each group containing 6 goats. The RG goats were fed a high concentrate palletized diet (6:4 concentrate/roughage ratio) with the addition of uniformly mixed buffer agents, while the TG goats were offered only a high concentrate powdered diet (6:4 concentrate/roughage ratio) for the 24-week experimental period. All the dairy goats were ruminally cannulated at 14th week. The animals were housed in individual tie stalls. Feed was offered two times daily at 08:00 and at 18:00 h, while fresh water was available to them at all times throughout the entire experimental period. The experiment was started after the animals made a full recovery, and no goat showed any clinical signs of infection during the experiment.

The ingredient details and buffer contents used in the diets are presented in Tables 1 and 2.

Sample Collection and Analysis

Blood samples and rumen fluid were collected at the 23rd week. Blood samples from all the goats were taken using 5-mL sterile Vacutainer tubes containing sodium heparin. The plasma was separated from the blood samples by centrifugation at $3000 \times g$ at 4°C for 15 min and stored at -20°C to determine the pro-inflammatory cytokines in the peripheral blood. The rumen fluid was collected in a depyrogenated, sterilized glass tube (previously heated at 250°C for 2 h). The pH of the ruminal fluid was immediately measured with a pH-metre (Sartorius, Basic pH Meter PB-10, PB-21, Goettingen, Germany) at 0 h, 1 h, 2 h, 4 h, 6 h, 8 h and 10 h after feeding after filtration through four layers of sterilized cheesecloth. At the end of the experiment, the goats were slaughtered after overnight fasting and uterine tissues from all the goats were carefully removed immediately after slaughter, flash and liquid nitrogen and stored at -70°C.

Determination of Ruminal pH

The ruminal pH was measured with a pH-metre.

Determination of LPS

The LPS concentrations in the plasma and rumen fluid were determined with Chromogenic endpoint Limulus Amebocyte Lysate assay kits, CE64406 and CE80545 (Chinese Horseshoe Crab Reagent Manufactory Co., Ltd., Xiamen, China), respectively, as reported by Dong *et al.* (2013). These assays have a minimum detection limit of 0.01 EU/mL.

Radioimmunoassay

Radioimmunoassay (RIA) kits were used to measure the concentration of the pro-inflammatory cytokines IL-1 β , IL-6 and TNF- α in the peripheral blood (IL-1 β , C09DJB; IL-6, C12DJB; TNF- α , C06PJB); Beijing North Institute of Biological Technology, Beijing, China.

Table 1: The ingredients and nutritional composition of the diets

| Items | Ingredients | Quantity |
|-------------------|---------------------------------|----------|
| Concentrate | Corn (g/100 g) | 23.07 |
| | Bran (g/100 g) | 28.03 |
| | Soyabean meal (g/100 g) | 2 |
| | Rape seed meal (g/100 g) | 3.7 |
| | Limestone (g/100 g) | 1.43 |
| | Dicalcium phosphate (g/100 g) | 0.6 |
| | Premix (g/100 g) | 0.4 |
| | Salt (g/100 g) | 0.5 |
| Coarse material | Leymaschinensis (g/100 g) | 32 |
| | Alfalfa (g/100 g) | 8 |
| Nutritional index | Net energy (MJ/kg) | 17.56 |
| | Water content | 1.89 |
| | Ash (%) | 8.22 |
| | Fat (%) | 3.6 |
| | Protein (%) | 16.76 |
| | Neutral detergent fiber NDF (%) | 42.64 |
| | Acid detergent fiber ADF (%) | 7.22 |
| | Ca (%) | 1.55 |
| P (%) | 0.76 | |

*The premix formulation per kilogram of diet containing VA, 6000ku/kg; VD, 2500 ku/kg; VE, 80 mg/kg; Cu, 6.25 mg/kg; Fe, 62.5 mg/kg; Zn, 62.5 mg/kg; Mn, 50 mg/kg; I, 125 mg/kg; Co, 0.125 mg/kg; Se, 27 mg/kg. M, 0.125 mg/kg

Table 2: Formulation of diets for the treatment group (TG) and the regulation group (RG)

| | Concentrate (Powdered) (g) | Concentrate (Pelletized) (g) | Roughage (g) | | | Buffer agents (g) | | |
|-----------------------|----------------------------|------------------------------|--------------|-----------|-----------------|--------------------|----------------|-----------------|
| | | | Alfalfa | Oat grass | Magnesium oxide | Sodium bicarbonate | Sodium acetate | Sodium butyrate |
| Treatment group (TG) | 600 | - | 133 | 267 | - | - | - | - |
| Regulation group (RG) | - | 600 | 133 | 267 | 6 | 9 | 10 | 10 |

Table 3: Primer sequences for real-time polymerase chain reaction (qRT-PCR) amplification of mRNA

| Gene name | Gene Bank accession | Primer sequences (5' -3') | Length (bp) |
|---------------------------------|---------------------|----------------------------|-------------|
| <i>β-actin</i> | JQ409458.1 | F: CTTGATGTCACGGACGATTT | 380 |
| | | R: CACGGCATGTGCACCAACT | |
| <i>NF-κB</i> | XM_00569893.1 | F: AGGTGGCGATCGTTGTCTA | 226 |
| | | R: TTGCCTTTGTCTTCCTGCC | |
| <i>MyD88</i> | XM_005695671.1 | F: TGGACACATACAAGCCCACT | 171 |
| | | R: AGTGTCACATCTTGTCAGG | |
| <i>TRAF-6</i> | XM_005690091.1 | F: TTTCAGAGACCCACCATCCC | 166 |
| | | R: GAAAACCCCTCCCTCCGAAGA | |
| <i>IL-1β</i> | XM_005686747.1 | F: CCGTGATGATGACCTGAGGAG | 303 |
| | | R: CAAGACAGGTATAGATTCTTGC | |
| <i>TLR-4</i> | NM_001285574.1 | F: CTGAGAACCGAGAGCTGGGAC | 207 |
| | | R: TTCGCATCTGGATAAATCCAGC | |
| <i>IL-6</i> | JF432019.1 | F: CGAAGCTCTCATTAAAGCACATC | 127 |
| | | R: CCAGGTATATCTGATACTCCAG | |

RNA Extraction and Quantitative Real-time PCR

RNA extraction: Total RNA from 100 mg of uterine tissue was extracted with the RNA iso Plus™ reagent (Takara Co., Otsu, Japan) using a Dounce homogenizer (Polytron PT 1200 E, Switzerland). The concentration of RNA in every sample was determined using a spectrophotometer (Eppendorf Biotechnology, Hamburg, Germany) at 260 nm. First-strand cDNA was prepared from 250 ng/ μ L of total RNA template using the Prime Script RT Master Mix Perfect Real Time kit (Takara Co., Otsu, Japan).

Quantitative Real Time PCR (qRT-PCR)

Primers for the *TLR4*, *MyD88*, *TRAF6*, and *NF- κ B*, *IL-1 β* , *IL-6* and *β -actin* genes were designed using Premier 6.0 (Premier Biosoft International, USA). The reaction mixture comprised 0.4 μ M primers and 2 μ L of cDNA in a final volume of 20 μ L of supermix. The thermal cycling parameters consisted of an initial denaturation step at 95°C for 15 s, followed by 40 cycles of annealing at 95°C for 5 sec. and primer extension at 60°C for 31 s. All the reactions were run in triplicate. The PCR products were verified by gel electrophoresis with 3.0% agarose gel.

Each cDNA sample was amplified using SYBR Green (Takara Co., Otsu, Japan) on the ABI 7300 Fast Real-Time PCR System (Applied Bio system, USA). The data were normalized to the mean of the internal reference β -actin gene to control unpredictability in expression levels and were evaluated using the $R = 2^{-\Delta\Delta C_t}$ method (Livak and Schmittgen, 2001). The primer sequences are presented in Table 3.

Western Blot Analysis

Approximately 100 mg of the frozen uterine tissue was homogenized in 1 mL of ice-cold RIPA protein isolation buffer (Beyotime, Shanghai, China) for 15 s, incubated on ice for 20 min and centrifuged at 12000 rpm for 30 min at 4°C. Care was taken to avoid disturbing the pellet at the bottom when the supernatant was shifted to another tube. The protein concentration was measured using a BCA Protein Assay kit (Thermo Fisher Scientific Inc., USA) and was adjusted to 4 μ g/ μ L. Equal protein amounts were loaded for separation with 10% SDS-polyacrylamide gel electrophoresis (PAGE) before being transferred to a nitrocellulose (NC) membrane (Pall Gelman Laboratory, Ann Arbor, USA) at 4°C. Afterward, the membranes were blocked in 5% skimmed milk (5% BSA for phosphorylated p65) for 2 h at room temperature, followed by incubation together anti-NF- κ B (p65) (1:300, SC-109, Santa Cruz Biotechnology, USA), anti-p NF- κ B (p-p65) (1:300, SC-33020, Santa Cruz Biotechnology, USA) and GAPDH primary antibodies (1:10000, Bioworld Technology, Inc., Louis Park, USA), at 4°C overnight. The membranes were subsequently washed in tris-buffered saline containing Tween20 (TBST) followed by incubation with horseradish peroxidase (HRP)-conjugated goat-anti-rabbit (1:5000, Santa Cruz Biotechnology, USA) and goat anti-mouse secondary antibodies (1:10000, Sunshine Biotechnology (Nanjing) Co., Ltd, China), at room temperature for 2 h. After washing the membranes 6 times (10 min/wash), the results were developed with an ECL Plus kit (Vazyme, Nanjing, China). The signals were visualized on an imaging system (LAS4000, USA) and evaluated with Quantity One software (Bio-Rad, USA).

Statistical Analyses

All analyses were performed with SPSS statistical software version 16.0 (SPSS Inc., Chicago, IL, USA) and an independent *t*-test was used to evaluate the differences in the mean \pm SEM between the groups. The mean differences were considered significant when $P < 0.05$.

Results

Ruminal pH and LPS Content of Ruminal Fluid and Blood Plasma

The pH of the rumen fluid is presented in Fig. 1. The ruminal pH in TG goats was less than 5.8 for more than 3 h,




Fig. 1: Comparison of the pH values in the ruminal fluid between regulation group (RG) and treatment group (TG) goats

The data were measured during the 23rd week using ruminal fluid samples collected at times ranging from 0 to 10 h (shown on the x-axis). Student's *t*-test was used to evaluate the significance of treatment; significant variations were observed across all of the sampling times ($P < 0.05$)

indicating that SARA was elicited effectively. Extended feeding of a high grain diet linearly reduced the dynamic pH curve in the TG goats relative to the RG goats throughout the experimental period. The ruminal pH of the TG goats was significantly less than that in the RG goats ($P < 0.05$). The LPS concentration in the rumen fluid of TG goats was 428702 ± 34505.6 EU/mL, while that in RG goats was 255027 ± 39756.2 EU/mL, indicating a significant increase in LPS level in TG goats compared to RG goats ($P < 0.02$). Similarly, the LPS level in the peripheral plasma of TG goats 0.032 ± 0.003 EU/mL was significantly higher than that in the RG goats 0.012 ± 0.003 EU/mL ($P < 0.002$) (Table 4).

Pro-inflammatory Cytokine Levels in the Peripheral Blood

The concentrations of the pro-inflammatory cytokines IL-1 β , IL-6 and TNF- α were determined with radioimmunoassay and the results are shown in Fig. 2. The concentration of IL-1 β was 0.2167 ± 0.0155 ng/mL in TG goats and 0.1700 ± 0.0041 ng/mL in RG goats ($P < 0.02$) (Fig. 2a); the IL-6 concentration was 607.3 ± 4.5 pg/mL in TG goats and 401.5 ± 69.9 pg/mL in RG goats ($P < 0.02$) (Fig. 2b) and the TNF- α concentration was 7.603 ± 0.531 fmol/mL in TG goats and 2.628 ± 1.120 fmol/mL in RG goats ($P < 0.007$) (Fig. 2c). The concentration of pro-inflammatory cytokines in the blood was significantly higher in the TG goats relative to the RG goats ($P < 0.05$).

Table 4: Lipopolysaccharide concentration in the rumen and plasma of treatment group (TG) and regulation group (RG) dairy goats fed a high concentrate diet

| LPS Concentration (EU/mL) | Treatment group (TG) | | Regulation group (RG) | | P-value |
|---------------------------|----------------------|---------|-----------------------|---------|---------|
| | Mean | SEM | Mean | SEM | |
| Rumen LPS (EU/mL) | 428702 | 34505.6 | 255027 | 39756.2 | 0.02 |
| Plasma LPS (EU/mL) | 0.032 | 0.003 | 0.012 | 0.003 | 0.002 |

SEM=Standard error of mean between the two groups; The LPS data were compared using Student's *t*-test

Expressions of Genes Involved in Inflammation of the Uterus

As presented in Fig. 3, the high concentrate diet induced SARA altered the expression of inflammation related genes in the uteri of TG goats relative to the RG goats. The gene expression of *TLR4* and *IL6* was significantly higher in the TG goats compared to RG goats ($P<0.01$). Similarly, the *IL-1 β* and *MyD88* levels were significantly higher in the TG goats than the RG goats ($P<0.01$). Moreover, the expression of *TRAF6* and *NF- κ B* was significantly elevated in TG goats relative to RG goats ($P<0.01$).

Protein expression of NF- κ B in the Goat Uterus of the Treatment Group and Regulation Group

Western blot analysis verified that high concentrate diet induced SARA significantly enhanced the levels of NF- κ B (p65) ($P<0.001$) and phosphorylated NF- κ B (p-p65) ($P<0.003$) protein in the uteri of TG goats compared to the RG goats (Fig. 4). Thus, it was confirmed that the *NF- κ B* p65 signaling pathway was activated.

Discussion

In the current research, we highlighted the pivotal role of inflammatory genes and the mechanism underlying their expression in the uteri of goats; this change resulted from the higher serum concentration of LPS induced by the extended feeding of a high concentrate diet. It has been documented that the feeding of a high concentrate diet results in digestive disturbance, especially, SARA (Gozho *et al.*, 2005), which is described as decreased feed intake, low milk fat and inflammation characterized by a rise in acute phase proteins (Khafipour *et al.*, 2009). The key indicator of SARA in many animals is the low pH of the rumen fluids (Khafipour *et al.*, 2009). The current study revealed that the dynamic pH curve for the ruminal fluid of TG goats was less than that of the RG goats, depicting the onset of SARA in the TG goats. This is in agreement with the results of Ghorbani *et al.* (2002), who proposed that the an average pH value below 5.8 in TG goats for more than three hours represented confirmation of SARA. However, there is little literature showing that SARA not only depends on pH but on the change in the microbial population that occurs secondarily to the feed type (Calsamiglia *et al.*, 2012). In fact low pH values in the rumen resulted in the lysis of gram-negative bacteria, which causes LPS to be released and free to enter the systemic circulation




Fig. 2: *IL-1 β* , *IL-6* and *TNF- α* levels in peripheral blood. The data are expressed as the mean \pm SEM; asterisks indicate differences between the goats in the regulation group (RG) and treatment group (TG) (* $P<0.05$, ** $P<0.01$, $n=12$)




Fig. 3: Uterine gene expression profile analysis with qRT-PCR

The genes involved in transcriptional regulation were measured in the uterine tissue. The error bars indicate the standard error of the mean and ** indicates significance at $P<0.01$

(Andersen, 2000). Moreover, feeding high concentrate diets to animals places their GIT at a high risk of impairment due to perturbations in the functions of the ruminal epithelium (Tao *et al.*, 2014). Due to epithelial damage and alterations in GIT permeability, LPS might also translocate into the circulation where this rise in LPS level triggers an immune response, which is in accord with earlier studies (Dong *et al.*, 2011; Khafipour *et al.*, 2009). This is the main reason why many studies on dairy animals recommended an appropriate coarse fibre buffered diet for good rumination and enhanced saliva secretions to overcome ruminal acidosis (Mertens, 1997; Calitz, 2009). The clinical evidence indicated that the supplementation of buffering agents, bicarbonates and carbonates including sodium bicarbonate (NaHCO_3),




Fig. 4: Western blot for NF- κ B (p65) and phosphorylated-p65 protein in the uterus

The NF- κ B level in the uteri of regulation group (RG) and treatment group (TG) goats was evaluated using western-blot (a). The protein quantities were measured as arbitrary units relative to GAPDH; fold alterations in NF- κ B (p65) and phosphorylated NF- κ B (p-p65) are shown (b). The data are expressed as the mean \pm SEM; asterisks indicate differences between the treatment group (TG) group and the regulation group (RG) goats (** $P < 0.01$)

magnesium oxide (MgO), potassium bicarbonate (KHCO₃) and potassium carbonate (K₂CO₃) are of significant value in increasing the ruminal pH of SARA affected animals (Erdman, 1988). It is well recognized that the cows fed a buffered diet have a higher ruminal pH (Ghorbani *et al.*, 1989). Notably, stromal and endometrial epithelial cells have an essential role in inflammatory gene expression in the reproductive tract due to their interactions with LPS (Sheldon and Roberts, 2010). This inflammation is mediated by the release of particular mediators which ultimately enhance the production and release of the immune cell infiltrates that are responsible for marked cellular changes in the epithelial cells (Chapwanya *et al.*, 2009).

In the present study, we sought to explore the influence of SARA derived endogenous LPS at the molecular level and observed increased expression of pro-inflammatory cytokines, probably because of a high LPS concentration in the peripheral plasma (Baumann and Gauldie, 1994). The LPS activates the immune system during the stimulation of cytokine release (Netea *et al.*, 2002). Our findings showed that the levels of the pro-inflammatory cytokines TNF- α , IL-1 β , and IL-6 were significantly higher in the peripheral blood of the TG goats. This finding was in agreement with the earlier reports of (Lukacs *et al.*, 1995; Dörffel *et al.*, 1999).

Moreover, RT-qPCR based analysis of inflammatory markers in the goat uterus confirmed that the TG goats suffered from uterine inflammation. TLRs recognize the PAMPs of gram-negative bacteria; this recognition subsequently enhances TLR expression, the most important

part of the innate immune system (Akira *et al.*, 2006). Our data confirmed that the LPS induced by TG goats, showed TLR4 elevation compared to the RG goats. In agreement with previous studies (Fu *et al.*, 2013; Ju *et al.*, 2014), our research illustrated that when bovine endometrial epithelial cells are challenged with LPS, TLR4 molecules are significantly regulated.

Furthermore, a significant increase in the expression of the adapter molecule MyD88 was observed in TG goats versus RG goats (Fu *et al.*, 2013; Ju *et al.*, 2014). Many reports have shown that IL-1 receptor-associated kinase (IRAK) is recruited by MyD88 triggering TRAF6, which stimulates the I κ B kinase (I κ K) complex (Fu *et al.*, 2013). Similar findings were also observed in this study, as higher TRAF6 expression was found in TG goats relative to RG goats. TRAF6 triggers inhibitory- κ B (I κ B) phosphorylation, resulting in the degradation of I κ B proteins; consequently, NF- κ B translocates from the cytoplasm to the nucleus to regulate the transcription of inflammatory genes (Kawai and Akira, 2007). Substantial evidence has proven that high expression of NF- κ B in the uterine tissue was due to inflammation of the endometrium (Chapwanya *et al.*, 2009). Our results also proved that SARA derived LPS triggers NF- κ B signaling causing endometrial inflammation.

The expression levels of both NF- κ B (p65) and phosphorylated p65 proteins were noticeably elevated in the TG goats compared with the RG goats. Taken together, the distinctive base of above mentioned facts firmly suggests that LPS triggered the activation of TLR4 signaling, resulting in an inflammatory response in the uterus of goats suffering from SARA. Overwhelmingly, the data in the current study indicate that LPS triggers the TLR4-NF- κ B signaling pathway, in accord with an earlier study (Chang *et al.*, 2015). The present study found that SARA increased the mRNA expression and protein concentrations of NF- κ B in the uterus of TG goats. It was also observed that TLR4 exerted a pro-inflammatory effect by activating NF- κ B and these results confirm that the TLR4-dependent activation of the NF- κ B pathway plays a key role in the endometrial immune response (Ju *et al.*, 2014).

Conclusion

It has been unmistakably suggested that the extended feeding of a high concentrate diet alters inflammatory gene expression in the uteri of dairy goats. The feeding of a comprehensive regulation buffered diet will be beneficial in reducing SARA and has been proved to be effective in combating inflammatory effects and ultimately endometritis in ruminants. However, further studies are needed to determine further prophylactic measures to overcome SARA-induced uterine inflammation.

Acknowledgements

This study was supported by grants from the National Natural Science Foundation of China (31172371) and the

Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD).

References

- Akira, S., S. Uematsu and O. Takeuchi, 2006. Pathogen recognition and innate immunity. *Cell*, 124: 783-801
- Andersen, P., 2000. Bovine endotoxemia: Aspects of relevance to ruminal acidosis. *Dr. Vet. Sci. Thesis*, The Royal Veterinary and Agricultural University, Copenhagen, Denmark
- Bannerman, D.D., M. Paape, W. Hare and J.C. Hope, 2004. Characterization of the bovine innate immune response to intramammary infection with *Klebsiella pneumoniae*. *J. Dairy Sci.*, 87: 2420-2432
- Baumann, H. and J. Gauldie, 1994. The acute phase response. *Immunol. Today*, 15: 74-80
- Bryant, M., 1970. Normal flora—rumen bacteria. *Am. J. Clin. Nutr.*, 23: 1440-1450
- Calitz, T., 2009. *The Effect of Acid Buf and Combinations of Acid Buf and Socium Bicarbonate in Dairy Cow Diets on Production Response and Rumen Parameters*. University of Stellenbosch, Stellenbosch
- Calsamiglia, S., M. Blanch, A. Ferret and D. Moya, 2012. Is subacute ruminal acidosis a pH related problem? Causes and tools for its control. *Anim. Feed Sci. Technol.*, 172: 42-50
- Chang, G., S. Zhuang, H.M. Seyfert, K. Zhang, T. Xu, D. Jin, J. Guo and X. Shen, 2015. Hepatic TLR4 signaling is activated by LPS from digestive tract during SARA, and epigenetic mechanisms contribute to enforced TLR4 expression. *Oncotarget*, 6: 38578-38590
- Chapwanya, A., K.G. Meade, M.L. Doherty, J.J. Callanan, J. F. Mee and C. O'Farrelly, 2009. Histopathological and molecular evaluation of Holstein-Friesian cows postpartum: Toward an improved understanding of uterine innate immunity. *Theriogenology*, 71: 1396-1407
- Cronin, J.G., M.L. Turner, L. Goetze, C.E. Bryant and I.M. Sheldon, 2012. Toll-like receptor 4 and MYD88-dependent signaling mechanisms of the innate immune system are essential for the response to lipopolysaccharide by epithelial and stromal cells of the bovine endometrium. *Biol. Reprod.*, 86: 51
- Dehority, B., 2003. *Starch Digesters, Other Less Numerous Species, and Facultative Anaerobes in the Rumen*. Nottingham University Press, (UK)
- Dong, G., S. Liu, Y. Wu, C. Lei, J. Zhou and S. Zhang, 2011. Diet-induced bacterial immunogens in the gastrointestinal tract of dairy cows: Impacts on immunity and metabolism. *Acta Vet Scand.*, 53: 48
- Dong, H., S. Wang, Y. Jia, Y. Ni, Y. Zhang, S. Zhuang, X. Shen and R. Zhao, 2013. Long-term effects of subacute ruminal acidosis (SARA) on milk quality and hepatic gene expression in lactating goats fed a high-concentrate diet. *PLoS ONE*, 8: e82850
- Dörffel, Y., C. Lätsch, B. Stuhlmüller, S. Schreiber, S. Scholze, G. R. Burmester and J. Scholze, 1999. Preactivated peripheral blood monocytes in patients with essential hypertension. *Hypertension*, 34: 113-117
- Erdman, R.A., 1988. Dietary buffering requirements of the lactating dairy cow: a review. *J. Dairy Sci.*, 71: 3246-3266
- Fu, Y., B. Liu, X. Feng, Z. Liu, D. Liang, F. Li, D. Li, Y. Cao, S. Feng and X. Zhang, 2013. Lipopolysaccharide increases Toll-like receptor 4 and downstream Toll-like receptor signaling molecules expression in bovine endometrial epithelial cells. *Vet. Immunol. Immunopathol.*, 151: 20-27
- Ghorbani, G., J. Jackson and R. Hemken, 1989. Effects of sodium bicarbonate and sodium sesquicarbonate on animal performance, ruminal metabolism, and systemic acid-base status. *J. Dairy Sci.*, 72: 2039-2045
- Ghorbani, G., D. Morgavi, K. Beauchemin and J. Leedle, 2002. Effects of bacterial direct-fed microbials on ruminal fermentation, blood variables, and the microbial populations of feedlot cattle. *J. Anim. Sci.*, 80: 1977-1985
- Gozho, G., J. Plaizier, D. Krause, A. Kennedy and K. Wittenberg, 2005. Subacute ruminal acidosis induces ruminal lipopolysaccharide endotoxin release and triggers an inflammatory response. *J. Dairy Sci.*, 88: 1399-1403
- Graham, C. and N.L. Simmons, 2005. Functional organization of the bovine rumen epithelium. *Am. J. Physiol. Regul. Integr. Comp. Physiol.*, 288: 19
- Huang, C., P. Song, P. Fan, C. Hou, P. Thacker and X. Ma, 2015. Dietary Sodium Butyrate Decreases Postweaning Diarrhea by Modulating Intestinal Permeability and Changing the Bacterial Communities in Weaned Piglets. *J. Nutr.*, 145: 2774-2780
- Jing, L., R. Zhang, Y. Liu, W. Zhu and S. Mao, 2014. Intravenous lipopolysaccharide challenge alters ruminal bacterial microbiota and disrupts ruminal metabolism in dairy cattle. *Brit. J. Nutr.*, 112: 170-182
- Ju, J., L. Li, J. Xie, Y. Wu, X. Wu and W. Li, 2014. Toll-like receptor-4 pathway is required for the pathogenesis of human chronic endometritis. *Exp. Ther. Med.*, 8: 1896-1900
- Kawai, T. and S. Akira, 2007. Signaling to NF- κ B by Toll-like receptors. *Trends Mol. Med.*, 13: 460-469
- Khafipour, E., D. Krause and J. Plaizier, 2009. A grain-based subacute ruminal acidosis challenge causes translocation of lipopolysaccharide and triggers inflammation. *J. Dairy Sci.*, 92: 1060-1070
- Le Ruyet, P. and W.B. Tucker, 1992. Ruminant buffers: temporal effects on buffering capacity and pH of ruminal fluid from cows fed a high concentrate diet. *J. Dairy Sci.*, 75: 1069-1077
- Livak, K.J. and T.D. Schmittgen, 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods*, 25: 402-408
- Lukacs, N.W., R.M. Strieter, S.W. Chensue, M. Widmer and S.L. Kunkel, 1995. TNF-alpha mediates recruitment of neutrophils and eosinophils during airway inflammation. *J. Immunol.*, 154: 5411-5417
- Mertens, D., 1997. Creating a system for meeting the fiber requirements of dairy cows. *J. Dairy Sci.*, 80: 1463-1481
- Metzler-Zebeli, B.U., S. Schmitz-Esser, F. Klevenhusen, L. Podstatzky-Lichtenstein, M. Wagner and Q. Zebeli, 2013. Grain-rich diets differently alter ruminal and colonic abundance of microbial populations and lipopolysaccharide in goats. *Anaerobe*, 20: 65-73
- Netea, M.G., M. van Deuren, B.J. Kullberg, J.M. Cavaillon and J.W. Van der Meer, 2002. Does the shape of lipid A determine the interaction of LPS with Toll-like receptors? *Trends Immunol.*, 23: 135-139
- Phy, T.S. and F.D. Provenza, 1998. Eating barley too frequently or in excess decreases lambs' preference for barley but sodium bicarbonate and lasalocid attenuate the response. *J. Anim. Sci.*, 76: 1578-1583
- Plaizier, J., D. Krause, G. Gozho and B. McBride, 2008. Subacute ruminal acidosis in dairy cows: The physiological causes, incidence and consequences. *Vet. J.*, 176: 21-31
- Sheldon, I.M., J. Cronin, L. Goetze, G. Donofrio and H.J. Schuberth, 2009. Defining postpartum uterine disease and the mechanisms of infection and immunity in the female reproductive tract in cattle. *Biol. Reprod.*, 81: 1025-1032
- Sheldon, I.M. and M.H. Roberts, 2010. Toll-like receptor 4 mediates the response of epithelial and stromal cells to lipopolysaccharide in the endometrium. *PLoS ONE*, 5: e12906
- Sohn, M.J., G.M. Hur, H.S. Byun and W.G. Kim, 2008. Cyclo (dehydrohistidyl-L-tryptophyl) inhibits nitric oxide production by preventing the dimerization of inducible nitric oxide synthase. *Biochem. Pharmacol.*, 75: 923-930
- Tao, S., Y. Duanmu, H. Dong, J. Tian, Y. Ni and R. Zhao, 2014b. A high-concentrate diet induced colonic epithelial barrier disruption is associated with the activating of cell apoptosis in lactating goats. *BMC Vet. Res.*, 10: 014-0235
- Turner, M., G. Healey and I. Sheldon, 2012. Immunity and inflammation in the uterus. *Reprod. Domest. Anim.*, 47: 402-409
- Wirtz, S. and M.F. Neurath, 2007. Mouse models of inflammatory bowel disease. *Adv. Drug. Deliv. Rev.*, 59: 1073-1083

(Received 11 July 2016; Accepted 01 September 2016)