



Full Length Article

Agroinfiltration Transient Assay to Evaluate the Potential of ihpRNA Constructs for Lily Mottle Virus Resistance

Pin-san Xu^{1,2}, Yue Qi², Xin-lei Wang² and Zheng-yao Zhang^{2*}

¹School of life science and biotechnology, Dalian University of Technology, Dalian, China

²School of Life Science and Medicine, Dalian University of Technology, Panjin, China

*For correspondence: zhengyaozhang@dlut.edu.cn

Abstract

In order to obtain a resistant strain of Lily mottle virus (LMoV), three RNA interference (RNAi) vectors with **intron hairpin RNA structure (ihpRNA)** were constructed and transformed into *Agrobacterium* GV 3101 strain. The gene hotspots of LMoV coated protein (CP) and cylindrical inclusion (CI) were selected, and the positive and negative fragments of the target gene were inserted into both sides of the Pdk intron fragment of plant expression vector pRNAi-LIC by one-step, zero-background ligation-independent cloning (OZ-LIC) method. Antiviral analysis was performed using *Agrobacterium*-mediated transient-expression assay. RT-PCR and indirect ELISA results showed that the hairpin RNA from the reverse repeats of CI-517 bp and CI-295 bp could interfere the virus effectively, while the CP-400 bp hairpin's impact was not significant. The work illustrates that the genetically modified genome could LMoV. © 2018 Friends Science Publishers

Keywords: Lily mottle virus; ihpRNA interference; Host-pathogen interactions; Resistance detection

Introduction

Plant antiviral genetic engineering provides a new strategy for synthesizing plant resistance viruses, among which RNA-mediated resistance virus is favored by increasing number of researchers (Kusaba, 2004). Studies have shown that ihpRNA with intron as a spacer sequence shows high silencing efficiency (Wesley *et al.*, 2001) with a maximum ihpRNA silencing efficiency of 300-600 bp length stem (Helliwell and Waterhouse, 2003). Therefore, the synthesis of an efficient ihpRNA interference vector is the key technology to apply the antiviral genetic engineering. The OZ-LIC method developed by Xu *et al.* (2010) enabled to build ihpRNA interference vector in one single step. Compared with the traditional Cleavage method and Gateway technique, this method has zero-background and low-cost advantages (Yan *et al.*, 2013). Chen *et al.* (2011) used this method to obtain an efficient ihpRNA.

LMoV is one of Potyvirus. **Once LMoV infects lily**, lily shows vein clearing, leaf mottle, twisted bifurcation, flower breaking and reduced bulb yields (Cees, 2000). Conventional detoxification is difficult to remove LMoV completely within a short time. With the development of molecular biology, antiviral genetic engineering has been increasingly applied to anti-LMoV (Zhang *et al.*, 2017). Xu *et al.* (2011) constructed virus resistant RNAi vector to Lily symptomless virus (LSV) and LMoV binary virus by Gateway technology. However, synthesizing virus that shows resistance to LMoV alone has not been reported.

Potyvirus protein function studies show that CI protein is responsible for cell-to-cell movement and ATPase/RNA helicase and CP protein is involved in genome replication, aphid transmission, cell-to-cell and systemic movement and virus assembly (Urcuqui-Inchima *et al.*, 2001). In this study, the genes of these two hot-spot sequence proteins were selected, and the RNAi vector which generates RNA hairpin structure was synthesized using OZ-LIC method.

Material and Methods

Plants, Bacterial Strains, Plasmids and other Reagents

LMoV, tobacco plants, pMDTM18-CI plasmid, pMDTM18-CP plasmid, *Escherichia coli* (*E. coli*) strain DH5a and *Agrobacterium* strain GV3101 were preserved in the lab. T4 DNA polymerase, dATP, dTTP, Gel purification kit and restriction enzymes were purchased from TAKARA (DaLian, China). The pRNAi-LIC vector was obtained from Professor Yule Liu's group of Tsinghua University.

Cloning of the Target Gene

According to the whole genome sequence of LMoV-DL (GeneBank accession number: HM222521), 3681-4197 and 4305-4599 of CI gene sequence (CI517, CI295) were selected and 1-400 of CP gene sequence (CP400) was used as the target fragments.

CI517-LIC1/CI517-LIC2 were used as primers to amplify CI517 positive fragment and CI517-LIC3/CI517-LIC4 were used to amplify CI517 reverse fragment with pMDTM18-CI plasmid as template. Similarly, CI295 and CP400 positive and negative fragments were chosen for amplification. (Table 1) The PCR products were purified using Gel purification kit.

Making ihpRNA Interference Vector

Synthesis of RNAi vector using OZ-LIC method was described in Xu *et al.* (2010). Purified forward and reverse fragments were treated with T4 DNA polymerase and dATP. The pRNAi-LIC vector was digested with Sma I and treated with T4 DNA polymerase and dNTP. Prepared forward/reverse fragments and pRNAi-LIC vector were mixed and incubated. The mixture was then transformed into *E. coli* DH5a competent cells. The plasmid was extracted and verified by restriction enzyme digestion according to the restriction sites of the vector and primer.

Agroinfiltration

RNAi vectors were transformed into *Agrobacterium* strain GV3101 competent cells by freeze-thaw method and plated in YEB liquid medium. The plasmids were extracted from *A. tumefaciens* for digestion. Leaves of tobacco plants were infiltrated by *Agrobacterium* engineering bacteria. For comparison, plants were agroinfiltrated with pRNAi-LIC vector. After 4 days, the agroinfiltrated leaves were challenged by LMoV.

RT-PCR and Indirect ELISA Analysis

Total RNA was extracted after 13 day post inoculation (dpi), and then RT-PCR was performed to detect LMoV. For RT-PCR, LMoV-CP1 and LMoV-CP2 were forward and reverse primers. Indirect ELISA was performed using a method described in Xu *et al.* (2017).

Results

Amplification of the Target Fragment

The sequencing PCR product fragments were shown in Fig. 1, which were ready for the following experiments.

Restriction Enzyme Analysis of RNAi Vector

BamH I/Sac I Digestion: the pRNAi-LIC vector has three restriction sites. Three bands occurred after digestion if the target fragment is not the cleavage site. Four bands occurred if the target fragment has one cleavage site (Fig. 2).

Pst I/Kpn I or Sal I/Kpn I Digestion: 2663 bp, 2219 bp and 2429 bp fragments were produced, respectively (Fig. 2). The plasmids were identified as RNAi-CI517, RNAi-CI295 and RNAi-CP400, respectively.




Fig. 1: Amplification of target genes
M: DL2 000 DNA marker; 1,2; 3,4; 5,6: sense and anti-sense fragments of CI₅₁₇, CI₂₉₅ and CP₄₀₀




Fig. 2: Restriction enzyme analysis of RNAi-CI₅₁₇, RNAi-CI₂₉₅ and RNAi-CP₄₀₀
M: DL15000 DNA marker; 1, 3, 5: RNAi-CI₅₁₇, RNAi-CI₂₉₅ and RNAi-CP₄₀₀ digested by BamH I/Sma I;
2,4: RNAi-CI₅₁₇, RNAi-CI₂₉₅ digested by Pst I/Kpn I; 6: RNAi-CP₄₀₀ digested by Sal I/Kpn I




Fig. 3: Identification of *Agrobacterium* engineering bacteria
M: DL15000 DNA marker; 1,2: RNAi-CI₅₁₇, RNAi-CI₂₉₅ digested by Pst I/Kpn I;
3: RNAi-CP₄₀₀ digested by Sal I/Kpn I

The Effect of Transient Interference

As shown in Fig. 3, the plasmids extracted from GV3101 produced 2663 bp, 2219 bp and 2429 bp bands, respectively. The interference vector was successfully transferred into *A. tumefaciens*.

Table 1: Primers used in this work

Name	Sequence
CI ₅₁₇ -LIC1	5'-CGACGACAAGACCCTCTGCAGTTGAAGAGCGTGAAATGGT-3'(Pst I)
CI ₅₁₇ -LIC2	5'-GAGGAGAAGAGCCCTCAAAGTCGCGTAGCTGGTT-3'
CI ₅₁₇ -LIC3	5'-CCAGCACGGAACCCTCAAAGTCGCGTAGCTGGTT-3'
CI ₅₁₇ -LIC4	5'-AGAGCACACGACCCTGGTACCTTGAAGAGCGTGAAATGGT-3'(Kpn I)
CI ₂₉₅ -LIC1	5'-CGACGACAAGACCCTCTGCAGGAGTTCACCACGCAGCATC-3'(Pst I)
CI ₂₉₅ -LIC2	5'-GAGGAGAAGAGCCCTATGATATTCGTGGCCACAACAA-3'
CI ₂₉₅ -LIC3	5'-CCAGCACGGAACCCTATGATATTCGTGGCCACAACAA-3'
CI ₂₉₅ -LIC4	5'-AGAGCACACGACCCTGGTACCGAGTTCACCACGCAGCATC-3'(Kpn I)
CP ₄₀₀ -LIC1	5'-CGACGACAAGACCCTGTCGACTGGCGTGTGGCTCATGATG-3'(Sal I)
CP ₄₀₀ -LIC2	5'-GAGGAGAAGAGCCCTCCAAGTAAGGAGTGCATGT-3'
CP ₄₀₀ -LIC3	5'-CCAGCACGGAACCCTCCAAGTAAGGAGTGCATGT-3'
CP ₄₀₀ -LIC4	5'-AGAGCACACGACCCTGGTACCTGGCGTGTGGCTCATGATG-3'(Kpn I)
LMoV-CP1	5'-TGGGCACCTTGTGAATAACA-3'
LMoV-CP2	5'-TGCTGTATGCCTTCCGTGT-3'

The RNAi-CI517 and RNAi-CI295 treatment groups were detected by RT-PCR without specific band amplification (Fig. 4), and compared with the empty vector control, the RNAi-CP400 treatment group, and leaf without chlorosis and shrinkage (Fig. 5). Meanwhile, indirect ELISA results (Table 2) showed similar conclusion. Although the symptoms were lighter than the control group in the RNAi-CP400 treatment group, the virus test results were positive. Therefore, we concluded that RNAi-CI517 and RNAi-CI295 showed good antiviral effect, while RNAi-CP400 demonstrated no obvious antiviral effect.

Discussion


Gene sequences derived from plant virus have been introduced into a wide variety of plant species to produce transgenic plants for protecting against virus infection (Sherman *et al.*, 2015). In this study, the RNAi vector containing ihpRNA was constructed using OZ-LIC method, and the feasibility and the effectiveness were verified. This method only requires four LIC linkers and a two-step polymerase treatment. With incubation overnight at 4°C, the connection efficiency was greatly enhanced. In order to investigate the effect of hairpin RNA derived from the reverse repeats of the pathogen, the invading virus by the transient expression method was mediated by *A. tumefaciens* (Ding *et al.*, 2011). The results showed that RNAi-CI517 and RNAi-CI295 showed significant interference effects, and the RNAi-CP400 had no obvious resistance to infection. The RNAi vectors' different antiviral effects depends on the length and the base sequence of stem and loop of ihpRNA, which can change the free energy of the secondary structure of the ihpRNA and as a consequence, affect its stability and silencing efficiency (Hirai *et al.*, 2007).

This work is novel in the field of LMoV disease resistance research. RNAi-CI517 and RNAi-CI295 show high silence efficiency, which can be transformed into the lily plant as the resistance strains. Meanwhile, dsRNA in vitro through synthesizing prokaryotic expression vector with ihpRNA are obtained, and plants


Table 2: Resistance effects of tobacco leaves with instant interfered of LMoV by indirect ELISA

Treatment	RNAi-CI ₅₁₇	RNAi-CI ₂₉₅	RNAi-CP ₄₀₀	pRNAi-LIC	Negative control	Blank
OD ₄₅₀	0.328	0.404	0.743	1.035	0.261	0.081
P/N	1.37	1.79	3.68	5.30	/	/
-/+	—	—	+	+	/	/

Note: P/N < 2 is negative; P/N > 2 is positive

**Fig. 4:** PCR detection of LMoV

M: DL2000 DNA marker; 1,2; 3,4; 5,6: Inoculate LMoV after agroinfiltration with RNAi-CI₅₁₇; RNAi-CI₂₉₅; RNAi-CP₄₀₀; 7,8: Vector alone; 9: Healthy plant

**Fig. 5:** Symptoms of tobacco leaves with instant interfered of LMoV

were imparted with anti-viral ability once the dsRNA enters into plants via the absorption of leaf epidermal cells (Tenllado *et al.*, 2004).

Acknowledgements

The authors are grateful for the financial support from National Natural Science Foundation of China (81503177) and they would like to thank the editors in The 4rd International Conference on Agricultural and Biological Sciences 2018 for editing our manuscript.

References

- Cees, J.A., 2000. Control of aphid-borne Lily symptomless virus and Lily mottle virus in Lilium in the Netherlands. *Vir. Res.*, 71: 23–32
- Chen, Y., W.T. Shen and P. Yan, 2011. Fast construction of dsRNA prokaryotic expression vector of Papaya ringspot virus CP gene. *J. Trop. Org.*, 2: 113–116
- Ding, Y.J., Z.S. Zhou and Y.F. Dong, 2011. Construction of the RNAi vector of GVA MP gene. *Chin. Agric. Sci. Bull.*, 27: 256–259
- Helliwell, C. and P. Waterhouse, 2003. Constructs and methods for high-throughput gene silencing in plants. *Methods*, 30: 289–295
- Hirai, S., S. Oka, E. Adachi and H. Kodama, 2007. The effects of spacer sequences on silencing efficiency of plant RNAi vectors. *Plant Cell Rep.*, 26: 651–659
- Kusaba, M., 2004. RNA interference in crop plants. *Curr. Opin. Biotech.*, 15: 139–143
- Sherman, J.H., S. Choudhuri and J.L. Vicini, 2015. RNAi technologies in agricultural biotechnology. *Regul. Toxicol. Pharm.*, 73: 671–680
- Tenllado, F., C. Llave and J.R. Díaz-Ruíz, 2004. RNA interference as a new biotechnological tool for the control of virus diseases in plants. *Vir. Res.*, 102: 85–96
- Urcuqui-Inchima, S., A.L. Haenni and F. Bernardi, 2001. Potyvirus proteins: a wealth of functions. *Vir. Res.*, 74: 157–175
- Wesley, S.V., C.A. Helliwell, N.A. Smith, M.B. Wang, D.T. Rouse, Q. Liu, P.S. Gooding, S.P. Singh, D. Abbott, P.A. Stoutjesdijk, S.P. Robinson, A.P. Gleave, A.G. Green and P.M. Waterhouse, 2001. Construct design for efficient, effective and high-throughput gene silencing in plants. *Plant J.*, 27: 581–590
- Xu, G.Y., N. Sui and Y. Tang, 2010. One-step, zero-background ligation-independent cloning intron-containing hairpin RNA constructs for RNAi in plants. *New Phytol.*, 187: 240–250
- Xu, P.S., J.F. Bai and J.W. Liu, 2011. Construction of LSV and LMoV binary virus resistant RNAi vector using Gateway technology. *Chin. Agric. Sci. Bull.*, 27: 144–147
- Xu, P.S., C.W. Zhao and Z.Y. Zhang, 2017. Identification of an RNA silencing suppressor encoded by lily symptomless virus. *Int. J. Agric. Biol.*, 19: 1259–1264
- Yan, P., W.T. Shen and X.Y. Li, 2013. Progress in construction of hpRNA vector for plant RNAi. *Biotechnol. Bull.*, 9: 7–12
- Zhang, Z.Y., X.Y. Xia, J. Song and P.S. Xu, 2017. Accumulation of Viral Coat Protein in Chloroplasts of Lily Leaves Infected with Lily Mottled Virus. *Int. J. Agric. Biol.*, 19: 1265–1269

(Received 02 June 2018; Accepted 04 August 2018)

LB PRO