



Full Length Article

Efficacy of *Bacillus thuringiensis israelensis* (Bti) Dunks in Different Qualities of Water Positive against Larvae of *Aedes aegypti* (L.) (Diptera: Culicidae) under Laboratory Conditions

Shabab Nasir^{1*}, Hina¹, Iram Nasir², Iqra Yousaf¹ and Mustapha Debboun³

¹Department of Zoology, Government College University, Faisalabad 38040, Pakistan

²Department of Statistics, Government College University, Faisalabad 38040, Pakistan

³Harris County Public Health & Environmental Services, 3330 Old Spanish Trail, Bldg. D. Houston, TX 77021

*For correspondence: flourenceshabab@yahoo.com

Abstract

Mosquitoes can breed in such varieties of water grades as fresh, turbid, foul and sewage water. Different water grades are found, on which mosquitoes can breed, due to environmental pollution caused by industry, inappropriate sewage system and lack of awareness. Therefore, the present studies demonstrated the efficacy of *Bacillus thuringiensis israelensis* (Bti) dunks against *Aede aegypti* larvae in different water grades under lab conditions. Different concentrations of Bti dunks (10, 20, 50 and 100 ppm) were used in six water grades viz., rich organic, foul, sewage, industrial, turbid and useable waters against *Ae. aegypti* larvae with control treatments in each experiment. Mortality of larvae was checked after 2, 4, 6, 12, 24 and 48 h. The mortality was highest (89%) in useable water followed by foul water (88%) after 24 h while low mortality (62% and 55%) was noted in turbid water and rich organic water due to the presence of organic matter, pollution and mud in water. Highest LC₅₀ (6.08 ppm) was observed in rich organic water while lowest LC₅₀ (3.63 ppm) was observed in sewage water after 48 h. It is concluded that use of Bti is an effective method to combat problem of mosquito borne diseases or mosquitoes themselves in all types of water but more effective in clear water. © 2018 Friends Science Publishers

Keywords: *Aedes* larvae; *Bacillus thuringiensis israelensis*; Water grades; Mortality

Introduction

Mosquitoes are vectors of the world's most hazardous pathogens of diseases such as filariasis, yellow fever, malaria, encephalitis, chikungunya, Zika virus and dengue (Reinert *et al.*, 2004; Radhika *et al.*, 2011). Dengue fever is an emerging health issue in Pakistan presently. *Aedes albopictus* and *Aedes aegypti* mosquitoes are the vectors of dengue fever. This disease emerged as an endemic all over the country (Pakistan) during 2011 particularly in Punjab province where >20,000 confirmed cases were found with 207 deaths in the city of Lahore only (Anonymous, 2013). Major reasons for outbreak of this disease were overloaded cities, inadequate water management and poor sanitary system. Prevention is the only way to avoid dengue fever; both at social and managerial levels (Junwei *et al.*, 2006). Mosquito population has been reduced through chemicals or source reduction or biological methods. The use of chemicals and source reduction cause severe burden on the ecosystem and the environment, the application of *Bacillus thuringiensis israelensis* (biological method) is an eco-friendly for the management of its juvenile stages. Toxicity to non-target flora and fauna was remarkably low (Lacey and Siegel, 2000; Lacey, 2007).

Mosquitoes can breed in a variety of habitats like polluted waters with organic and/or inorganic matter, small containers, tires and tree holes, roadside and irrigation drains, grasslands, woodland pools, tidal waters, salt marshes, dumped pools, bird baths, roofs, clogged gutters, etc. nevertheless stagnant water is prerequisite for successful breeding (Lacey and Siegel, 2000).

A number of biotic and abiotic factors (water temperature, sunlight, quantity of larvae, larval instar, application of larval food, and some water chemistry elements such as water pollution, dissolved oxygen, dust particles and concentration of dissolved organic matter) affect the action of *Bacillus thuringiensis* var. *israelensis* (Bti) (Becker *et al.*, 2010). Generally water temperature affects the toxic activity of Bti due to larval feeding rate. The Bti efficacy increases as a larvicide at higher temperature up to certain limit (35°C) due to high feeding rate of mosquito larvae (Majambere *et al.*, 2010). But the high temperature beyond a certain limit (>35°C) may decrease the activity of Bti due to the biodegradation of Bti (Karch *et al.*, 1992). Moreover, the developmental time of the mosquito vector shortened with increase in temperature and the possibility for retreatment of breeding places increases (Becker *et al.*, 1992). Every geographical locality attribute a specific combination

of above said factors that urge the need for testing the efficacy of *Bti* under those situations (Romi *et al.*, 1993). In Punjab, the environmental pollution exists due to industrialization. So, variety of water grades are present in different localities of Punjab. Therefore, the current study was proposed to investigate efficacy of *Bti* in various water grades under laboratory conditions.

Material and Methods

Collection of Mosquitoes

Adult mosquitoes were collected with aspirator from Faisalabad (Nasir *et al.*, 2017) during monsoon season. The collected specimens were brought on the same day to the Department of Zoology, Government College University, Faisalabad, by means of a plastic bottle (250 mL) knotted up with muslin cloth to the Department of Zoology, Government College University, Faisalabad.

Rearing of Mosquitoes

After identification with the help of keys (Darsi and Pradhan, 1990), the mosquitoes were reared (Ahmad *et al.*, 2017) under laboratory conditions; $25\pm 2^\circ\text{C}$ and 75% RH in rearing cages. Female mosquitoes (Nasir *et al.*, 2015) were fed with blood meals from rat and males were fed on 10% sucrose solution. The larvae were shifted in plastic trays (30x15x6 cm) after hatching of eggs and used for bioassays (20 larvae in 250 mL of water).

Collection of Water Samples

Different water grades including water with rich organic matter (from water channel in a park), tap water (from tap), turbid water (from water channel), foul water (from a small deserted pond along bank of water channel), sewage water (from sewage pond) and industrial water (from the pond near an industry) were collected and their physio-chemical properties (Table 1) were studied in the laboratory before experiment. Mosquito larvae were found in all waters except industrial water.

Test Material for Larval Bioassays

The test strains of bacterial larvicide *Bti* dunks/briquets by Summit, USA by 7000 ITU (International Toxic Unit/mg) (Fansiri *et al.*, 2006) were tested for their efficacy in different water grades against *Ae. aegypti* larvae. *Bti* EPA Registration No is 6218-47 and EPA Est. No is 6218-MD-2 was used. The net weight is 0.458 oz. per Briquet/Dunk or 13 grams per Briquet/Dunk. Different concentrations (10, 20, 50, 100 ppm) of *Bti* dunks were prepared in distilled water and used accordingly against mosquito larvae (Jahan *et al.*, 2011).

Bioassay

For each treatment, twenty larvae were taken into glass

beakers having 250 mL of water. These larvae were acclimatized for 1 h before adding *Bti*. Each experiment was replicated three times with a control (containing only water of respective grade with no *Bti*) under lab conditions ($25\pm 1^\circ\text{C}$ temperature & $70\pm 5\%$ R.H.) Dead larvae were counted after 2, 4, 6, 12, 24 and 48 h. Death of mosquito larvae was determined as a complete lack of movement, even with gentle prodding with a probe (Fansiri *et al.*, 2006).

Statistical Analysis

Abbott's Formula was used to calculate the corrected mortality percentage. Then data was examined by using statistical method, software Design Expert 9 and Minitab 15. Probit Analysis was used to evaluate the LC_{50} , LT_{50} , slope, standard error; chi square and probability.

Results

Hundred percent (100%) mortality occurred in industrial water during acclimatizing stage, so only five types of water are included. No effect of *Bti* was seen during first 2 h except in foul water (20%) and maximum mortality (85%) was noticed after maximum time interval (48 h) with 10 ppm. Less mortality was observed with low concentrations (10 and 20 ppm) than high concentrations (50 and 100 ppm). So, 100% mortality was observed with 100 ppm in almost all waters in case of 2nd instar larvae. Higher mortality was observed in 2nd instar larvae than 3rd instar larvae in almost all waters. Less than 6% mortality was also seen in control treatments. Relatively low mortality was seen in rich organic water than other water grades and maximum mortality was seen in sewage water.

In concentration response experiments where water grades were different and larvae were checked at 2, 4, 6, 12, 24 and 48 h, respectively after the start of experiment, the mortality was observed in both instars (2nd and 3rd) at all dosages including control treatment. In different water grades the lethal concentration of *Bti* to kill 50% population (LC_{50}) of *Aedes* larvae was calculated in laboratory conditions as shown in Table 2, 3 and 4. The highest LC_{50} value (34.83 ppm) with (0.034) p-values after 2 h and the lowest (6.08 ppm) after 48 h in rich organic water for 3rd instar larvae were almost double from the values of sewage water as shown in Table 2 and 4.

In our experiment *Bti* showed highest efficacy in sewage water. In foul water, LC_{50} value was 29.82 ppm after 2 h while the LC_{50} , chi square, and p-values were 10.83 ppm, 0.0573 and 0.97 respectively in case of 3rd instar larvae after 24 h as shown in Table 2 and 4. In sewage water 17.52 ppm (12.52-23.56) was required to kill 50% population of 3rd instar larvae after 2 h. The least LC_{50} (3.63 ppm) was noted after 48 h in sewage water that is almost three times less than turbid and useable water. The highest value of LC_{50} (29.31 and 21.81 ppm) was noted after 2 and 12 h in useable water.

Table 1: Physico-chemical status of water in the treated and control trials

Water grades	pH	DO (%)	DO (ppm)	TDS (ppm)	EC (m bar)
Rich organic water	7.84	8.32	0.58	2550	977
	8.82	11	0.84	2358	975
	8.51	23	1.61	1840	969
Foul Water	8.64	1.40	1.48	3189	976
	8.74	19.6	1.38	2400	989
	8.33	9.2	1.02	2728	970
	8.54	7.10	0.58	1659	976
Sewage water	8.50	7.05	0.09	2006	960
	8.16	11.8	0.85	2000	969
	7.80	11.6	0.81	1557	976
Useable Water	7.50	23	1.87	1793	975
	8.00	19.5	1.24	1670	969
	8.15	12.45	0.94	317	976
Turbid Water	5.70	26	1.88	240	975
	8.54	22	1.25	230	970
	0.90	72	4.13	80.55	976
Industrial water	0.80	96	3.70	41.25	980
	0.65	87	3.65	69.27	956

Table 2: Susceptibility of laboratory reared *Aedes aegypti* larvae to *Bacillus thuringiensis israelensis* in different water grades after 2 and 4 h

Water grades	Larval instar	2-h lethal concentration (ppm)		4-h lethal concentration (ppm)	
		LC ₅₀	P-value	LC ₅₀	P-value
Rich organic water	2 nd instar	29.08(22.80-31.72)	0.012	22.99(19.32-25.28)	0.006
	3 rd instar	34.83(28.41-36.22)	0.034	27.63(23.90-31.60)	0.147
Foul water	2 nd instar	26.01(23.03-29.25)	0.004	19.46(16.01-21.01)	0.03
	3 rd instar	29.82(27.61-34.77)	0.23	24.09(17.48-26.86)	0.023
Sewage water	2 nd instar	12.01(10.07-14.20)	0.013	9.98(7.59-12.05)	0.023
	3 rd instar	17.52(12.52-23.56)	0.459	14.90(9.99-19.09)	0.238
Useable water	2 nd instar	25.01(22.64-28.87)	0.031	24.22(22.09-28.05)	0.003
	3 rd instar	29.31(23.07-34.55)	0.012	28.89(23.69-37.15)	0.039
Turbid water	2 nd instar	20.01(18.01-23.35)	0.013	12.02(2.89-5.05)	0.003
	3 rd instar	24.25(17.09-28.64)	0.423	15.10(12.22-17.15)	0.091

Table 3: Susceptibility of laboratory reared *Aedes aegypti* larvae to *Bacillus thuringiensis israelensis* in different water grades after 6 and 12 h

Water grades	Larval instar	6-h lethal concentration (ppm)		12-h lethal concentration (ppm)	
		LC ₅₀	P-value	LC ₅₀	P-value
Rich organic water	2 nd instar	16.08(12.28-18.72)	0.013	13.99(11.34-15.28)	0.003
	3 rd instar	21.93(18.51-24.83)	0.869	18.65(15.93-21.72)	0.45
Foul water	2 nd instar	15.09(12.03-19.25)	0.004	11.46(9.01-13.61)	0.039
	3 rd instar	19.43(17.72-25.71)	0.006	15.66(10.93-19.32)	0.209
Sewage water	2 nd instar	5.99(6.07-9.20)	0.014	3.99(4.59-7.05)	0.028
	3 rd instar	9.95(5.16-13.05)	0.493	6.48(4.71-9.60)	0.136
Useable water	2 nd instar	19.21(18.04-24.87)	0.003	16.92(15.09-18.65)	0.003
	3 rd instar	23.36(18.72-27.72)	0.002	21.81(16.90-27.82)	0.003
Turbid water	2 nd instar	8.51(6.01-10.35)	0.014	5.02(3.89-6.05)	0.003
	3 rd instar	11.15(8.63-15.85)	0.297	8.90(5.80-11.14)	0.017

Table 4 showed that lowest LC₅₀ value of Bti was noted in sewage water against 2nd instar (3.31 ppm) and 3rd instar (5.20 ppm) larvae while its highest values (10.21 and 15.42 ppm) were noted in useable water, that were almost triple as compared to sewage water. After 48 h, highest LC₅₀ value was noted in turbid water as shown in Table 4.

Discussion

Bacillus thuringiensis israelensis is safe and effective biocontrol agent used widely to control *Culex*, *Anopheles*

and *Aedes* mosquitoes for the last twenty years (Lacey and Siegel, 2000; Mittal, 2003; Junwei *et al.*, 2006; Lacey, 2007). The situation of Dengue fever and dengue hemorrhagic fever has become worst in Pakistan and *Aedes* is responsible for these endemic diseases due to unplanned urbanization and pollution. In Punjab, the climatic conditions fluctuate whole the year. The changes in temperature and wide range of water grades may have a significant effect on the life cycle of mosquitoes (Tun-lin *et al.*, 2000) and biological control agents.

Table 4: Susceptibility of laboratory reared *Aedes aegypti* larvae to *Bacillus thuringiensis israelensis* in different water grades after 24 and 48 h

Water grades	Larval instar	24-h lethal concentration (ppm)		48-h lethal concentration (ppm)	
		LC ₅₀	P-value	LC ₅₀	P-value
Rich organic water	2 nd instar	8.08 (6.28-9.72)	0.012	3.99 (3.34-6.28)	0.007
	3 rd instar	11.17 (8.47-14.60)	0.869	6.08 (2.94-11.13)	0.397
Foul water	2 nd instar	7.09 (6.03-9.25)	0.003	3.46 (2.01-5.61)	0.04
	3 rd instar	10.83 (8.07-15.08)	0.972	5.21 (3.01-9.71)	3.01
Sewage water	2 nd instar	3.31 (2.07-5.20)	0.015	1.99 (1.59-2.05)	0.038
	3 rd instar	5.20 (2.38-7.59)	0.012	3.63 (1.25-7.19)	0.003
Useable water	2 nd instar	10.21 (8.64-14.87)	0.003	6.02 (5.99-8.65)	0.002
	3 rd instar	15.42 (13.93-19.54)	0.453	8.12 (7.67-10.94)	0.031
Turbid water	2 nd instar	7.51 (4.01-8.35)	0.012	6.92 (4.89-8.05)	0.003
	3 rd instar	9.63 (7.21-11.91)	0.001	8.38 (5.98-11.99)	0.021

Thus, the efficacy of *Bti* was checked in different water grades against *Ae. aegypti* larvae in the current study. In this research work, different concentrations (10, 20, 50, 100 ppm) of *Bti* dunks were prepared as done by previous workers (Abdalmagid *et al.*, 2012, Jahan and Shahid, 2012). Abdalmagid *et al.* (2012) checked the efficacy of *Bti* dunks in field water and studied the physico-chemical properties of water. They concluded that these properties have no impact on the efficacy of *Bti* ($P > 0.05$). Mulla (1990) reported high mortality rate in 1st instar larvae during handling. Due to this reason, 2nd and 3rd instar larvae were used in the present experiments. High mortality rate in case of early instar larvae (2nd) and less mortality in case of late instar larvae (3rd) was also concluded by Rodrigues *et al.* (1999). Ramathilaga *et al.* (2012) studied the impact of *Bti* against 3rd instar larvae of *Ae. aegypti* as was recorded in the present study against 2nd & 3rd instar larvae. In the present study, 63% and 79% mortality was recorded for 50 and 100 ppm of *Bti* respectively after 24 h in useable water while Ramathilaga *et al.* (2012) recorded (16%) mortality at 1mg concentration of *Bti* for 24 h treatment in tap water. Haung *et al.* (1993) recorded 52.1, 69.5 and 78.2% mortality after 12, 24 and 48 h respectively in 0.10 ppm against *Ae. aegypti* larvae while 97.1, 97.1 and 97.1% mortality after 12, 24 and 48 h in 0.20 ppm. Gbehou *et al.* (2010) compared the efficacy of *Bti* on mosquitoes (*Aedes*, *Culex* and *Anopheles*) and observed 40, 80 & 100% mortality after 2, 4 and 6 h against *Aedes*. Many other factors (genera susceptibility, feeding behavior of larvae, instar susceptibility to *Bti*, suspended organic matter, water temperature, larval density, water depth) also affect the efficacy of *Bti* against mosquitoes (Boisvert, 2005). Some of these factors like organic, inorganic, muddy, food and floating particles decreased the efficacy of *Bti* due to adsorption of *Bti* onto suspended particles followed by a slow sedimentation (Margalit and Bobroglo, 1984; Margalit *et al.*, 1985). Hence, fewer toxic crystals are ingested by larvae due to suspended particles. So, higher concentration of *Bti* will be needed for larval control in case of high density of suspended particles and pollution (Ohana *et al.*, 1987; Mulla, 1990). The interruption in the efficacy of *Bti* was found to be caused by bacterial adsorption to soil particles, but the inactivation

could be inverted by washing the mud away (WHO, 2004). Due to these reasons, mean value of LC₅₀ was higher in rich organic water (34.83 ppm) and turbid water (24.25 ppm) while low in sewage water (17.52 ppm) after 2 h in the present study. Mean LC₅₀ values were 5.20 and 10.83 ppm after 24 h for sewage water (full of organic matter) and foul water, respectively. The results of the present study revealed the higher mortality in sewage water and useable water because these were free of any particles while least was observed in rich organic water due to suspended particles. *Bti* formulation along with these factors affects its efficacy. *Bti* formulations originated as a water dispersible powder (WDP) formulation using fly ash (FA) as a carrier material was developed and was found effective against the larval stages of major mosquito vector species (Tamilselvan *et al.*, 2017).

Conclusion

Use of *Bti* in different water grades is an effective method to control larvae of dengue mosquito, *Aedes aegypti*.

References

- Abdalmagid, M.A., S.M. Elkhalfa, A.B. Ismail, S.M. Ismail, A.H. Abdalrahman, A.E. Jamal, M. Brair and I.H. Elnaeim, 2012. The mosquito dunk (*Bacillus thuringiensis israelensis*) as a control agent against mosquito larvae in Khartoum State, Sudan. *Sudan J. Public Health*, 7: 51–55
- Ahmad, T., S. Nasir, I. Nasir, T. Nawaz, A. Rafique and I. Yousaf, 2017. Response surface modeling for West Nile viral encephalitis mosquito control experiments. *Pak. Vet. J.*, 37: 465–469
- Anonymous, 2013. *Evaluation Report on Project: Prevention and Control Program of Epidemics in Punjab*. P&D department, Civil Secretariat, Lahore–Punjab, Pakistan
- Becker, N., D. Petric, M. Zgomba, C. Boase, M. Madon, C. Dahl and A. Kaiser, 2010. *Mosquito and their Control*. Heidelberg, Springer, Germany
- Becker, N., M. Zgomba, M. Ludwig, D. Petric and F. Rettich, 1992. Factors influencing the activity of *Bacillus thuringiensis* var. *israelensis* treatments. *J. Am. Mosq. Control Assoc.*, 8: 285–289
- Boisvert, M., 2005. Utilization of *Bacillus thuringiensis* var. *israelensis* (*Bti*)-Based Formulations for the Biological Control of Mosquitoes in Canada. 6th Pacific-Rim Conf. on Biotechnol. of *Bacillus thuringiensis* and its Environmental Impact. Victoria BC

- Darsi, Jr.R.F. and S.P. Pradhan, 1990. The mos-quitoes of Nepal: Their identification, distribution and biology. *Mosq. Syst.*, 22: 69–130
- Fansiri, T., U. Thavara, A. Tawatsin, S. Krasaesub and R. Sithiprasasna, 2006. Laboratory and semi-field evaluation of mosquito dunks against *Aedes aegypti* and *Aedes albopictus* larvae (diptera: culicidae). *Southeast Asian J. Trop. Med. Public Health*, 37: 62–66
- Gbehou, N.A., H.S. Christophe and L. Yilian, 2010. Effect of *Bacillus thuringiensis* var. *israelensis* (H-14) on *Culex*, *Aedes* and *Anopheles* larvae (Cotonou; Benin). *Stem Cell*, 1: 60–67
- Haug, R.N., I.P. LO, C. Ho, J.S. Haug and E.L. Hsu, 1993. Effectiveness of two forms of *Bacillus thuringiensis* var. *israelensis* on mosquitoes larvae. *Chin. J. Entomol.*, 13: 177–185
- Jahan, N., J. Razaq and A. Jan, 2011. Laboratory evaluation of chitin synthesis inhibitors (Diflubenzuron and buprofezin) against *Aedes aegypti* larvae from Lahore, Pakistan. *Pak. J. Zool.*, 43: 1079–1084
- Jahan, N. and A. Shahid, 2012. Evaluation of resistance against *Bacillus thuringiensis israelensis* WDG in dengue vector from Lahore, Pakistan. *Pak. J. Zool.*, 44: 995
- Junwei, Z., Z. Xiaopeng, T.L. Yanma, L. Ting, Q. Kuen, H. Yuhua, X. Suqin and T. Brad, 2006. Adult repellency and larvicidal activity of five plant essential oils against mosquitoes. *J. Am. Mosq. Control Assoc.*, 3: 515–522
- Karch, S., N. Asidi, M. Manzambi and J.J. Salaun, 1992. Efficacy of *Bacillus sphaericus* against the malaria vector *Anopheles gambiae* and other mosquitoes in swamps and rice fields in Zaire. *J. Am. Mosq. Control Assoc.*, 8: 376–380
- Lacey, L.A. and J.P. Siegel, 2000. Safety and ecotoxicology of entomopathogenic bacteria. In: *Entomopathogenic Bacteria: from Laboratory to Field Application*, pp: 253–273. Charles, J.F., A. Delecluse and C. Nielsen- LeRoux (eds). Kluwer Academic Publishers, Dordrecht, The Netherlands
- Lacey, L.A., 2007. *Bacillus thuringiensis* serovariety *israelensis* and *Bacillus sphaericus* for mosquito control. *J. Am. Mosq. Control Assoc.*, 23: 133–163
- Majambere, S., M. Pinder, U. Fillinger, D. Ameh, D.J. Conway and C. Green, 2010. Is mosquito larval source management appropriate for reducing malaria in areas of extensive flooding in the Gambia? A cross-over intervention trial. *Am. J. Trop. Med. Hyg.*, 82: 176–84
- Margalit, J. and H. Bobroglo, 1984. The effect of organic materials and solids in water on the persistence of *Bacillus thuringiensis* var. *israelensis*. *Z. Angew Entomologie-J. App. Entomol.*, 97: 516–520
- Margalit, J.C., H. Pascar-gluzman, Z. Bobroglo, Barak and L. Lahkim-tsror, 1985. Biocontrol of mosquitoes in Israel. In: *Integrated Mosquito Control Methodologies*, pp: 361–374. Laird, M. and J.W. Miles (eds) San-93 Diego, Academic Press, USA
- Mittal, P.K., 2003. Biolarvicides in vector control: Challenges and prospects. *J. Vector Borne Dis.*, 40: 20–32
- Mulla, M.S., 1990. Activity, field efficacy and use of *Bacillus thuringiensis israelensis* against mosquitoes. In: *Bacterial control of mosquitoes and black flies: Biochemistry, genetics and applications of Bacillus thuringiensis israelensis and Bacillus sphaericus*, pp. 134–160. Barjac, H. and D.J. Sutherland (eds.). Rutgers University Press, New Brunswick, New Jersey
- Nasir, S., M. Batool, S.M. Hussain, I. Nasir, F. Hafeez and M. Debboun, 2015. Bioactivity of oils from medicinal plants against immature stages of dengue mosquito *Aedes aegypti* (Diptera: Culicidae). *Int. J. Agric. Biol.*, 17: 843–847
- Nasir, S., I. Nasir and I. Yousaf, 2017. Medicinal plant extracts: control strategy against dengue mosquitoes. *Bang. J. Bot.*, 46: 27–33
- Ohana, B., J. Margalit and Z. Barak, 1987. Fate of *Bacillus thuringiensis* subsp. *israelensis* under Simulated Field Conditions. *App. Environ. Microbiol.*, 53: 828–831
- Radhika, D., A. Ramathilaga and C.S. Prabu, 2011. Evaluation of larvicidal activity of soil microbial isolates (*Bacillus* and *Acinetobacter* sp.) against *Aedes aegypti* (Diptera: Culicidae), the vector of Chikungunya and Dengue. *Proc. Int. Acad. Ecol. Environ. Sci.*, 1: 169–178
- Ramathilaga, A., A.G. Murugesan, C. Sathesh and Prabu, 2012. Biolarvicidal activity of *Peainbacillus macerans* and *Bacillus subtilis* isolated from the dead larvae against *Aedes aegypti* – Vector for Chikungunya. *Proc. Int. Acad. Ecol. Environ. Sci.*, 2: 90–95
- Reinert, S.F., R.E. Harbaach and I.J. Kitching, 2004. Phylogeny and classification of *Aedini* (Diptera: Culicidae), based on morphological character of all life stages. *Zool. J. Linn. Soc.*, 142: 289–368
- Rodrigues, I.B., W.P. Tadei and J.S. Dias, 1999. Larvicidal activity of *Bacillus sphaericus* 2362 against *Anopheles nuneztovari*, *Anopheles darlingi* and *Anopheles braziliensis* (Diptera: culicidae). *Rev. Inst. Med. Trop. S. Paulo*, 41: 2
- Romi, R., B. Ravoniharimelina, M. Ramiakajato and G. Majori, 1993. Field trials of *Bacillus thuringiensis* H-14 and *Bacillus sphaericus* (strain 2362) formulations against *Anopheles arabiensis* in the central highlands of Madagascar. *J. Am. Mosq. Control Assoc.*, 9: 325–329
- Tamilselvan, S., A.M. Manonmani and P. Jambulingam, 2017. Fly ash-based water dispersible powder formulation of *Bacillus thuringiensis* var. *israelensis*: Development & laboratory evaluation against mosquito immatures. *Ind. J. Med. Res.*, 146: 714–721
- Tun-Lin, W., T.R. Burkot and B.H. Kay, 2000. Effects of temperature and larval diet on development rates and survival of the dengue vector *Aedes aegypti* in north Queensland Australia. *Med. Vet. Entomol.*, 14: 31–37
- WHO, 2004. *Report of the Seventh WHOPES Working Group Meeting*. WHO/ CDS/ WHOPES/ 8: 3–28

(Received 12 April 2018; Accepted 28 June 2018)