



## Full Length Article

# Genome-wide Identification, Classification and Expression of Flavonol Synthase from *Nelumbo nucifera* in Defense against Various Stresses

Chen Dong<sup>1\*</sup>, Xingfei Zheng<sup>2</sup>, Yannan Shi<sup>1</sup>, Shuang Chen<sup>1</sup>, Die Liu<sup>1</sup>, Peng Lin<sup>1</sup> and Zhongli Hu<sup>2\*</sup>

<sup>1</sup>College of Biological Engineering, Henan University of Technology, Zhengzhou 450001, China

<sup>2</sup>State Key Laboratory of Hybrid Rice, Lotus Engineering Research Center of Hubei Province, College of Life Science, Wuhan University, Wuhan 430072, China

\*For correspondence: chen.dong@haut.edu.cn; huzhongli@whu.edu.cn

## Abstract

Four full-length cDNA of flavonol synthase (*NnFLS*) named as *NnFLS1*, *NnFLS2*, *NnFLS3* and *NnFLS4* was identified in *Nelumbo nucifera*, encoding proteins with sizes between 331 and 362 amino acids. His-x-Asp-x<sub>n</sub>-His (HxDx<sub>n</sub>H) domain was found in putative *NnFLS1*, *NnFLS2* and *NnFLS3*, while the first His was replaced by Arg in *NnFLS4*. Arg-x-Ser (RxS) domain was highly conserved in all *NnFLSs*. Four *NnFLSs* genes comprised of 3 exons and 2 introns, and the different exon-intron junctions of *NnFLS4* was investigated. Additionally, phylogenetic analysis indicated *NnFLS2* was grouped together with dicotyledons, and other three *NnFLSs* were classed into monocotyledon. The second structures and homology models indicated that four *NnFLSs* were highly conserved. Moreover, the highest levels of four *NnFLSs* mRNAs were detected in embryo with the lowest levels in the stems. Furthermore, the levels of four *NnFLSs* mRNA were significantly elevated in response to various stresses including chilling, ultraviolet-B radiation (UV-B) and short-time mechanical wounding. These results are likely to serve as fundamental research for future evolutionary and functional characterization studies of the *NnFLS* genes in lotus. © 2019 Friends Science Publishers

**Keywords:** Flavonol synthase; Flower opening; Gene expression; *Nelumbo nucifera*

## Introduction

Flavonoids are well-known types of polyphenols in high plants, exhibiting antioxidant, anti-proliferative and anti-inflammatory functions in defending against cancer and cardiovascular disease (Kaur *et al.*, 2008). Considering as the most widely distributed flavonoids, Flavonols play imperative roles in plant physiology, growth and development including protection from UV damage or change of flower color (Stracke *et al.*, 2010a). Additionally, flavonols were involved in the ethylene-signaling pathway (Lewis *et al.*, 2011).

Three common types of flavonols including myricetin, quercetin and kaempferol are generally converted from the dihydromyricetin, dihydroquercetin or dihydrokaempferol by flavonol synthase (FLS), which is usually classified as bifunctional dioxygenase and defined as having hydroxylation and desaturation activity (Akita *et al.*, 2018). This process requires coordination of gene expression in the flavonoid pathway including chalcone synthase (CHS), flavanone 3 $\beta$ -hydroxylase (F3H), anthocyanidin synthase (ANS) and FLS. All of these genes belong to the second largest family of higher plants known as 2-oxoglutarate-dependent dioxygenase (2-ODD), which catalyse the oxidation reaction using ferrous iron or 2-oxoglutarate as cofactor (Akita *et al.*,

2018). *FLS* cDNAs had been isolated from high plants including *Zea mays*, *Oryza sativa*, *Arabidopsis thaliana*, *Petroselinum crispum*, lisianthus, strawberry and soybean (Takahashi *et al.*, 2007). Studies have indicated that *FLS* was encoded by a muticopy gene family in plants, and the different copies were expressed in an organ-specific pattern (Preuß *et al.*, 2009; Ferreyra *et al.*, 2010; Kim *et al.*, 2010). Overexpression of *ANS* accumulated a mixture of flavonoids in transgenic rice (Reddy *et al.*, 2007). Healthy phytochemicals treatment also led to the production of novel flavonoids in tomato fruit (Schijlen *et al.*, 2006).

As one of the oldest plants surviving from the last ice age, sacred lotus (*Nelumbo nucifera* Gaertn) is considered as a traditional ornamental and medicinal plant in China. Its seeds have exceptional longevity, remaining viable as long as thousands of years. Each parts of sacred lotus could be used in herbal medicines for curation of cancer, depression, diarrhea, heart disease as well as insomnia. It is also well-known as healthy food rich in proteins, lipids, dietary fiber, minerals and flavonoids. Moreover, petals and stamens of lotus rich in flavonoids and alkaloids are widely used in Chinese herbal medicine.

Although flavonoids in lotus were considered as the key composition in Chinese herbal medicine, the biosynthetic synthase of flavonoids is still unveiled in *N. nucifera*.

We hypothesize that comprehensive analysis of *FLS* in *N. nucifera* (NnFLS) is thought to benefit for illustrating the mechanism of biosynthetic pathway of flavonoids in *N. nucifera*. Here, we identified four *NnFLSs* genes in lotus genome. A systematic analysis of *NnFLS* genes was performed, including their organization of intron and exon, evolutionary relationship, and expression profile in various tissues as well as stresses. The findings of this study will serve as foundation for functional characterization studies of NnFLS in lotus.

## Materials and Methods

### Plant Material

Seed of *N. nucifera* “Taikonglian-36” was collected from Lotus Engineering Research Center of Hubei Province, Wuhan University, and planted in Henan University of Technology, China. Rhizome is the storage organ of *N. nucifera*, which is cultured in soil of experimental ponds for asexual propagation. All the materials were collected and frozen in liquid nitrogen immediately.

### Identification and Bioinformatics Analysis of NnFLS

To scan NnFLS in the genome of *N. nucifera*, BLASTP searches were conducted using proteins sequences of FLS from *Arabidopsis thaliana* against the NCBI database (E value <  $1e^{-6}$ ). Another search for NnFLS was performed using the key word “flavonol synthase” in the Sacred lotus (*N. nucifera*) genome database (<http://lotus-db.wbgcas.cn/>). The putative NnFLS was further searched in the NCBI conserved domain database (<https://www.ncbi.nlm.nih.gov/Structure/bwrpsb/bwrpsb.cgi>) to confirm the presence of conserved domains of FLS. The ProtParam program (<http://web.expasy.org/protparam/>) was performed to evaluate the putative molecular weight (MW), isoelectric point (pI) and grand average of hydropathicity (GRAVY) for NnFLS. Protcomp Version 9.0 software was used to predict the Sub-cellular location of NnFLS.

### Analysis of Exon–intron Structures and Conserved Motifs

The exon–intron structures of NnFLS were analyzed by Gene Structure Display Server (GSDS; <http://gsds.cbi.pku.edu.cn/index.php>) (Hu *et al.*, 2014). Protein structures of NnFLS were predicted by SMART online tools (<http://smart.embl-heidelberg.de/>). Conserved motifs were indicated using online MEME program (<http://meme.nbcr.net/meme/cgi-bin/meme.cgi>).

### Phylogenetic Analysis

The putative protein sequences of NnFLS were compared with other FLS of high plants by Clustal W software.

Additionally, the phylogenetic relationship of FLS family in high plants was drawn using the neighbour-joining (NJ) method by the MEGA software version 4 (Tamura *et al.*, 2007).

### Secondary Structures and Three Dimensional Models of NnFLS

The secondary structures of NnFLS were predicted using the PSIPred program (Expasy tools). ANS of *A. thaliana* (PDB.1gp6\_A) was selected as the homology template from RSCB protein data bank (Wilmouth *et al.*, 2002). About fifty homology models of NnFLS were produced by the MODELLER software with the default parameters (Sanchez and Sali, 1997), which were further assessed by PROCHECK 3.5 for testing the qualities of the models (Laskowski *et al.*, 1993). The best model was selected and shown by Swiss-pdbviewer 4.1.0.

### Differential Expression of NnFLS in Tissues

For examining the level of *NnFLS* mRNAs in various tissues, experimental materials such as young leaves, stem, roots and embryo were harvested.  $\beta$ -actin (GenBank accession no. EU131153) was selected as a reference gene. The gene-specific primers for *NnFLS* (NnFLS F and NnFLS R) and  $\beta$ -actin ( $\beta$ -actin F and  $\beta$ -actin R) were synthesized using Primer Premier 5 software (Premier) (Table S1). DNA binding dye SYBR GreenI (TOYOBO) was used for detection of PCR products in Real-time PCR (Dong *et al.*, 2017, 2018). The PCR program included one cycle at 94°C for 30 s and then 30 cycles of 94°C for 15 s and 60°C for 15 s. According to the method published (Livak and Schmittgen, 2001), the fold change of *NnFLSs* was calculated using  $\beta$ -actin as a reference gene by  $2^{-\Delta\Delta Ct}$  method.

### Expression Pattern of NnFLS in Response to Stresses

To unveil the roles of *NnFLS* in the face of environmental stresses, the levels of *NnFLS* mRNAs were detected by Real-time PCR. The seedlings in a closed chamber were irradiated under 1000  $\mu\text{J}/\text{m}^2$  ultraviolet-B radiation (UV-B) and separately collected at 2 h post treatment (hpt), 4 hpt and 6 hpt and the plants placed in a dark closed chamber were used as control. For chilling stress, the seedlings were transferred to a cold (4°C) growth chamber for 2, 4 and 6 h, respectively and the control plants continued to grow at room temperature (25°C) in normal condition. In term of short-time mechanical wounding treatment, the leaves of *N. nucifera* were treated by pressure-stress with a needle puncher when the leaves had unfolded for one week. The leaves were separately collected at 2, 4 and 6 hpw, using the intact leaves as the control. The total RNAs of materials were isolated, and the levels of *NnFLS* mRNA were examined by Real-time PCR as method described above.

### Statistical Analysis

Three independent biological replicates were performed to calculate the fold change of *NnFLS* by Real-time PCR. Data were expressed as the mean ± SD from three independent biological replicates. Significance was determined based on one-way analysis of variance (ANOVA) and the least significant difference (LSD) was performed to find out differences between groups ( $p < 0.05$ ). The experiments were repeated twice each time to get reproducible data.

### Results

#### Identification and Conserved Domains of *NnFLS*


Four *NnFLSs* named as *NnFLS1* (XP\_010256287.1), *NnFLS2* (XP\_010272845.1), *NnFLS3* (XP\_010274342.1) and *NnFLS4* (XP\_019055362.1) were identified in our study, encoding the putative proteins between 331 and 362 amino acids (Table 1). The MW of different *NnFLSs* was between 37.70 KD and 41.03 KD, with pI between 5.18 and 5.85. Moreover, all the four *NnFLSs* were predicted to be located in cytoplasm (Table 1). Alignment of conserved domains and functional analysis illustrated all *NnFLSs* belonged to the 2OG-FeII\_Oxy superfamily. Three highly conserved amino acids (His, Asp and His) were detected in *NnFLS1*, *NnFLS2* and *NnFLS3*, which were involved in binding with a ferrous iron II while the first His was replaced by Arg in *NnFLS4* (Fig. 1). Moreover, two amino acids as Arg and Ser were the putative 2-oxoglutarate binding sites and highly conserved in four *NnFLSs*. The conserved region of 2-ODD superfamily was also detected in *NnFLSs* (Fig. 1).

#### Exon-intron Structures of *NnFLS*


According published DNA sequence of *NnFLS* genes in NCBI, the detailed comparisons suggested that all *NnFLSs* had 3 exons and 2 introns, and the two introns showed different sizes (Fig. 2). The lengths of exon 1 and exon 3 were various, and all the exon 2 of four *NnFLSs* was 328 bp (Table 1). The exon-intron junctions of *NnFLS* genes were consistent with the GT-AG rule, except GC-AG splicing way in intron 1 of *NnFLS4* (Fig. 2).

#### Phylogenetic Analysis of *NnFLSs*

A phylogenetic tree was constructed to investigate the evolution relationship of 2-ODD superfamily. The topology of the phylogenetic tree was an asymmetric clover shape, including FLS, F3H and ANS families. Compared with F3H, FLS family had closer relationship with ANS. Among the 2-ODD superfamily, *NnFLSs* showed higher identity with other FLS from high plants, and had lower similarity with F3H or ANS (Fig. 3). In subgroup of FLS, *NnFLS2* was belonged to dicotyledons. However, other three *NnFLSs* were grouped together with monocotyledon (Fig. 3).



**Fig. 1:** Amino acid sequence alignment and characteristics of *NnFLS*. The 2-ODD superfamily conserved region was shadowed by grey. The conserved residues binding with a ferrous iron II were represented by yellow. The putative 2-oxoglutarate binding motif consisting of Arg and Ser was described by red. Several amino acids (Phe, Phe, Lys, and Ser) involved in binding to substrate were represented by green



**Fig. 2:** Exons-intron architecture of *NnFLS* genes, CDS, intron and UTR were labeled. The exon-intron junctions of *NnFLS* genes were examined

#### Second Structures and Homology Model of *NnFLSs*



Second structures indicated that both *NnFLS1* and *NnFLS3* had 11  $\alpha$ -helix and 9  $\beta$ -strands. *NnFLS2* composed of 10  $\alpha$ -helix and 8  $\beta$ -strands, with 10  $\alpha$ -helix and 10  $\beta$ -strands detected in *NnFLS4*. The putative three-dimensional structures of *NnFLSs* were constructed using crystal structure of *A. thaliana* ANS (PDB. 1gp6\_A) as homology template, which shared high degree of homology with *NnFLSs*. Four *NnFLSs* had the similar three-dimensional structures, consisting of  $\alpha$ -helix and  $\beta$ -strands in N-terminal and C-terminal of proteins (Fig. 4).

#### Expression of *NnFLS* in Various Tissues

*NnFLS* mRNAs were examined in different tissues of *N. nucifera* including young leaves, roots, stems and embryo.

**Table 1:** identified NnFLSs and their sequence characteristics

Gene name	NCBI accession no	Exon1	Intron1	Exon2	Intron2	Exon3	ORF	Amino acids	MW (KD)	pI	GRAVY	location
NnFLS1	XP_010256287.1	462	73	328	558	364	999	332	37.82	5.18	-0.344	Cytoplasmic
NnFLS2	XP_010272845.1	556	114	328	2078	386	1008	335	38.08	5.56	-0.410	Cytoplasmic
NnFLS3	XP_010274342.1	544	1854	328	1172	439	996	331	37.70	5.41	-0.456	Cytoplasmic
NnFLS4	XP_019055362.1	557	1042	328	849	326	1089	362	41.03	5.85	-0.340	Cytoplasmic

**Fig. 3:** Phylogenetic analysis of NnFLS and other 2-ODD superfamily including ANS as well as F3H by neighbour-joining method using MEGA software version 4**Fig. 4:** The homology model of NnFLS1 (a), NnFLS2 (b), NnFLS3 (c) and NnFLS4 (d)

The expression pattern of *NnFLS2*, *NnFLS3* and *NnFLS4* was similar. The highest level of *NnFLS2*, *NnFLS3* and *NnFLS4* mRNA was detected in embryo, with the moderate level of *NnFLSs* mRNA in young leaves. Additionally, low expression of *NnFLS2*, *NnFLS3* and *NnFLS4* mRNA was


detected in roots while it was still more than stems as control (Fig. 5). However, different expression profile of *NnFLS1* was found. The most expression of *NnFLS1* was indicated in embryo and higher level of *NnFLS1* mRNA was found in roots than young leaves using stems as control (Fig. 5).

### *NnFLS* Expression in Response to Various Stresses


To find out the role of NnFLS in various stresses, Real-time PCR was used to study the expression pattern of *NnFLS* mRNA. UV-B treatment increased the expression of four *NnFLSs* mRNA at 2 hpt using Tween 20 treated seeding as control (Fig. 6). The level continued to increase at 4 hpt and reached the highest level at 6 hpt. Additionally, mRNAs level of *NnFLSs* was also induced by chilling treatment (4°C) using the seeding in room temperature as control (Fig. 7). Their expression started to increase at 2 hpt except *NnFLS3* and significantly increased at 4 hpt with the highest expression at 6 hpt. Moreover, short-time mechanical wounding augmented the expression of *NnFLS* mRNA. Real-time PCR indicated that *NnFLS* mRNA reached the highest level as soon as 2 hpt using normal seeding as control, but declined at 4 hpt and 6 hpt (Fig. 8).

### Discussion

Since FLS roles as the major enzyme involved in the flavonoid pathway, and several *FLS* genes in high plants have been studied at the chemical, genetical or enzymatic levels (Nguyen *et al.*, 2016). However, few reports are available on the identification of *FLS* genes in aquatic plant. The flavonoids are known as the most imperative medicinal component in *N. nucifera*, whereas biosynthetic pathway of flavonoids is still unclear. In this study, four *NnFLSs* were identified and studied in lotus (Table 1), which possessed all the conservative domains and active sites of FLS family, including ferrous iron II binding residues and putative 2-oxoglutarate binding residues (Fig. 1). All *NnFLS* genes contained 3 exons and 2 introns. The exon-intron junctions of *NnFLS* genes were consistent with the GT-AG rule, except intron 1 of *NnFLS4* (Fig. 2). Moreover, phylogenetic tree analysis indicated NnFLS was located in FLS group (Fig. 3). Interesting, only NnFLS2 was grouped into dicotyledons, and other three NnFLSs were classed into monocotyledon. Considering *N. nucifera* was the species between dicotyledons and monocotyledon in evolutionary relationship, NnFLS1, NnFLS3 and NnFLS4 were probably evolved from NnFLS2.




**Fig. 5:** The expression patterns of *NnFLS* mRNAs in different tissues. Fold change of *NnFLS* mRNAs in different tissues. Total RNAs from roots, young leaves, embryo, and stems were isolated. *NnFLS* mRNAs were detected using  $\beta$ -actin as a reference gene by Real-time PCR




**Fig. 6:** The mRNA level of *NnFLS* in response to UV-B treatment. The seedlings in a closed chamber were irradiated under 1000  $\mu\text{J}/\text{m}^2$  UV-B, and the plants placed in a dark closed chamber were used as control. Relative quantities of *NnFLS*s mRNA at various hours post treatment (hpt) were calculated. Different letters were used to represent significant difference at  $p < 0.05$

*NnFLS*s contained several  $\beta$ - strands, forming a hydrophobic active center known as jellyroll fold. Additionally,  $\alpha$ -helix was detected in both N-terminus and C-terminus of *NnFLS* (Fig. 4). The absence of the first  $\alpha$ -helix in *AtFLS1* could cause loss of activity (Li *et al.*, 2012) and the mutation or deletion of this region in other *AtFLS*s would cause the inactivation of expression products (Owens *et al.*, 2008). Both *FLS* and *ANS*



**Fig. 7:** The mRNA level of *NnFLS* during chilling stress. For chilling stress, the seedlings were transferred to a cold (4°C) growth chamber for 2 h, 4 h and 6 h respectively, and the control plants continued to grow at room temperature (25°C) in normal condition. The relative expression of *NnFLS*s was calculated by Real-time PCR



**Fig. 8:** The mRNA level of *NnFLS* in term of short-time mechanical wounding. The leaves of *N. nucifera* were treated by pressure-stress with a needle puncher when the leaves had unfolded for one week, using the intact leaves as the control. The relative expression of *NnFLS*s was examined by Real-time PCR

belonged to the 2-ODD superfamilies with similar sequences and functions. Previous study illustrated *FLS* shared the same substrates with *ANS in vitro*, which produced the same products.

Flavonols are normally rich in leaves, flower, branches and heartwood of *A. confusa* (Hsieh and Chang, 2010). *AtFLS1* could be stimulated by transcription factor such as MYB11, MYB12 and MYB111 in *A. thaliana*,

which result in various spatial accumulation of specific flavonol derivatives in leaves, stems and roots (Stracke *et al.*, 2010b). A similar expression profile of various *NnFLSs* was investigated in this study. Significant change of *NnFLS* mRNA was detected with the highest level in embryo, while lowest expression of *NnFLSs* was found in stems (Fig. 5).

Plants have complex mechanisms to cope with environmental factors including UV irradiation, chilling, wounding, nutrient depletion as well as plant hormones, which advanced the accumulation of flavonols in plants (Ferreira *et al.*, 2010; Lewis *et al.*, 2011). It had been reported that *FLS* in poplar was not stimulated by wounding treatment (Mellway *et al.*, 2009). We therefore concluded that *FLS* exhibited various expression patterns in different plants. Our study indicated *NnFLS* mRNAs were stimulated by mechanical wounding, chilling and UV-B treatment (Fig. 6, 7 and 8). All of these results proved that *NnFLS* played an imperative role in defense against environmental stresses.

## Conclusion

Although four *NnFLSs* were induced by environmental stresses including wounding, chilling and UV-B treatment, transgenic plants should be developed for understanding the mechanism of *NnFLS* in aquatic plant.

## Acknowledgments

This work is financially supported by National Natural Science Foundation of China (31601366), Foundation of Henan Educational Committee (16A180024) and Doctoral Scientific Research Start-up Foundation from Henan University of Technology (31400855).

## Conflicts of Interest

The authors declare that they have no conflicts of interest.

## References

- Akita, Y., S. Kitamura, R. Mikami and H. Ishizaka, 2018. Identification of functional flavonol synthase genes from fragrant wild cyclamen. *J. Plant Biochem. Biotechnol.*, 27: 147–155
- Dong, C., M.G. Yang, H.H. Wang and J.P. Mi, 2017. Identification and expression analyses of two lotus (*Nelumbo nucifera*) dehydrin genes in response to adverse temperatures, ABA and IAA treatments. *Biologia*, 72: 745–752
- Dong, C., R. Wang, X.F. Zheng, X.W. Zheng, L.F. Jin, H.J. Wang, S. Chen, Y.N. Shi, M.Q. Wang, D. Liu, Y.H. Yang and Z.L. Hu, 2018. Integration of transcriptome and proteome analyses reveal molecular mechanisms for formation of replant disease in *Nelumbo nucifera*. *RSC ADV*, 8: 32574–32587
- Ferreira, M.L.F., S. Rius, J. Emiliani, L. Pourcel, A. Feller, K. Morohashi, P. Casati and E. Grotewold, 2010. Cloning and characterization of a UV-B-inducible maize flavonol synthase. *Plant J.*, 62: 77–91
- Hsieh, C.Y. and S.T. Chang, 2010. Antioxidant activities and xanthine oxidase inhibitory effects of phenolic phytochemicals from *Acacia confusa* twigs and branches. *J. Agric. Food Chem.*, 58: 1578–1583
- Hu, B., J. Jin, A.Y. Guo, H. Zhang, J. Luo and G. Gao, 2014. GSDB 2.0: an upgraded gene feature visualization server. *Bioinformatics*, 31: 1296–1297
- Kaur, P., S. Shukla and S. Gupta, 2008. Plant flavonoid apigenin inactivates Akt to trigger apoptosis in human prostate cancer: an in vitro and in vivo study. *Carcinogenesis*, 29: 2210–2217
- Kim, B.G., E.J. Joe and J.H. Ahn, 2010. Molecular characterization of flavonol synthase from poplar and its application to the synthesis of 3-O-methylkaempferol. *Biotechnol. Lett.*, 32: 579–584
- Laskowski, R.A., M.W. MacArthur, D.S. Moss and J.M. Thornton, 1993. Procheck: a program to check the stereochemical quality of protein structures. *J. Appl. Crystallogr.*, 26: 283–291
- Lewis, D.R., M.V. Ramirez, N.D. Miller, P. Vallabhaneni, W.K. Ray, R.F. Helm, B.S. Winkel and G.K. Munday, 2011. Auxin and ethylene induce flavonol accumulation through distinct transcriptional networks. *Plant Physiol.*, 156: 144–164
- Li, C.L., Y.C. Bai, S.J. Li, H. Chen, X.Y. Han, H.X. Zhao, J.R. Shao, S. Park and Q. Wu, 2012. Cloning, Characterization, and Activity Analysis of a Flavonol Synthase Gene *PtFLS1* and Its Association with Flavonoid Content in Tartary Buckwheat. *J. Agric. Food Chem.*, 60: 5161–5168
- Livak, K.J. and T.D. Schmittgen, 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method. *Methods*, 25: 402–408
- Mellway, R.D., L.T. Tran, M.B. Prouse, M.M. Campbell and C.P. Constabel, 2009. The wound-, pathogen-, and ultraviolet B-responsive MYB134 gene encodes an R2R3 MYB transcription factor that regulates proanthocyanidin synthesis in poplar. *Plant Physiol.*, 150: 924–941
- Nguyen, N.H., J.H. Kim, J. Kwon, C.Y. Jeong, W. Lee, D. Lee, S. Hong and H. Lee, 2016. Characterization of *Arabidopsis thaliana* flavonol synthase 1 (*FLS1*)-overexpression plants in response to abiotic stress. *Plant Physiol. Biochem.*, 103: 133–142
- Owens, D.K., A.B. Alerding, K.C. Crosby, A.B. Bandara, J.H. Westwood and B.S.J. Winkel, 2008. Functional analysis of a predicted flavonol synthase gene family in *Arabidopsis*. *Plant Physiol.*, 147: 1046–1061
- Preuß, A., R. Stracke, B. Weisshaar, A. Hillebrecht, U. Matern and S. Martens, 2009. *Arabidopsis thaliana* expresses a second functional flavonol synthase. *FEBS Lett.*, 583: 1981–1986
- Reddy, A.M., V.S. Reddy, B.E. Scheffler, U. Wienand and A.R. Reddy, 2007. Novel transgenic rice overexpressing anthocyanidin synthase accumulates a mixture of flavonoids leading to an increased antioxidant potential. *Metab. Eng.*, 9: 95–111
- Sanchez, R. and A. Sali, 1997. Evaluation of comparative protein structure modeling by MODELLER-3. *Proteins (Suppl.)*, 1: 50–58
- Schijlen, E., C.H. Ric de Vos, H. Jonker, H. van den Broeck, J. Molthoff, A. van Tunen, S. Martens and A. Bovy, 2006. Pathway engineering for healthy phytochemicals leading to the production of novel flavonoids in tomato fruit. *Plant Biotechnol. J.*, 4: 433–444
- Stracke, R., J.J. Favory, H.E. Gruber, L. Bartelniewoehner, S. Bartels, M. Binkert, M. Funk, B. Weisshaar and R. Ulm, 2010a. The *Arabidopsis* bZIP transcription factor HY5 regulates expression of the *PFG1/MYB12* gene in response to light and ultraviolet-B radiation. *Plant Cell Environ.*, 33: 88–103
- Stracke, R., O. Jahns, M. Keck, T. Tohge, K. Niehaus, A.R. Fernie and B. Weisshaar, 2010b. Analysis of production of flavonol glycosides-dependent flavonol glycoside accumulation in *Arabidopsis thaliana* plants reveals MYB11-, MYB12- and MYB111-independent flavonol glycoside accumulation. *New Phytol.*, 188: 985–1000
- Takahashi, R., S.M. Githiri, K. Hatayama, E.G. Dubouzet, N. Shimada, T. Aoki, S. Ayabe, T. Iwashina, K. Toda and H. Matsumura, 2007. A single-base deletion in soybean flavonol synthase gene is associated with magenta flower color. *Plant Mol. Biol.*, 63: 125–135
- Tamura, K., J. Dudley, M. Nei and S. Kumar, 2007. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol. Biol. Evol.*, 24: 1596–1599
- Wilmouth, R.C., J.J. Turnbull, R.W.D. Welford, I.J. Clifton, A.G. Prescott and C.J. Schofield, 2002. Structure and mechanism of anthocyanidin synthase from *Arabidopsis thaliana*. *Structure*, 10: 93