



Full Length Article

Organophosphorous Solubilizing Mechanism of *Pseudomonas* sp. Isolated from Rhizosphere of Chinese Pine

Guofen Zhao^{1,*,#}, Wei Zhao^{1,#}, Chunbo Yang^{1,#}, Shen Duan², Tongtong Zhao¹ and Dan Wang¹

¹Life Sciences College, Inner Mongolia Agricultural University, Erdos Street, Saihan District, Hohhot City, Inner Mongolia Province, China

²Life Sciences College, Inner Mongolia University, 24 Zhaojun Road, Saihan District, Hohhot City, Inner Mongolia Province, China

*For correspondence: GuofenZhao302@163.com

#These authors contributed equally to this article

Abstract

Phosphorus is an essential element for plants, whereas they cannot use most of phosphorus, which is fixed by calcium, aluminum, iron and organic compounds. Phosphorus solubilizing microorganism (PSM) and its secreting extracellular enzymes have been widely reported in the process of converting soil fixed phosphorus into a bioavailable form. Hence in this study two strains of organophosphate solubilizing bacteria (YP-1, YP-4) were isolated from the rhizosphere soil of Chinese pine (*Pinus tabulaeformis*) in Qingshuihe county (in Hohhot, P. R. China). The organophosphate solubilizing activity was determined to $24.42 \pm 0.52 \mu\text{g mL}^{-1}$ and $23.47 \pm 0.38 \mu\text{g mL}^{-1}$ for YP-1 and YP-4, respectively. YP-1 was identified as *Pseudomonas* sp. YP-4 was identified as *Pseudomonas reingkei* based on phenotypic and molecular characterization, respectively. The most favorite carbon and nitrogen sources were sucrose and $(\text{NH}_4)_2\text{SO}_4$ for YP-1, whereas glucose and KNO_3 for YP-4 in the cultural medium. The extracellular catalytic proteins secreted by microorganisms have the ability to solubilize phosphorus. The extracellular enzymes of two strains were also extracted and determined by Sephadex G-150 method. The molecular weights were 39.8 kDa (Kilodalton) from YP-1 and 38.0 kDa from YP-4. As for the optimized conditions for enzyme catalyzing activities, except for 40°C and 60 min reaction time, the variations were discovered at pH 3.0 and pH 9.0 for YP-1 and YP-4, respectively. By using mass spectrometry, it was found that the corresponding enzyme derived from YP-4 could match 25% to alkaline phosphatase. We conclude that the two strains and their extracellular enzymes can solubilize organophosphate. It is envisioned that the two strains PSM and their related enzymes may find wide applications in future agricultural and biotechnological fields. © 2019 Friends Science Publishers

Keywords: Organophosphate; Bacteria; Enzyme; Chinese pine

Introduction

Chinese pine (*Pinus tabulaeformis*) is one of the most widespread tree species and the main species for afforestation, with economic, social, environmental, and ornamental values in Northwestern China (Yuan *et al.*, 2016). However, the deficiency of available phosphorus in soil environment leads to growth retardation, dark leaves and inhibition of seeding and rooting development system (Li *et al.*, 2006). Phosphorus is a main component of soil nutrients and is one of the indispensable elements for plant growth, which profoundly affect the overall growth progress of plants by influencing metabolic rate-limiting enzyme such as cell division and development, energy transport, signal transduction, macromolecular biosynthesis and photosynthesis (Khan *et al.*, 2002; Shenoy and Kalagudi, 2005; Ahemad *et al.*, 2009; Wang *et al.*, 2009; Hafiza *et al.*, 2013; Rubya and Lqbal, 2016).

Phosphorus is one of the deficient elements, compared to the other major essential macronutrients (with the exception of N). On average, organophosphate is between 30% and 65% of total phosphorus in mineral soils. In organic soils (>20–30% organic matter), organophosphate can approach up to 90% of the total phosphorus (Jones and Oburger, 2011). In addition, phosphate fertilizer is used to promote plant growth in a long-term and later become fixation insoluble form, which results in low absorption by the plants and large accumulation of phosphorus in the soil (Sundara *et al.*, 2002).

The rhizosphere phosphorus microorganisms are capable of solubilizing organophosphate into a bioavailable form via uptake and release of organic and inorganic ions and molecules, including mineral equilibria (the release of complexing or mineral dissolving compounds, such as

organic acid anions, siderophores, protons, hydroxyl ions, and CO₂) and mineralization (the conversions of phosphorus between organic and inorganic forms mediated by extracellular enzymes) (McGill and Cole, 1998; Sims and Pierzynski, 2005). Almost 1-50% of soil bacteria can be classified as phosphorus converting microorganisms (Gyaneshwar et al., 2002). Organophosphate solubilizing bacteria can efficiently degrade organophosphate, including species of the common soil bacteria (*Pseudomonas*, *Azotobacter*, *Burkholderia*, *Bacillus* and *Rhizobium*). Organophosphate mineralization is that the organophosphate converting bacteria convert organophosphate into a bioavailable form by secreting some enzymes (e.g., phosphatase, nuclease and phytase) in soil environment (Jin et al., 2016). Mineralization of organophosphate like nucleic acids and lecithin under favorable conditions are carried out by nuclease and lecithinase. The hydrolysis of both esters and anhydrides of H₃PO₄ can be catalyzed by phosphatases or phosphohydrolases (Tabatabai, 1994). These enzymes are classified as acid and alkaline phosphatases because their maximum activities are related to pH of reaction conditions. The phytic acid to myoinositol and phosphate can be broken by phytase via intermediate inositol polyphosphate (Abul and Charles, 1993; Ramani, 2014). A broad group of enzyme converts organophosphate pesticides into small molecules (non-toxic and soluble in water) by cutting off molecular bonds in organic compounds such as P-F bonds, P-CN (Cheng et al., 1993). These enzymes are widely found in microorganism (*Eschericia coli*, *Bacillus subtilis*, *Pseudomonas*, etc.), animals and plants. In addition, these enzymes probably have potential application for environmental monitoring, pollutant handling, and biosensors (Labry et al., 2005; Fernandez and Kidney, 2007; Tian and Zhou, 2007).

Keeping the above notions in consideration, this study focused on organophosphate solubilizing bacteria identified and characterized from the rhizosphere soil of Chinese pine in Qingshuihe County, Hohhot, Inner Mongolia. Furthermore, their extracellular enzymes to solubilize organophosphate has been purified, identified and characterized, which may have potential industrial, agricultural and biotechnological application.

Materials and Methods

Bacteria Isolation and Biochemical Characterization

Organophosphate solubilizing bacteria were isolated from soil samples which were obtained from mixed soils of the different locations within Chinese pine in Qingshuihe County, Hohhot, Inner Mongolia. Soil samples (10 g) were serially diluted and coated in triplicates on Egg Yolk Solid Medium. The organophosphate solubilizing bacteria can produce phosphoric acid by solubilizing the organophosphate, so that a transparent circle is formed

around the colony. Colonies forming clear surrounding halos were selected as organophosphate solubilizing bacteria and named as YP. The morphology of organophosphate solubilizing bacteria was observed by Gram staining. The physiological and biochemical identification was done according to the bacteriological manual, including the contact enzyme test, methyl red test (MR test), acetyl-methyl methanol test (V-P test), indole test, gelatin liquefaction test, starch hydrolysis test, nitrate reduction test, lecithin test, malonate test, citrate utilization test and sugar fermentation tests (Du et al., 2008).

Molecular Identification of the Bacterial Strain

Extraction of genomic DNA from organophosphate solubilizing bacteria by the CTAB (cetyltrimethylammonium Ammonium Bromide) method, was as described by Chen et al. (2006). The 16S rRNA gene of organophosphate solubilizing bacteria were amplified by PCR (polymerase chain reaction), using universal 27F forward primer (5'-AGAGTTTGATCTGGCTCAG-3') and 1492R reverse primer (5'-CTACGGCTACCTTGTACGA-3'). PCR product was purified using TIANGel Midi Purification Kit, TIANGEN, Beijing (China), then ligated with the cloning vector pMD19-T and transferred to *E. coli* (DH5 α). Nucleotide sequences were determined by HuaDa Gene Company, Beijing. The final partial sequence was submitted to NCBI gene bank in order to obtain the similar sequence. High sequence similarity was selected as reference strains. The phylogenetic tree of the tested strain and the reference strain were constructed by MEGA7.0 software, using the neighbour-joining DNA distance algorithm with a bootstrap of 1000.

Quantitative Estimation of Bacterial Ability to Degrade Organophosphate

The organophosphate solubilizing bacteria were isolated in triplicate, from egg yolk solid medium after 4 days at 30°C. The diameter of dissolved phosphorus transparent circle (D) and the diameter of the colony (d) were determined by a vernier caliper. Preliminary determination of the ability of strains to solubilize phosphorus was determined by calculating the size of D/d. The organophosphate solubilizing bacteria were inoculated into the beef extract peptone liquid medium, containing: Beef extract 3.0 g; NaCl 5.0 g; Peptone 10.0 g. The flasks were incubated in an incubator shaker at a shaking speed of 160 rpm for 37°C, the bacteria fermentation broth was measured at 660 nm every half hour until the OD was 0.5. After 1% inoculation was added to the 30 mL Monkina's medium, (glucose 10 g; CaCO₃ 5 g; NaCl 0.3 g; KCl 0.3 g; (NH₄)₂SO₄ 0.5 g; MnSO₄·H₂O 0.03 g; FeSO₄·7H₂O 0.03 g; pH 7.0 ~ 7.2), the flasks were incubated at 30 °C in an orbital shaker at 160 rpm for 6 days. In triplicate, non-inoculated medium served as a control. Subsequently, centrifugation for 20 min

(4°C, 10000 rpm). 5 mL reagent (containing: Ammonium molybdate 1.0 g; H₂SO₄: 15.3 mL; Antimony potassium tartrate: 10 mL; Ascorbic acid; 1.5 g), and 20 mL distilled water was added to 25 mL of the supernatant in 50 mL volumetric flask, and mixed. After setting aside for 30 min at room temperature, the absorbance was measured at 660 nm. The corresponding phosphorus concentration was checked on the standard curve and the phosphorus content was calculated. The difference of soluble phosphorus content between the experimental and the control group was used as the ability of the tested strain to solubilize phosphorus ($\mu\text{g L}^{-1}$). Three replicates were used. With (NH₄)₂SO₄ as the nitrogen source, carbon sources were glucose, sucrose or maltose (the carbon source concentration was 10 g L⁻¹), and with glucose as the carbon source, nitrogen sources were (NH₄)₂SO₄, urea or KNO₃ (the nitrogen source concentration was 0.5 g L⁻¹). The phosphorus contents were then detected.

Isolation and Purification of Extracellular Enzyme with Ability to Solubilize Phosphorus

The organophosphate solubilizing bacteria were inoculated in Monkina's medium, 37°C in an orbital shaker at 160 rpm for 48 h. Then centrifuged (4°C, 10000 rpm) for 20 min to obtain cell-free culture supernatant. The supernatant was lyophilized by vacuum freeze-dried and used 5 mL Tris-HCl (50 mM, pH 7.0) to dissolve the lyophilized powder. Lyophilized supernatant was purified by filtration chromatography, using the Sephadex G-150 gel (GE Healthcare, Sweden). The elute was Tris-HCl (50 mM, pH 7.0) at a flow rate of 23 mL min⁻¹. The peak at 280 nm was detected by a nucleic acid protein analyzer and then collected the elute with protein, thereby obtaining purified extracellular enzyme. The reaction between 1 mL of each peak collection solution and 24 mL Monkina's medium as substrate in erlenmeyer flasks containing were started at 30°C for 90 min; then added 20% trichloroacetic acid to terminate the reaction, with 5 mL reagent and distilled water to the volume of 50 mL. After setting aside for 30 min at room temperature, absorbance was measured at 660 nm and the phosphorus content was calculated. The purity of the obtained enzyme with ability to solubilize phosphorus was analyzed by 12% SDS-PAGE. The samples were stained with Coomassie Blue R-250. The standard curve constructed by measuring the migration distance between protein standard and target protein with indicator, respectively. The molecular weight of the protein with the ability to solubilize phosphorus was calculated according to the standard curve and was initially identified. A single target protein band was cut for mass spectrometry analysis with a mobile phase: A: 0.1% formic acid solution; B: 0.1% formic acid-acetonitrile solution; flow rate: 300 mL min⁻¹; elution gradient: 3 to 8% B elution for 1 min, 8 to 40% B elution for 5 min, 40 to 85% B for 1 min and finally eluted

with 85% B for 1 min; Full MS (resolution: 70000; scanning range: 350-1600); dd-MS2: (resolution: 17500); and injection volume of 10 μL .

Characterization of Partially Purified Extracellular Enzyme

The characterization of partially purified enzymes activity was studied by changing the reaction parameters, such as pH from 3.0 to 11.0, temperature (at 10°C intervals) from 20 to 60°C, and the reaction time of 0 to 120 min. The reaction between 1 mL of each peak collection solution and 24 mL Monkina's medium as substrate in erlenmeyer flasks containing were started at 30°C for 90 min; then added 20% trichloroacetic acid to terminate the reaction, with 5 mL reagent and distilled water to the volume of 50 mL. After setting aside for 30 min at room temperature, absorbance was measured at 660 nm and the phosphorus content was calculated by the formula (Tsang *et al.*, 2007).

Statistical Analysis

Data on organophosphorus solubilizing capacity under different conditions were analyzed of multiple comparisons (ANOVA) (SPSS 20.0; [Koneri *et al.*, 2008](#)). Significant effects of treatments were determined by the magnitude of F value ($P \leq 0.05$).

Results

Isolation and Identification of Organophosphate Solubilizing Bacteria

Organophosphate solubilizing bacteria were isolated from the rhizosphere soil of Chinese Pine. Two strains of organophosphate solubilizing bacteria were isolated and named YP-1 and YP-4. Based on the morphological and biochemical analysis, YP-1 was rod-shaped, gelatin liquefaction, citrate, and fermentation of glucose were positive, whereas gram staining, methyl red, VP tests, indole tests, amylase, nitrate reduction tests, lecithin enzyme tests and fermentation of lactose, sucrose and maltose were negative. YP-4 was found to be resembled with YP-1 about morphological and biochemical analysis, except the fermentation of glucose, which was negative. Further, genus *Pseudomonas* was confirmed by BLAST analysis data of the 16S rRNA gene sequence. YP-1 showed 99% similarity with the genus *Pseudomonas* sp. (KC355362). YP-4 showed 99% similarity with the *Pseudomonas reinekei* (LT629709.1). By comparing the nucleotide sequences of 16S rRNA sequences of the isolates YP-1 and YP-4 with different *Pseudomonas* sp. from NCBI database, a phylogenetic tree was constructed, we found that the YP-1 is most closely related to *Pseudomonas* sp., and YP-4 is most closely related to *Pseudomonas reinekei* (Fig. 1).




Fig. 1: Phylogenetic tree based on 16S rRNA gene sequences by Neighbor Joining method (using MEGA 7.0), showing the relationship between (A) YP-1, (B) YP-4 and other members of the *Pseudomonas* sp. The Genbank nucleotide accession numbers are listed next to the strain names. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) is shown next to the branches. The scale bars represent 0.5 substitution/site

Solubilizing of Organophosphate

In evaluation of the organophosphate solubilizing activity of the strain *Pseudomonas* sp., the D/d of the YP-1 was 3.15, and the YP-4 was 2.27. Further, the definite activity of the bacteria was detected by combining with quantitative methods. The soluble phosphorus content in the supernatant was used to indicate ability to solubilize organophosphate of the bacteria. When the carbon source was sucrose, the nitrogen source was $(\text{NH}_4)_2\text{SO}_4$ in the Monkina's medium, the organophosphate solubilizing activity of the bacteria strain YP-1 was the highest. With the addition of glucose and KNO_3 to the culture medium, YP-4 had highest activity for solubilizing organophosphate (Fig. 2A and B).

Purification of Extracellular Enzyme

The extracellular proteins secreted by YP-1 or YP-4 were isolated and purified by Sephadex G-150 gel (Fig. 3). It was found that only one protein was secreted into the cell fermentation supernatant by the YP-1, was $(21.94 \pm 3.82 \mu\text{g mL}^{-1})$. The YP-4 was capable of secreting two proteins,




Fig. 2: Influence of different culture medium: (A) Nitrogen sources and (B) Carbon sources on crude enzymes production by the degrade, YP-1 and YP-4. Different superscripts indicate significant difference among different strain (Lowercase letters are significant, ANOVA, Duncan's test; $p < 0.05$)

while only one of the enzyme had ability to degrade organophosphate, was $(16.14 \pm 0.98 \mu\text{g mL}^{-1})$. The smaller protein of YP-4 had no activity.

Identification of Protein that Solubilize Organophosphate

According to the standard curve of protein electrophoresis ($y = -0.9954x + 5.1857$), the molecular weight of the extracellular enzymes of YP-1 and YP-4 were calculated as 39.8 kDa and 38.0 kDa, respectively (Fig. 5). After mass spectrometry determination, the extracellular enzyme secreted by YP-1 did not find a matching protein, whereas the extracellular enzyme secreted by YP-4 was matched with an alkaline phosphatase, had the highest coverage 25%, protein accession for WP_016986280.1.

Characterization of Extracellular Organophosphate Solubilizing Enzymes

Enzymes activities were greatly influenced by the incubation temperature (Fig. 6A). The optimum temperature




Fig. 3: The extracellular enzyme of YP-1 was purified by Sephadex G-150 gel chromatography. A peak appeared at 1.8 h




Fig. 4: The extracellular enzyme of YP-4 was purified by Sephadex G-150 gel chromatography. Two peaks (1 and 2) appeared at 1.2h and 2.2h




Fig. 5: SDS-PAGE analysis of extracellular proteins were secreted by YP-1 and YP-4. The purified proteins (lane YP-1 and YP-4) were separated by 12% (w/v) SDS-PAGE followed by staining with Coomassie Blue R-250. Lane M: molecular weight markers

for the enzymes activity were 40°C, with enzymes activities of $67.68 \pm 0.18 \mu\text{g mL}^{-1}$ (YP-1) and $23.78 \pm 0.59 \mu\text{g mL}^{-1}$ (YP-4), respectively. The enzymes activities of YP-1 had a large difference at different






Fig. 6: Effect of different reaction parameters (A) Temperature, (B) pH, and (C) Reaction time on purified enzymes production by the isolate, YP-1 and YP-4. (Lowercase letters are significant. ANOVA, Duncan's test; $p < 0.05$)

temperature conditions. The enzymes activities of YP-1 showed “decrease first and then increase” with increasing pH (Fig. 6B). The maximum activity of enzyme was recorded ($11.01 \pm 0.18 \mu\text{g mL}^{-1}$) at pH 3.0. At pH 9.0, the enzyme of YP-4 had the greatest ability to solubilize organophosphate, the content of soluble phosphorus was $22.05 \pm 0.37 \mu\text{g mL}^{-1}$ in the reaction system. By the assay of enzymes activities with increasing reaction time (Fig. 6C), the reaction between enzyme and organophosphate was accomplished within 30 min.

Discussion

The research of organophosphate solubilizing bacteria is focused on soil microorganisms, and the amount is greater in the rhizosphere (20 - 40% of the total) as compared to non-rhizosphere (10 - 15% of the total amount) region (Swaby and Sperber, 1958). Two strains of YP-1 and YP-4 with the ability to degrade organophosphate were obtained from rhizosphere of Chinese pine, were *Pseudomonas* sp. and *Pseudomonas reinekei*, respectively. In comparison to the present study, Jorquera *et al.* (2008) isolated organophosphate solubilizing bacteria from rhizosphere soils of different plants (ryegrass, white clover, wheat, oats and lupines), which exhibited maximum ability of solubilizing phosphorus for $29.92 \pm 1.45 \mu\text{mol L}^{-1}$. Tao *et al.* (2008) isolated ten organophosphate solubilizing bacteria from subtropical irrigation and temperate non-irrigated soils, which could degrade phosphorus in the range of 13.8 ~ 62.8 $\mu\text{g mL}^{-1}$. Rhizosphere symbiotic bacteria, *Pseudomonas* sp. and *Bacillus* sp. have strong ability to solubilize organophosphate in the soil (Wei *et al.*, 2014). Lin *et al.* (2000) found that *Bacillus* sp. was the main part of the organophosphate solubilizing bacteria, followed by *Pseudomonas* sp., in the rhizosphere soil of farmland (wheat-maize), lawn (fescue), woodland (elm) and vegetable land (cauliflower).

According to the size of the halo zone, the YP-4 ability of solubilizing organophosphate was higher than YP-1. However, the organophosphate solubilizing ability of two bacterium were quantitatively determined in liquid culture by molybdenum blue colorimetric method, showing the opposite result. The size of the halo zone around colony is not always accurate to determine the ability about solubilize organophosphate of the bacteria (Behera *et al.*, 2017). This may be due to various diffusion rates of different organic acids secreted by an organism (Nautiyal, 1999). When the lecithin in the Monkina medium became soluble phosphorus and reacted with ammonium molybdate to form ammonium salt, the ascorbic acid was reduced to molybdenum blue (absorbance at 660 nm), organophosphate solubilizing efficiency of microorganisms can be easily detected. Hence, the molybdenum blue colorimetric method is better way to evaluate organophosphate solubilizing efficiency.

The ability to solubilize organophosphate of microorganisms will change as the living environment changes. A wide range of carbon source can be used by microorganisms; differences in carbon sources are significant for specific bacterial strains. As a kind of life element, nitrogen is an important constituent element of proteins and nucleic acids. For this reason, this study further explored the effects of different carbon and nitrogen sources on the organophosphate solubilizing efficiency of YP-1 and YP-4. Similar studies have been conducted by many reachers, they also analyzed the different carbon and nitrogen sources on the growth and physiological function, *Tricholoma mongolicum* Imai (Chang *et al.*, 2011),

Trichoderma harzianum (Rajput *et al.*, 2018) and *Beauveria bassiana* (Dale and Shinde, 2018) which indicated the carbon and nitrogen is the major component on the growth and development of organisms.

Determining the catalytic characteristics of enzyme, is not only beneficial to expound the enzyme function and the mechanism of enzyme action, but also has important practical significance in science developing. The activity of purified enzyme is greatly influenced by reaction parameters such as pH, temperature, and reaction time. Many enzyme researchers have also shown that temperature and pH are crucial to enzyme activity (Hirata *et al.*, 2015; Joana and Alejandra, 2015; Lorenzen *et al.*, 2017). In order to optimize the soluble phosphorus production, it is necessary to evaluate optimum conditions of these parameters. The pH can affect the ionization of binding and catalyzing sites of enzyme to substrate, alter the charge distribution, thereby affecting activity of the enzyme (Chen *et al.*, 2009; Sasirekha *et al.*, 2012). The extracellular enzyme of YP-1 was stable with the different conditions of pH, thus it is an acid-alkali resistant protein. Since the extracellular enzyme YP-4 had the highest activity at pH 9.0, it is speculated that YP-4 produced an alkaline phosphatase. The number of activated substrate molecules will increase as the temperature increases the reaction speed will accompany by faster. When the temperature reaches the appropriate level, continued increase in temperature will cause the enzyme denaturation, resulting in reaction rate is reduced (Peterson *et al.*, 2007). In addition, it had invalid on enzyme activity with the prolongation of the reaction time, indicating that whole process about the binding and catalyzing of the enzyme and substrate is stable and quick.

In this study, it was found that only one protein was secreted into the cell fermentation supernatant by YP-1, and had the solubilizing organophosphate activity. YP-4 was capable of secreting two proteins, while only one of the enzyme had ability to solubilize organophosphate. These two purified enzyme were detected by SDS-PAGE, protein bands were single, indicating that the protein contains sufficient purity. The corresponding enzymes derived from YP-4 can match 25% to alkaline phosphatase. This result is consistent with the previous speculation that enzyme derived from YP-4 had best activity with pH 9.0. Unfortunately, as the protein sequence is unknown, the enzyme secreted by YP-1 did not find a matching protein. The reliable protein sequence will be detected by others proteomics technologies in the next study. The gene encoding phosphatase will be tried to be cloned from the genome of YP-4 and YP-1 by PCR, then the structure of phosphatase will be analyzed to understand solubilizing mechanism of organophosphate in details in the future study.

Conclusion

In this study, the organophosphorous solubilizing bacteria

were isolated in the rhizosphere soil of Chinese pine (*Pinus tabulaeformis*) in Qingshuihe county (in Hohhot, P. R. China). Two strains were identified as *Pseudomonas* sp., and *Pseudomonas reinekei*. We found that the bacteria were secreted extracellular enzymes to solubilize organophosphorous. Further, the extracellular enzymes were isolated and purified, and had great abilities of organophosphorous solubilizing. It may have probable use as bio-inoculants to increase soil fertility by minimizing fertilizer application, which can promote sustainable agriculture and help to meet future needs.

Acknowledgements

This study was supported by the Inner Mongolia forestry key science and technology projects. The help and cooperation of the field staff of forest department of Chinese pine is grate-fully acknowledged.

References

- Abul, H.J. and H.D. Charles, 1993. Identification of active site residues in *Aspergillus ficuum* extracellular pH 2.5 optimum acid phosphate. *Biochem. Biophys. Res. Commun.*, 192: 754–759
- Ahemad, M., A. Zaidi, M.S. Khan and M. Oves, 2009. Biological importance of phosphorus and phosphate solubilizing microorganisms—an overview. In: *Phosphate Solubilizing Microbes for Crop Improvement*, pp: 1–4. Khan, M.S. and A. Zaidi (eds.). Nova, New York
- Behera, B.C., H. Yadaz, S.K. Singh, R.R. Mishra and B.K. Sethi, 2017. Phosphate solubilization and acid phosphatase activity of *Serratia* sp. isolated from mangrove soil of Mahanadi river delta, Odisha, India. *J. Genet. Eng. Biotechnol.*, 15: 169–178
- Chang, X., Yangling (China), C.O.L. Sciences, 2011. Effect of different carbon sources and nitrogen sources on the growth of *tricholoma mongolicum* imai. *J. Northwest. A. F. Univ.*, 39: 203–207
- Chen, Y.P., P.D. Rekha, A.B. Arun, F.T. Shen, W.A. Lai, and C.C. Young, 2006. Phosphate solubilizing bacteria from subtropical soil and their tricalcium phosphate solubilizing abilities. *Appl. Soil. Ecol.*, 34: 33–41
- Chen, H.Q., Q.Y. Xu, Q. Ye, W. Chen and H. Zhang, 2009. Bioconversion kinetics of conjugated linoleic acid by *Lactobacillus plantarum* ZS2058. *Acta. Microbiol. Sin.*, 49: 174–179
- Cheng, T.C., S.P. Harvey and A.N. Stroup, 1993. Purification and properties of a highly active organophosphate acid anhydrolase from *Alteromonas undina*. *Appl. Environ. Microb.*, 59: 3138–3140
- Dale, N.S. and S.S. Shinde, 2018. Studies on effect of different concentration of carbon and nitrogen sources on growth and biomass production of *Beauveria bassiana*. *Internat. J. Acad. Res. Dev.*, 2: 457–458
- Du, Y.S., X. Wei, Y.Z. He, G. Wei, H. Hampel, R. Dodel, and M.R. Farlow, 2008. P2-380: Identification and characterization of human autoantibodies that may be used for the treatment of prion diseases. *Alzheimers. Dement.*, 4: 484
- Fernandez, N.J. and B.A. Kidney, 2007. Alkaline phosphatase: beyond the liver. *Vet. Clin. Pathol.*, 34: 68–70
- Gyaneshwar, P., G.N. Kumar, L.J. Parekh and P.S. Poole, 2002. Role of soil microorganisms in improving P nutrition of plants. *Plant Soil*, 245: 83–93
- Hafiza, M.S., G.Z. Jahangir, I.A. Nasir, M. Iqtidar and MIqbal, 2013. Isolation and characterization of phosphate-solubilizing bacteria from rhizosphere soil. *Biotechnol. Biotec. Eq.*, 27: 4248–4255
- Hirata, A., S. Kishino, B. Park, M. Takeuchi, N. Kitamura and J. Ogawa, 2015. A novel unsaturated fatty acid hydratase toward c16 to c22 fatty acids from *Lactobacillus acidophilus*. *J. Lipid. Res.*, 56: 1340–1350
- Jin, T.T., J.H. Ren and R.X. Liu, 2016. Identification, Characterization and Growth-promoting Effects of an Organophosphate-solubilizing Bacterium from *Taxus chinensis* var. *mairei* Rhizosphere. *Acta Bot. Boreali-Occidentalia Sin.*, 36: 1819–1827
- Jones, D.L. and E. Oburger, 2011. Solubilization of phosphorus by soil microorganisms. In: *Heidelberg: Phosphorus in Action*, pp: 169–198. Bünemann, E., A. Oberson and E. Frossard (eds.). Berlin
- Jorquera, M.A., M.T. Hernández, Z. Rengel, P. Marschner and M.L. Mora, 2008. Isolation of culturable phosphobacteria with both phytate-mineralization and phosphate-solubilization activity from the rhizosphere of plants grow in a volcanic soil. *Biol. Fertil. Soil.*, 44: 1025–1034
- Khan, M.S., A. Zaidi and M. Aamil, 2002. Biocontrol of fungal pathogens by the use of plant growth promoting rhizobacteria and nitrogen fixing microorganisms. *J. Ind. Bot. Soc.*, 81: 255–263
- Koneri, R., R. Balaraman, K.M. Vinoth and Hariprasad, 2008. Cardioprotective effect of *Momordica cymbalaria* fenzl against experimental myocardial injury induced by isoproterenol. *Inter. J. Pharmacol.*, 2: 1–6
- Labry, C., D. Delmas and A. Herbland, 2005. Phytoplankton and bacterial alkaline phosphatase activities in relation to phosphate and DOP availability within the Gironde plume waters (Bay of Biscay). *J. Exp. Mar. Biol. Ecol.*, 318: 213–225
- Li, B.Q., O. WahLiew and A.K. Asundi, 2006. Pre-visual detection of iron and phosphorus deficiency by transformed reflectance spectra. *J. Photoch. Photobio. B*, 85: 131–139
- Lin, Q.M., X.R. Zhao, Y.X. Sun and J. Yao, 2000. Community characters of soil phosphobacteria in four ecosystems. *Soil. Environ. Sin.*, 9: 34–37
- Lorenzen, J., R. Driller, A. Waldow, F. Qoura, B. Loll and T. Brück, 2017. *Rhodococcus erythropolis* oleate hydratase: a new member in the oleate hydratase family tree—biochemical and structural studies. *Chemcatchemistry*, 10: 407–414
- McGill, W.B. and C.V. Cole, 1998. Comparative aspects of cycling of organic C, N, P through soil organic matter. *Geoderma*, 26: 267–268
- Nautiyal, C.S., 1999. An efficient microbiological growth medium for screening phosphate solubilizing microorganisms. *FEMS Microbiol. Lett.*, 170: 265–270
- Joana O.A. and H.S. Alejandra, 2015. Functional characterization of a fatty acid double-bond hydratase from *Lactobacillus plantarum* and its interaction with biosynthetic membranes. *Biochimica. Et. Biophysica. Acta.*, 1848: 3166–3174
- Peterson, M.E., R.M. Daniel, M.J. Danson and R. Eiseenthal, 2007. The dependence of enzyme activity on temperature: determination and validation of parameters. *Biochem. J.*, 402: 331–337
- Rajput, A.Q., M.A. Khanzada and S. Shahzad, 2018. Effect of different organic substrates and carbon and nitrogen sources on growth and shelf life of *trichoderma harzianum*. *Int. J. Agric. Biol.*, 16: 893–898
- Ramani, V., 2014. Organic phosphate mineralization by *Bacillus sphaericus* and *Pseudomonas cepacia*. *J. Agr. Sci. Tech-Iran*, 10: 913–922
- Ruby, S. and M.T. Lqbal, 2016. Phosphorus use efficiency by wheat plants that grown in an acidic soil. *Braz. J. Sin. Technol.*, 3: 18
- Sasirekha, B., T. Bedashree and K.L. Champa, 2012. Optimization and partial purification of extracellular phytase from *Pseudomonas aeruginosa* p6. *J. Exp. Biol.*, 2: 95–104
- Shenoy, V.V. and G.M. Kalagudi, 2005. Enhancing plant phosphorus use efficiency for sustainable cropping. *Biotechnol. Adv.*, 23: 501–513
- Sims, J.T. and G.M. Pierzynski, 2005. Chemical of phosphorus in soil. In: *Madison: Chemical Processes in Soil*, pp: 151–192. Tabatabai, A.M. and D.L. Sparks, (eds.). Soil Science Society of America
- Sundara, B., V. Natarajan and K. Hari, 2002. Influence of phosphorus solubilizing bacteria on the changes in soil available phosphorus and sugarcane and sugar yields. *Field Crop Res.*, 77: 43–49
- Swaby, R.J. and J. Sperber, 1958 Phosphate dissolving microorganisms in the rhizosphere of Legumes. In: *New York: Nutrition of the Legumes*, pp: 289–294. Hallsworth, E.G. (eds.)

- Tabatabai, M.A., 1994. Soil enzymes. In: *madison: Methods of Soil Analysis, Microbial and Biochemical Properties, Soil Science Society of America Book Series 5*, pp: 775–833. Weaver, R.W. (eds.). Soil Science Society of America
- Tao, G.C., S.J. Tian, M.Y. Cai and G.H. Xie, 2008. Phosphate-solubilizing and mineralizing abilities of bacteria isolated from soils. *Pedosphere*, 18: 515–523
- Tian, J.R. and P.J. Zhou, 2007. Phosphorus fractions of floodplain sediments and phosphorus exchange on the sediment-water interface in the lower reaches of the Han River in China. *Ecol. Eng.*, 30: 264–270
- Tsang, S., F. Phu, M.M. Baum and G.A. Poskrebyshev, 2007. Determination of phosphate/arsenate by a modified molybdenum blue method and reduction of arsenate by S₂O₄²⁻. *Talanta*, 71: 1560–1568
- Wang, X.R., Y.X. Wang, J. Tian, B.L. Lin, X.L. Yan and H. Liao, 2009. Overexpressing AtPAP15 enhances phosphorus efficiency in soybean. *Plant Physiol.*, 151: 233–240
- Wei, S., G. Jie, Y.D. Li, X. Qian and X.J. Wang, 2014. Screening, identification and growth characteristics of soil organic phosphate converting bacterial strains. *J. Northwest A F Univ.*, 42: 199–206
- Yuan, H.W., S.H. Niu, X.Q. Zhou, Q.P. Du, Y. Li and W. Li, 2016. Evaluation of seed production in a first-generation seed orchard of Chinese pine (*Pinus tabuliformis*). *J. For. Res-Jpn.*, 27: 1–6

(Received 16 July 2018; Accepted 12 September 2018)

IB Press