



**Full Length Article**

## Genome-wide Identification and Expression Analysis of the PIN Gene Family during Abiotic Stress in Tomato (*Solanum lycopersium*)

Chunli Zhang, Qi Yuan, Tingting Zhao, Di Xiao and Xiangyang Xu\*

College of Horticulture & Landscape Architecture, Northeast Agricultural University, Harbin 150030, China

\*For correspondence: 1009821258@qq.com; xxy709@126.com

### Abstract

It is widely believed that the PIN gene family is the most important auxin efflux carrier in plants since the separation of pin-formed mutant from *Arabidopsis thaliana* in 1991. Auxin transport and accumulation can be indirectly measured by the expression level of PIN genes. In recent years, members of the PIN gene family have been cloned in different species. Ten PIN genes (SIPINs) were identified in the genome data of tomato using the Arabidopsis PIN protein family for reference sequences. Phylogenetic analysis showed that 10 members were divided into “short” and “long” PINs. Chromosomal distribution map revealed that the SIPIN genes were unevenly distributed in eight chromosomes. We also studied the expression of 9 SIPIN genes under cold, drought and salinity by qRT-PCR, (SIPIN2 was not expressed in tomato leaves). The obtained levels of expression of the same genes were different under different stresses. Some genes were up-regulated under some stresses but were down-regulated under other stresses. The relative expression of SIPIN6 clearly changed under cold and drought stress. The results here would provide the theoretical basis for the molecular cloning and resistance breeding of PIN genes in tomato. © 2018 Friends Science Publishers

**Keywords:** Abiotic stresses; Auxin efflux carriers; PIN; Relative expression; Tomato

### Introduction

In early 1880, it was found that there is a transport signal substance – auxin. Auxin is the only hormone that undergoes polar transport in plants and is an indispensable regulator of plant growth and development. Auxin is synthesized in many plant tissues *via* several different pathways, including the stem apex, young leaves, and flowers as well as during the growth of lateral root vascular tissue (Zhao, 2010), and then transported to the other tissue parts of the plant. The concentration gradient of auxin is typically established by the different sites of synthesis and methods of transportation. Auxin is involved in plant apical dominance, tropism, vascular differentiation, floral tissue differentiation, photomorphogenesis, embryo formation, root development and plant responses to internal and external stimuli (Dubrovsky *et al.*, 2008; Xi *et al.*, 2016). The interaction of auxin, gibberellins and cytokinins promote the growth of plants by promoting the growth of cells (Fleet and Sun, 2005; Růžička *et al.*, 2009). Different auxin concentrations in different tissues are caused by polar auxin transport (PAT). The PAT study using coleoptile as a test material improved that auxin is transported from the morphological top to the morphological bottom of the plant, and this transportation direction is not affected by gravity. There is auxin carrier protein at the bottom of each cell membrane, but not on the top of the cell membrane (Yu and Cui, 2009). Multiple

classes of auxin transport proteins participate in cell-to-cell auxin transport. Auxin is transported by the efflux carriers pin-formed (PIN) and Multidrug-Resistant/P-glycoprotein (MDR/PGP), and the influx carriers auxin resistant 1/like aux1 (AUX/LAX). The PIN protein has a speed limit function during the output process of auxin, and sensitivity to auxin output inhibitory factor. Recently, some members of the PIN gene family are localized to the endoplasmic reticulum and participated in the regulation of intracellular auxin homeostasis, the plasma membrane (Barbez *et al.*, 2012; Feraru *et al.*, 2012). Other members are located in the plasma membrane and involved in various tropic responses and apical shoot establishment in *Arabidopsis*.

There have been some studies on auxin transport genes in monocotyledons and dicotyledonous plants. Many PIN genes have been reported in detail in dicotyledonous plants, but there are also a few PIN genes whose research is rare in monocotyledons. PIN1 was one of the most studied and earliest members of the PIN gene family (Goto *et al.*, 1987). Gälweiler *et al.* (1998) cloned the PIN1 gene as an auxin transporter from *Arabidopsis thaliana* by transposon tagging. The PIN gene has been predicted in potato, *Medicago truncatula* (Schnabel and Frugoli, 2004), maize (Carraro *et al.*, 2006), *Brassica juncea* (Ni *et al.*, 2002) and *Mangifera indica* (Li *et al.*, 2012). In higher plants, all PIN proteins are polar distributed on one side or another side of the cell, which is consistent with auxin transport. PIN proteins are

characterized as cell specific and tissue specific. Different members of the PIN family are expressed in different plant parts, and the function of PIN proteins changes with a change in polarity position. The subcellular localization of PIN genes also has the characteristics of dynamic change and not only is affected by the growth and development of plants but also is restricted by external environmental conditions. Presently the information of PIN-dependent polar auxin transport in plants mainly comes from the extensive investigation of the PIN gene family in rice (*Oryza sativa*) (Ma and Jie, 2010) and *Arabidopsis*. In *Arabidopsis*, the PIN family comprises of 8 members and is divided into two groups. AtPIN1-AtPIN4, AtPIN6 and AtPIN7 belonged to the typical long PINs. These long PIN proteins are located in the plasma membrane, and have a relatively long central hydrophilic loop, which share high sequence similarity especially in the hydrophobic domains of both N- and C-termini (Roumeliotis *et al.*, 2013). These proteins are primarily responsible for the transport of auxin to the outside of cell and unevenly distributed on the cell membrane and participated in various tropic responses (Robert *et al.*, 2013). In contrast to the long PINs, the short PINs include AtPIN5 and AtPIN8, which lack a central loop domain. They are localized in the endoplasmic reticulum and mediated the auxin communication of the cytoplasm and endoplasmic reticulum. They also participate in the homeostasis of auxin and compartmental localization by working together with members of the PIN auxin efflux carriers (Ding *et al.*, 2012; Mravec *et al.*, 2009; Cazzonelli *et al.*, 2013). In addition, the PIN gene has been studied to be involved in abiotic stress responses, including those that involve dehydration, salt and drought (Shen *et al.*, 2010).

Tomato (*Solanum lycopersicum*) is one of the most important vegetable crops and is also a model plant for molecular biology research in *Solanaceae* plant. Tomato has high economic and medicinal value. In 2012, the tomato genome sequencing project was completed (Consortium 2012), which laid the foundation for the whole-genome bioinformatics analysis of the PIN gene family, and studies on the bioinformatics analysis of the PIN gene in tomato have been reported (Zhao *et al.*, 2017). Recently, the expression of three PIN genes was examined in tomato vegetative and reproductive organs. Nishio *et al.* (2010) focused on PIN gene expression patterns and auxin distribution patterns during early fruit development. The tissue-specific expression and the spatial and temporal expression patterns of auxin accumulation with respect to the tomato PIN gene family have also been reported; these studies have especially emphasized the process of fruit growth (Pattison and Catalá 2012). However, no studies have analysed the effects of abiotic stresses on the tomato PIN family. In sorghum, the expression of 5 SbPIN genes highly increased under ABA, salt and drought treatments, whereas the rest of the 6 SbPIN genes were almost inhibited. Here, we present a genome-wide of the tomato PIN gene family and its expression profiling under abiotic stresses such as salt, cold and drought. This work identified the tomato PINs associated with abiotic

stresses responses. Some of these PINs would be candidate genes for further functional studies.

## Materials and Methods

### Identification of PIN Gene Family Members in Tomato

Protein sequences of the tomato PIN gene were obtained from the SOL Genomics Network (SGN, <http://solgenomics.net/search/loci>). According to analysing the conserved PIN domains (PF03547), all candidate protein sequences of tomato were further confirmed using the software programs HMMER (<http://www.ebi.ac.uk/Tools/hummer/search/hmmsearch>) and Pfam (<http://pfam.janelia.org>). Essential on exons and chromosome locations of tomato PIN members were obtained from the SGN sequence database. Theoretical pI, molecular weight and other physicochemical properties of PIN amino acids were calculated using the online program ProtParam (<http://expasy.org/tools/protparam/>).

### Prediction of the Secondary Structure of Tomato PIN Members

Tomato PIN signal peptides were predicted using the SignalP 4.1 (<http://www.cbs.dtu.dk/services/SignalP/>) online tool. The secondary structure of tomato PIN members were forecast utilizing SOPMA online software ([https://npsa-prabi.ibcp.fr/cgi-bin/npsa\\_automat.pl?page=npsa\\_sopma.html](https://npsa-prabi.ibcp.fr/cgi-bin/npsa_automat.pl?page=npsa_sopma.html)).

### Phylogenetic Tree and Multiple Sequence Alignment

The sequence alignment of tomato, sorghum, rice and *Arabidopsis* was performed by the software ClustalX (Sun *et al.*, 2015). The results of the sequence alignments were used to construct an unrooted phylogenetic tree using the neighbour joining method with a bootstrap analysis of 1000 replicates by MEGA 5.2 program (Tamura *et al.*, 2011; Chai and Subudhi, 2016). The multiple sequence alignment was performed using the DNAMAN program.

### Conserved Motifs and Chromosomal Mapping

Conserved motifs of the tomato PIN members were statistically identified by the MEME program (<http://meme.nbcr.net/meme/>) according to the searched amino acid sequence. Each SIPIN gene chromosome localization were determined based on the protein sequence from SGN. The SIPIN genome location was drawn from top to bottom by the MapInspect software.

### Plant Growth Conditions and Quantitative RT-PCR (qRT-PCR) Analysis

The experiment was performed in the horticultural station of Northeast Agricultural University (Harbin, China), during 2016-2017. Tomato seeds (*S. lycopersicum* cv. Moneymaker)

were grown in a greenhouse. Unified managements of watering, pest and disease controls were used during the whole growth period (Liu *et al.*, 2015). For abiotic stress treatments, five- or six-leaf seedlings were treated with salinity (100 mM NaCl), cold (5°C) and drought (10% PEG6000). The leaves of seedlings were saved at 0, 2, 4, 8, 12 and 24 h after stimulation of the salt and cold treatments, and three biological replicates were performed for every treatment. For drought treatment, the young leaves were collected separately at 0, 4, 6, 8 and 12 h after treatment. The samples were frozen immediately in liquid nitrogen after collection and kept at -80°C, and each sample was repeated three times.

Total RNA was extracted with using Trizol reagent (Invitrogen) according to the manufacturer's guidances. First-strand cDNA was synthesized from 3 mg of total RNA from each sample using the TransScript® One-Step gDNA removal kit and cDNA Synthesis SuperMix reverse transcriptase. Quantitative real-time PCR performed in an iQ 5 system using SYBR® Green I. Primer pairs (Table 1) for individual gene families were designed using Primer Premier 5.0 program. Among them, SIPIN1, 5, and 6 primer pairs have been published (Pattison and Catalá 2012). The results of RNA were checked using agarose gel electrophoresis and analysed by AlphaEaseFC image (Zhang *et al.*, 2014), and images were collected under UV light. The qPCR reaction mixture contained SYBR® Green Master Mix (10 µL), each primer (0.5 µL), ROX Reference Dye one (0.5 µL), cDNA template (1 µL) and sterile distilled water up to a total volume of 20 µL. The thermal conditions were as follows: 95°C for 10 min and 40 cycles of 95°C for 5 s, 56°C for 15 s, and 72°C for 30 s. Slacton was employed as an internal control, and qRT-PCR (quantitative reverse transcription-polymerase chain reaction) data were analysed in accordance with the  $2^{-\Delta\Delta Ct}$  method (Livak and Schmittgen, 2001).

## Results

### Identification of SIPIN Gene Family in Tomato

Based on gene uniqueness, a total of ten SIPIN genes were identified and analysed, these genes are listed in Table 2. The 10 PIN genes were orderly named from SIPIN1 to SIPIN10 in the later work. The longest gene for encoding amino acids is SIPIN4, 653 amino acids (aa). The shortest gene for encoding amino acids is SIPIN10, 347 aa, with an average length of 527 aa. The molecular weights of SIPIN proteins ranged from 38411.53 (SIPIN10) Da to 71197.03 (SIPIN4) Da, and the variation range isoelectric points was 6.76 to 9.19. The majority of amino acids were basic amino acids; only SIPIN3, 4, and 10 proteins were acidic amino acids. The gene structure was highly conserved for most SIPINs. Six of 10 SIPINs contain 6 exons. These conserved exon structures of PIN genes were also found in other land plant species (Křeček *et al.*, 2009; Forestan *et al.*, 2012; Wang *et al.*, 2015). In addition, the structural change of the N-terminus was

stable, and the carboxyl terminal of the PIN protein changed greatly. Analysis of the instability coefficient of the coding protein showed that SIPIN2 and SIPIN4 proteins are instability proteins (instability coefficient >40); the other 8 SIPIN proteins are stable proteins. It was found that SIPIN5, 8 and 10 were hydrophobic proteins, as their grand averages of hydropathicity were more than 100. SignalP online tool reveal all of the identified SIPIN proteins that had no signal peptide were non-secreted proteins.

### The Secondary Structure of the PIN Protein Family Members in Tomato

Similar to other plant PINs proteins, SIPIN proteins have a typical conserved transmembrane domain, and the 10 SIPIN proteins are composed of two hydrophobic regions connected by a central hydrophilic loop at both ends of the protein. At the N-terminus, all SIPINs have 4 to 5 transmembrane helices, except for SIPIN7 and SIPIN8. The C-terminus end has 3 to 4 transmembrane helices (Zhao *et al.*, 2017). The 10 SIPIN proteins consisted of an alpha helix, turn, coil and beta sheet, and the ratio of the turn is smaller (Table 3). There are significant similarities (Coil > Alpha helix > Beta sheet > Turn) in the secondary structure of the analysed proteins except for that of SIPIN9 (Coil > Beta sheet > Alpha helix > Turn). SIPIN8 and SIPIN10 have the same structure characteristics (Alpha helix > Coil > Beta sheet > Turn). The secondary structure of proteins determines the folding and spatial configurations to some extent. It is of great significance to understand the structure and function of proteins by analysing and predicting their secondary structure.

### Phylogenetic Analysis of SIPIN Genes

To investigate evolutionary relationships among PINs in tomato and other species, 41 PIN protein sequences from tomato, Arabidopsis, rice and sorghum were comprehensive aligned by ClustalX and the MEGA 5.2 Software (Fig. 1). According to the difference of the length central hydrophilic loop, the SIPIN proteins were divided into two groups (Křeček *et al.*, 2009; Shen *et al.*, 2010): long PINs (SIPIN1, 2, 3, 4, 7 and 9) and short PINs (SIPIN5, 6, 8 and 10). The fact that SIPIN7 and SIPIN9 are very similar in sequence level and are located on the same branch indicates that they come from a common ancestor. SIPIN3, SIPIN4 and SIPIN5, SIPIN10 are also clustered on the same branch respectively. Meanwhile, SIPIN1 and AtPIN1, SIPIN8 and AtPIN8, SIPIN6 and AtPIN6 are clustered on the same branch respectively. In addition, the number of PIN genes in tomato, Arabidopsis, rice and sorghum was almost similar.

### Analysis of PIN Gene Structure

The multiple sequence alignment indicated that the sequences of transmembrane helices in SIPIN proteins were highly conserved at both ends of the protein and that the central hydrophilic loop was high heterogeneity (Fig. 2).

**Table 1:** Primers for qRT-PCR

Primer name	Squence (5'-3')	Primer name	Squence(5'-3')
SIPIN1-F	GCTGCAGGCTGGTCTAGATT	SIPIN1-R	AACAATGGCAACAAAGCACA
SIPIN3-F	GTGGGAACACTGTGGCTACT	SIPIN3-R	TGCATTGGCCTAATACATCTCTA
SIPIN4-F	TGCTTCAATTGCTGTTGGGC	SIPIN4-R	TTGAACCAACAATTTTAATGCAACA
SIPIN5-F	ACATTGAGCTGGCATTTTTGG	SIPIN5-R	TCCACTACCAGCCTTTTGACA
SIPIN6-F	GCAGCTCTCCCAAGGAAT	SIPIN6-R	GCGAAGACAAAATGGAACGAT
SIPIN7-F	CATCAGCGGTCCAGCAGTCA	SIPIN7-R	TGTTTCCGAAGGGTCTCAGTT
SIPIN8-F	CAGCCCTTCCCAAGGAATC	SIPIN8-R	CCCTGCAATCAGAATGAAAACA
SIPIN9-F	TCTTTTAGGTGGAATGTTCAAATGC	SIPIN9-R	CGACAATGCCATGAACAAACC
SIPIN10-F	AGAGCATGTTTGGCTCAGCTT	SIPIN10-R	ACCCCTACCCAGCTTTTAAC
Slaction-F	GAAATAGCATAAGATGGCAGACG	Slaction-R	ATACCCACCATCACACCAAGTAT

**Table 2:** The distribution of PIN gene family members on the scaffolds of the genome and physico-chemical analysis in tomato

Gene name	Locus name	Amino acids (No.)	Exons (No.)	Molecular weight (Da)	Theoretical PI	Instability index (II)	Grand hydrophaticity (GRAVY)	Average Aliphatic Index (AI)
SIPIN1	Solyc03g118740.2.1	611	6	66938	9.09	38.93	0.038	87.15
SIPIN2	Solyc07g006900.1.1	631	7	68565.38	9.19	44.16	0.180	97.34
SIPIN3	Solyc04g007690.2.1	613	6	67612.20	6.76	37.33	0.151	93.83
SIPIN4	Solyc05g008060.2.1	653	6	71197.03	6.98	40.74	0.073	89.62
SIPIN5	Solyc01g068410.2.1	358	5	39673.22	8.80	34.09	0.732	120.67
SIPIN6	Solyc06g059730.1.1	521	7	56876.39	8.61	34.08	0.379	105.71
SIPIN7	Solyc10g080880.1.1	586	6	63858.08	8.93	34.49	0.240	97.03
SIPIN8	Solyc02g087660.2.1	357	6	38966.79	8.89	35.40	0.778	136.30
SIPIN9	Solyc10g078370.1.1	594	6	64304.33	9.16	33.22	0.195	95.54
SIPIN10	Solyc04g056620.1.1	347	5	38411.53	6.99	33.37	0.698	111.01

**Table 3:** The secondary structure of PIN protein family members in tomato

Gene	Alpha helix	tum	coil	Beta sheet
SIPIN1	193 (31.59%)	70 (11.46%)	222 (36.33%)	126 (20.62%)
SIPIN2	204 (32.33%)	72 (11.41%)	219 (34.71%)	136 (21.55%)
SIPIN3	167 (27.24%)	63 (10.28%)	234 (38.17%)	149 (24.31%)
SIPIN4	177 (27.11%)	69 (10.57%)	259 (39.66%)	148 (22.66%)
SIPIN5	224 (31.11%)	89 (12.36%)	227 (31.53%)	180 (25.00%)
SIPIN6	158 (30.38%)	56 (10.77%)	171 (32.88%)	135 (25.96%)
SIPIN7	183 (31.23%)	73 (12.46%)	206 (35.15%)	124 (21.16%)
SIPIN8	143 (40.06%)	32 (8.96%)	94 (26.33%)	88 (24.65%)
SIPIN9	147 (24.75%)	69 (11.62%)	223 (37.54%)	155 (26.09%)
SIPIN10	159 (45.82%)	26 (7.49%)	84 (24.21%)	78 (22.48%)

Similar to the AtPIN (Paponov *et al.*, 2005), OsPIN (Wang *et al.*, 2009) and SbPIN (Shen *et al.*, 2010) proteins, the central hydrophilic loop consists of C1, C2, C3 structural domains and two variable regions (V1, V2) for long PINs. The hydrophilic region of the short PINs group has only one constant C1 region and a variable V1 region. The length of the central hydrophilic loop is approximately 350 amino acids for members of the long PINs. However, the length of the short PINs members is 47–243 amino acids. The conserved structure NPNXY is found between the hydrophilic loop and the C-terminal hydrophobic domain of all SIPIN proteins (except SIPIN8). The last one amino acid of SIPIN8 is histidine rather than tyrosine. NPXXY plays an important role in clathrin dependent endocytosis (Yu and Cui, 2009). The difference in length between the proteins is the result of the difference in length of the hydrophilic region located between the transmembrane helice domains present at C- and N-terminal of the protein. Most SIPINs contain two phosphorylation active sites that are marked in the figure by


black triangles. These sites can be phosphorylated by serine/threonine protein kinases (Křeček *et al.*, 2009).

### Motifs Analysis and Chromosome Mapping of the SIPINs

To further study the multiformity of SIPINs, conserved motifs in PIN genes were identified by MEME software. Based on the distribution of the 8 predicted motifs, the tomato PIN genes were also categorized into two groups based on the distribution of the 8 predicted motifs, which was consistent with the phylogenetic analysis. The essential information of the logo and sequence of each motif are shown in Fig. 3 and Table 4. We found that the tomato PIN gene family members not only have a typical conserved motif but also have some relatively conserved motifs that consist of 29–50 amino acid residues. We also found that different clades shared similar motifs: 8 motifs were distributed in the long PINs, and 6 motifs were in the short group. Motifs 1, 2, 3, and 4 encoded PIN domains in all the studied genes.

**Table 4:** The conserved motif of PIN protein family from tomato

Motif	Width	Best possible match
1	50	PLYVAMILAYGSVKWWKIFSPDQCSGINRFVALFAVPLLSFHFASNNPY
2	49	LLHVAIVQAALPQGIVPFVFAKEYNVHPDILSTAVIFGMLIALPITLVYY
3	49	RKLIRNPNTYSSLJGLIWSLISFRWNVQMPKIIIEKSISILSDAGLGMAMF
4	50	GSLEWSITLFLSLTLPNTLVMGIPLLKAMYGDYSGSLMVQIVVLQCIWY
5	41	FMALQPKIIACGKSVATFAMA VRFLTGPVMAAAASIAVGLR
6	42	DTAGSIVSFKVDSVIVSLDGREPLETDAEIGDDGKLVHTVRK
7	41	MTPRPSNLTGAEIYSLQSSRNPTPRGSNFNHTDFYSMVGGK
8	29	MNYRFIAADTLQKVIVLFLVLAIWANVSKR



**Fig. 1:** Phylogenetic relationships of PIN auxin transporters from tomato (SI), *A. thaliana* (At), Rice (Os) and Sorghum (Sb). Triangles, squares, circles and rhombi represent the PINs of Rice, Sorghum, *A. thaliana* and tomato, respectively

The internalizational NPXXY structure is located within motif 3. The N-terminus of the SIPIN protein family has a conserved motif 1. Motif 2 was present in the C-terminus of all members, and the rest motifs were unevenly distributed in two groups.

Chromosome map positions of tomato PINs that were identified are reported in Fig. 4. Our results showed that 10 SIPIN genes were unevenly distributed on eight chromosomes in the whole tomato genome. The number of SIPIN genes per chromosome ranged from zero to two, with zero genes on chromosomes 8, 9, 11 and 12; one gene each on chromosomes 1, 2, 3, 5, 6 and 7; and two genes on chromosomes 4 and 10. In tandem replication, the distances of adjacent genes on the same chromosome were less than 100 kb, thus, tandem replication did not occur.

**Expression of SIPIN Genes under Abiotic Stresses Treatment**

To understand the expression patterns of SIPIN genes under

different stresses conditions, the expression profiles of 9 selected SIPIN genes (excluding SIPIN2) were studied in response to drought, salt and cold treatments by qRT-PCR experiments based on tissue-specific expression patterns of SIPIN genes. A histogram representation for transcript expression fold changes in response to abiotic stresses is shown in Fig. 4. The expression level of the PIN genes was affected under one or more treatments. More SIPINs were up-regulated in response to drought compared with cold and salt stress.

Gene expression was slightly induced in tomato plants treated with salt (Fig. 5a). Expression of SIPIN3 decreased over time. SIPIN1, 5, and 7 decreased first but then increased. The expression of SIPIN1 and 7 reached the minimum value at 4 h, and SIPIN8 reached the minimum value at 8 h. SIPIN4 significantly decreased compared to the high expression at the normal 0-h point. The change patterns of SIPIN6 and 10 were consistent, as both reached their maximum at the 12-h point. The relative expression of partial PIN genes were remarked by cold stress (Fig. 5b). The expression of SIPIN7, 8 and 10 increased only at 8 h; nevertheless, their expression was low at the other time points. The relative expression of PIN genes SIPIN3 and 6 increased rapidly at the 2 h post-stimulus point and showed a tendency of increasing first and then decreasing. SIPIN1 was up-regulated compared to the low expression at the normal 0 h point, whereas the SIPIN genes were moderately induced. Under drought stress, all PIN genes possessed different expression patterns (Fig. 5c). The expression of SIPIN1, 6, 8 and 9 was up-regulated at first but then decreased after 8 h. SIPIN10 was also up-regulated over time but decreased at the 12 h point. The expression of SIPIN4 and 5 increased at 4, 6 and 12 h. The expression of SIPIN7 increased at the 4 h point, and then held steady, with moderate changes.


**Discussion**

Based on the current tomato genome sequence and the SGN database, we identified 10 PIN genes and synthetically analysed the SIPIN family of the tomato genome. Our results showed that 10 SIPIN proteins are stable proteins, most of which are composed of basic amino acids, and are non-secreted proteins. However, the protein length of SIPIN3 and 6 and the exon numbers of SIPIN2 were different from those reported in previous studies (Zhao *et al.*, 2017),






**Fig. 2:** Multiple sequence alignment of the SIPIN gene family. Two hydrophobic domains in the SIPIN proteins are underlined with solid blue boxes, while the hydrophilic loop region is underlined with a solid red box. The predicted transmembrane helix-formed regions in the primary structure of SIPIN proteins are marked. The possible phosphorylation sites are marked with triangles. The NPXXY structure is represented with a green ellipse




**Fig. 3:** Schematic distribution of conserved motifs in the SIPIN proteins. The distribution of conserved motifs in tomato identified using the MEME program is shown. Eight conserved motifs are shaded in different colours. Details of the individual motifs are shown in Supplementary Table 4. Among them, motif 3 contains the conserved structure NPXXY

which may be due to differences in information collected by different databases. The secondary structure of SIPIN proteins was highly similar. It is speculated that the SIPIN genes have a similar function. The function of genes can be inferred according to the phylogenetic relationships among homologues from different species, thus Arabidopsis, sorghum and rice PIN protein families were

selected as reference sequences. Compared with rice and sorghum, the evolutionary relationship between tomato and *A. thaliana* was relatively closer, which may be because they are dicotyledons. The cluster analysis of proteins based on similarity between AtPIN and SIPIN genes sequence indicates a similar function in the species-dependent developmental processes.



**Fig. 4:** Chromosomal distributions of the identified tomato PIN genes. Chromosomal positions of the PIN genes in tomato were mapped on the basis of the SGN tomato genome database. There were no PIN genes distributed on tomato chromosomes 8, 9, 11, or 12. The scale is according to physical position (Mb)



**Fig. 5:** Expression profiles of SIPIN genes in response to different abiotic stresses. The relative expression of SIPIN genes was analysed using qRT-PCR. The Y axis is the scale of the relative expression level; the X axis is the time course of cold stress treatment. The statistical bars indicate the relative expression of SIPIN genes in different treatments. (a) shows the expression levels of SIPIN genes under salt treatment. (b) shows the expression levels of SIPIN genes under cold treatment. (c) shows the expression levels of SIPIN genes under drought treatment

In agreement with the very recent study of Mounet (Mounet *et al.*, 2012). AtPIN1 is located in root and shoot vascular tissues and embryos. This gene is involved in the regulation of the development of various organs that are involved with

roots and tillers. It is speculated that SIPIN1 may take on these functions. At present, the functions of AtPIN6 and AtPIN8 are not very clear; many experiments are needed to further verify and analyse these functions. Subsequent studies

can refer to the expression and function of *Arabidopsis* PIN proteins and carry out the related research of tomato PIN proteins. Motif analysis showed that motifs 1, 2, 3 and 4 were present in all the members of SIPIN. In addition, different groups shared similar motifs, indicating that these conserved motifs might play vital roles in specific functions. Alignment of amino acid sequences of SIPIN proteins showed that the N-terminal and C-terminal regions were highly conserved, divided by the hydrophilic loop. The PIN protein hydrophilic loop can modulate intracellular auxin homeostasis, which is dependent on cell type and developmental stage (Ganguly *et al.*, 2014; Yue *et al.*, 2015). The relatively high amino acid identity between PIN proteins indicated that all the PIN genes evolved from a common ancestral sequence. Both the conserved domain and variable domain of the SIPINs may determine the specific function of these genes. These bioinformatic analysis are useful for studying the biological functions of genes.

Auxin transport plays important roles in plant growth and development by controlling a large number of auxin-responsive genes (AUX/LAX, ABCB, PIN and PILS) (Zhang *et al.*, 2014). It has been studied that various abiotic treatments can alter auxin polarity distribution by modulating PIN protein (Friml, 2010). At the same time, PIN families are regulated under various abiotic stresses, including dehydration drought, low temperature, high temperature, salinity, and hormones in the leaves (Ranawake *et al.*, 2012) and roots (Yue *et al.*, 2015). However, there are few investigations of tomato PIN genes under abiotic stress. Therefore, we investigated the change of relative expression of SIPIN genes under abiotic stress using the above analysis as a foundation. The relative expression of SIPIN4 decreased rapidly at the 2 h point after salt treatment, which indicated that SIPIN4 might be involved in the mechanism of salt tolerance in tomato. Only the expression of SIPIN3 decreased gradually under salt stress treatment without any rebound in this study, but the range of overall change of this gene was small. It has recently been reported that salt stress causes severe yield loss in salt-affected areas (Guan *et al.*, 2014) and promotes auxin accumulation in the developing primordia of *Arabidopsis*. Low temperature has different effects on plant growth, photosynthesis, dry matter accumulation and carbon metabolism. The results suggested that the relative expression of most the SIPIN genes peaked at 8 h after cold treatment, except for SIPIN3. SIPIN3 and SIPIN6 were greatly affected by cold stress, which implied that these SIPIN genes play a part in the mechanism that helps tomato plants tolerate cold stress. Kyohei Shibasaki (Shibasaki *et al.*, 2009) also showed that cold stress influences the polar transport of auxin through selectively inhibiting the intracellular trafficking of proteins, including auxin influx carriers and efflux carriers, and indicated that the AtPIN3 proteins might be responsible for the inhibited auxin polar transport under cold stress conditions. The expression levels of SIPIN7, 8 and 10 were low at the any time points, except for 8 h point. Their more distinguishingly fluctuations were

complex and should be considered as a basic information for further investigation. In maize, the expression of most ZmPIN genes was up-regulated by drought in the shoots but was down-regulated in the roots. Generally in a tissue-specific manner, many soybean PIN genes were responsive to drought conditions at the transcriptional level, at different degrees of stress. Our data showed that the relative expression level of all SIPIN genes was enhanced compared with that of untreated plants under drought conditions. In particular, the expression of SIPIN1 and SIPIN6 varied greatly under drought stress. It is suggested that tomato responds to drought stress through complicated network, which necessitates the mediated regulation of most SIPINs. It is important that the relative expression of SIPIN6 significantly increased under cold and drought stress, as SbPIN6 was inhibited by salt and drought treatments. These results show that SIPIN6 were obvious relative to cold and drought. The molecular mechanism of SIPIN6 will be studied further. Candidate genes would provide a useful reference for further functional investigation of PIN gene family in *Solanaceae* crops.

## Conclusion

In this study, we identified total 10 PIN genes in the genome of tomato and classified them into two groups ("short" and "long" PINs), according to the known *Arabidopsis thaliana*, rice and sorghum PIN protein family as reference sequences, and the results of the motif analysis are in agreement with the phylogenetic tree analysis results. There are significant similarities in the secondary structure of the 10 PIN members. Alignment of amino acid sequences showed that the tomato PIN domain were highly conserved. Chromosome map positions showed that 10 SIPIN genes were unevenly distributed on eight chromosomes. In addition, the expression levels of SIPIN family members were affected by salt, drought and cold stresses to different degrees. More SIPIN genes were involved in response to drought and cold than to salt stress. They might participate in the response to environmental stresses. These results provide fundamental information for further exploration of PIN genes in tomato.

## Acknowledgements

This work was supported by the National Key R & D Program of China (2017YED0101900), The China Agriculture Research System (CARS-23-A-16), Natural Science Foundation of Heilongjiang Province (C2017024) and Youth Talent support program of Northeast Agricultural University (17QC07).

## References

- Barbez, E., M. Kubeš, J. Rolčík, C. Béziat, A. Pěnčík, B. Wang, M.R. Rosquete, J.S. Zhu, P.I. Dobrev, Y. Lee, E. Zažímalová, J. Petrášek, M. Geisler, J. Friml and J. Kleine-Vehn, 2012. A novel putative auxin carrier family regulates intracellular auxin homeostasis in plants. *Nature*, 485: 119–122



- Cazzonelli, C.I., M. Vanstraelen, S. Simon, K. Yin, A. Carron-Arthur, N. Nisar, G. Tarle, A.J. Cuttriss, I.R. Searle, E. Benkova, U. Mathesius, J. Masle, J. Friml and B.J. Pogson, 2013. Role of the Arabidopsis PIN6 auxin transporter in auxin homeostasis and auxin-mediated development. *Plos One*, 8: 1–14
- Carraro, N., C. Forestan, S. Canova, J. Traas and S. Varotto, 2006. ZmPIN1a and ZmPIN1b encode two novel putative candidates for polar auxin transport and plant architecture determination of maize. *Plant Physiol.*, 142: 254–264
- Chai, C.L. and P.K. Subudhi, 2016. Comprehensive analysis and expression profiling of the OsLAX and OsABC B auxin transporter gene families in rice under phytohormone stimuli and abiotic stresses. *Plant Sci.*, 7: 593
- Consortium, T.T.G., 2012. The tomato genome sequence provides insights into fleshy fruit evolution. *Nature*, 485: 635–641
- Ding, Z.J., B. Wang, I. Moreno, N. Duplakova, S. Simon, N. Carraro, J. Reemmer, A. Pěnčík, X. Chen, R. Tejos, P. Skůpa, S. Pollmann, J. Mravec, J. Petrášek, E. Zažímalová, D. Honys, J. Rolčík, A. Murphy, A. Orellana, M. Geisler and J. Friml, 2012. ER-localized auxin transporter PIN8 regulates auxin homeostasis and male gametophyte development in Arabidopsis. *Nat. Commun.*, 3: 941–942
- Dubrovsky, J.G., M. Sauer, S. Napsucially-Mendivil, M.G. Ivanchenko, J. Friml, S. Shishkova, J. Celenza and E. Benková, 2008. Auxin acts as a local morphogenetic trigger to specify lateral root founder cells. *Proc. Natl. Acad. Sci. USA*, 105: 8790–8794
- Feraru, E., S. Vosolobě, M.I. Feraru, J. Petrášek and J. Kleine-Vehn, 2012. Evolution and structural diversification of PILS putative auxin carriers in plants. *Front. Plant Sci.*, 3: 1–13
- Fleet, C.M. and T.P. Sun, 2005. A DELLAcate balance: the role of gibberellin in plant morphogenesis. *Curr. Opin. Plant Biol.*, 8: 77–85
- Forestan, C., S. Farinati and S. Varotto, 2012. The maize PIN gene family of auxin transporters. *Front. Plant Sci.*, 3: 1–16
- Friml, J., 2010. Subcellular trafficking of PIN auxin efflux carriers in auxin transport. *Eur. J. Cell. Biol.*, 89: 231–235
- Gälweiler, L., C. Guan, A. Müller, E. Wisman, K. Mendgen, A. Yephremov and K. Palme, 1998. Regulation of polar auxin transport by AtPIN1 in Arabidopsis vascular tissue. *Science*, 282: 2226–2230
- Ganguly, A., M. Park, M.S. Kesawat and H.T. Cho, 2014. Functional analysis of the hydrophilic loop in intracellular trafficking of Arabidopsis PIN-FORMED proteins. *Plant Cell*, 26: 1570–1585
- Goto, N., M. Starke and A.R. Kranz, 1987. Effect of gibberellins on flower development of the pin-formed mutant of *Arabidopsis thaliana*. *Arabidopsis Inf. Ser.*, 23: 66–71
- Guan, R.X., Y. Qu, Y. Guo, L.L. Yu, Y. Liu, J. Jiang, Y.L. Ren, G.Y. Liu, L. Tian, L.G. Jin, Z.X. Liu, H.L. Hong, R.Z. Chang, M. Gilliam and L.J. Qiu, 2014. Salinity tolerance in soybean is modulated by natural variation in GmSALT3. *Plant J.*, 80: 937–950
- Křeček, P., P. Skůpa, J. Libus, S. Naramoto, R. Tejos, J. Friml and E. Zažímalová, 2009. The Pinformed (PIN) protein family of auxin transporters. *Genom. Biol.*, 10: 249–260
- Liu, C., A. Chang and C. Du, 2015. Genetic, physiological and biochemical analysis of the formation of yellow-green leaf color of burley tobacco (*Nicotiana tabacum*). *Int. J. Agric. Biol.*, 17: 767–772
- Li, Y.H., M.H. Zou, B.H. Feng, X. Huang, Z. Zhang and G.M. Sun, 2012. Molecular cloning and characterization of the genes encoding an auxin efflux carrier and the auxin influx carriers associated with the adventitious root formation in mango (*Mangifera indica* L.) cotyledon segments. *Plant Physiol. Biochem.*, 55: 33–42
- Livak, K.J. and T.D. Schmittgen, 2001. Analysis of relative gene expression data using realtime quantitative PCR and the  $2^{-\Delta\Delta CT}$  method. *Methods*, 25: 402–408
- Ma, H. and Z. Jie, 2010. Genome-wide identification classification and expression analysis of the arabinogalactan protein gene family in rice. *J. Exp. Bot.*, 61: 2647–2668
- Mounet, F., A. Moing, M. Kowalczyk, J. Rohrmann, J. Petit, V. Garcia, M. Maucourt, K. Yano, C. Deborde, K. Aoki, H. Bergès, A. Granell, A.R. Fernie, C. Bellini, C. Rothan and M. Lemaire-Chamley, 2012. Down-regulation of a single auxin efflux transport protein in tomato induces precocious fruit development. *J. Exp. Bot.*, 63: 4901–4917
- Mravec, J., P. Skůpa, A. Bailly, K. Hoyerová, P. Křeček, A. Bielach J. Petrášek, J. Zhang, V. Gaykova, Y.D. Stierhof, P.I. Dobrev, K. Schwarzerová, J. Rolčík, D. Seifertová, C. Luschnig, E. Benková, E. Zažímalová, M. Geisler and J. Friml, 2009. Subcellular homeostasis of phytohormone auxin is mediated by the ER-localized PIN5 transporter. *Nature*, 459: 1136–1140
- Ni, W.M., X.Y. Chen, Z.H. Xu and H.W. Xue, 2002. A Pin gene families encoding components of auxin efflux carriers in *Brassica juncea*. *Cell Res.*, 12: 247–255
- Nishio, S., R. Moriguchi, H. Ikeda, H. Takahashi, N. Fujii, T.J. Guilfoyle, K. Kanahama and Y. Kanayama, 2010. Expression analysis of the auxin efflux carrier family in tomato fruit development. *Plantarum*, 232: 755–764
- Paponov, I.A., W.D. Teale, M. Trebar, I. Blihou and K. Palme, 2005. The PIN auxin efflux facilitators: evolutionary and functional perspectives. *Trends Plant Sci.*, 10: 170–177
- Pattison, R.J. and C. Catalá, 2012. Evaluating auxin distribution in tomato (*Solanum lycopersicum*) through an analysis of the PIN and AUX/LAX gene families. *Plant J.*, 70: 585–598
- Ranawake, A.L., N. Mori and C. Nakamura, 2012. Expression profiles of stress responsive genes in rice (*Oryza sativa* L.) under abiotic stresses. *Biotechnol. Eq.*, 26: 2838–2843
- Robert, H.S., P. Grones, A.N. Stepanova, L.M. Robles, A.S. Lokerse, J.M. Alonso, D. Weijers and J. Friml, 2013. Local auxin sources orient the apical-basal axis in *Arabidopsis* embryos. *Curr. Biol. Cb.*, 23: 2506–2512
- Roumeliotis, E., B. Kloosterman, M. Oortwijn, R.G.F. Visser and C.W.B. Bachem, 2013. The PIN family of proteins in potato and their putative role in tuberization. *Front. Plant Sci.*, 4: 524–531
- Růžicka, K., M. Šimásková, J. Duclercq, J. Petrášek, E. Zažímalová, S. Simon, J. Friml, M.C. Van and E. Benková, 2009. Cytokinin regulates root meristem activity via modulation of the polar auxin transport. *P. Natl. Acad. Sci. USA*, 106: 4284–4289
- Schnabel, E.L. and J. Frugoli, 2004. The PIN and LAX families of auxin transport genes in *Medicago truncatula*. *Mol. Genet. Genom.*, 272: 420–432
- Shen, C.J., Y.H. Bai, S.K. Wang, S.N. Zhang, Y.R. Wu, M. Chen, D.A. Jiang and Y.H. Qi, 2010. Expression profile of PIN AUX/LAX and PGP auxin transporter gene families in Sorghum bicolor under phytohormone and abiotic stress. *FEBS J.*, 277: 2954–2968
- Shibasaki, K., M. Uemura, S.J. Tsurumi and A. Rahman, 2009. Auxin response in arabidopsis under cold stress: underlying molecular mechanisms. *Plant Cell*, 21: 3823–3838
- Sun, X., T. Lei, J.B. Du and W.Y. Yang, 2015. Identification and characterization of two paralogous plastid terminal oxidase genes in soybean. *Int. J. Agric. Biol.*, 17: 1275–1278
- Tamura, K., D. Peterson, N. Peterson, G. Stecher, M. Nei and S. Kumar, 2011. MEGA5: Molecular evolutionary genetics analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. *Mol. Biol. Evol.*, 28: 2731–2739
- Wang, Y.Q., C.L. Chai, B.B. Valliyodan, C. Maupin, B. Annen and H.T. Nguyen, 2015. Genome-wide analysis and expression profiling of the PIN auxin transporter gene family in soybean (*Glycine max*). *BMC. Genom.*, 16: 951
- Wang, J.R., H. Hu, G.H. Wang, J. Li, J.Y. Chen and P. Wu, 2009. Expression of PIN genes in rice (*Oryza sativa* L.): tissue specificity and regulation by hormones. *Mol. Plant*, 2: 823–831
- Xi, W.Y., X.M. Gong, Q.Y. Yang, H. Yu and Y.C. Liou, 2016. Pin 1 At regulates PIN1 polar localization and root gravitropism. *Nat. Commun.*, 7: 10430
- Yu, S.N. and J.Z. Cui, 2009. Role of PIN proteins in polar auxin transport. *Biotechnol. Bull.*, 3: 20–24
- Yue, R.Q., S.G. Tie, T. Sun, L. Zhang, Y.J. Yang, J.S. Qi, S.F. Yan, X.H. Han, H.Z. Wang and C.J. Shen, 2015. Genome-wide identification and expression profiling analysis of ZmPIN, ZmPILS, ZmLAX and ZmABC B auxin transporter gene families in maize (*Zea mays* L.) under various abiotic stresses. *Plos One*, 10: 276–289

- Zhao, R.R., L. Shen and J.P. Sheng, 2017. Bioinformatic analysis of the PIN gene family in tomato and its expression pattern. *Food Sci.*, 38: 1–5
- Zhao, Y., 2010. Auxin biosynthesis and its role in plant development. *Annu. Rev. Plant Biol.*, 61: 49–64
- Zhang, Z.Z., C.L. Chen, X. Guan, Y. Liu, H.Y. Chen, T.T. Wang, L.D.O. Mouekouba, J.F. Li and A.X. Wang, 2014. A genome-wide survey of homeodomain-leucine zipper genes and analysis of cold-responsive HD-Zip I members expression in tomato. *Biosci. Biotech. Biochem.*, 78: 1337–1349
- Zhang, Y.X., A. Paschold, C. Marcon, S.Z. Liu, H.H. Tai, J. Nestler, C.T. Yel, N. Opitz, C. Lanz, P.S. Schnable and F. Hochholdinger, 2014. The Aux/IAA gene *rum1* involved in seminal and lateral root formation controls vascular patterning in maize (*Zea mays* L.) primary roots. *J. Exp. Bot.*, 65: 4919–4930

(Received 23 January 2018; Accepted 30 May 2018)

IB Press