



Full Length Article

Influence of Seed Priming and Sowing Methods on Growth and Productivity of Wheat Cultivars Differing in Seed Size in Rice-Wheat Cropping system of Punjab, Pakistan

Wardah Muzaffar¹, Riaz Ahmad^{1*}, Muzzammil Hussain³, Muhammad Farooq² and Abdul Wakeel⁴

¹Department of Agronomy, Faculty of Agriculture, University of Agriculture, Faisalabad 38000, Pakistan

²College of Agricultural and Marine Sciences, Sultan Qaboos University, P.O. Box 34, AlKhoud 123, Oman

³Senior Subject Matter Specialist, Adaptive Research Farm, Gujranwala 52250, Pakistan

⁴Department of Soil Science, University of Agriculture, Faisalabad 38000, Pakistan

*For correspondence: riazahmadagr@gmail.com

Abstract

Poor and non-uniform stand establishment causes significant yield loss in wheat sown after puddled rice in rice-wheat cropping system. Such losses can be overcome by sowing of wheat with appropriate cultivar after seed priming by conservation tillage practices. Field experiment were conducted during 2016-2017 and 2017-2018 comprising of three different cultivars *i.e.*, AS-2002, FSD-2008, Lasani-2008 having bold, medium and small seed size, respectively, two priming techniques *i.e.*, hydropriming, osmopriming (1.2%) and dry seeds (control) and three different sowing methods *i.e.*, plough tillage, happy seeder (zero tillage) and turbo seeder. Results showed that wheat planted with conservation tillage using happy seeder improved the grain yield and other parameters. Bold seeded variety 'AS-2002' had better stand establishment, improved yield related traits and grain yield of wheat than small and medium grain varieties. Crop sown with happy seeder using hydroprimed seeds completed its maturity earlier through rapid phenological developments. Hydropriming of medium-size seed variety (FSD-2008) sown using happy seeder produced maximum economic return followed by osmoprimed seed of Lasani-2008 (small seed size) using happy seeder. In conclusion, sowing of medium seed size cultivar after seed priming (hydropriming/osmopriming) using happy seeder improved the productivity of wheat sown in conventional rice-wheat system. © 2019 Friends Science Publishers

Keywords: Seed size; Seed priming; Conservation tillage; Rice-wheat cropping systems

Introduction

In Asian subtropics, rice-wheat is the most important cropping system; and farmers in Pakistan grow wheat (*Triticum aestivum* L.) followed by rice (*Oryza sativa* L.) in this system (Ahmad *et al.*, 2013). Rice is grown in puddled fields thus formation of hardpan due to excessive use of machinery for puddling results in poor root development in succeeding wheat crop thus reduce the nutrient and water uptake from soil (Farooq *et al.*, 2011; Jabran *et al.*, 2015). Due to soil compaction, leaching and de-nitrification increase which ultimately leads towards the low availability of nitrogen (N) to plant (Fugelsnes and Lie, 2011). Moreover, late harvesting of rice and crop residues management is the major problem for timely preparation of seedbed of following wheat crop that delay timely wheat sowing resulting in low productivity of wheat (Ahmad and Meisner, 1996). These problems may be solved by sowing wheat with zero tillage (ZT). ZT technology used for sowing of wheat crop can help to enhance the ultimate yield, increase input use efficiency and finally the net income of growers (Gupta and Seth, 2007; Erenstein *et al.*, 2008; Shahzad *et al.*, 2017). Zero tillage in

wheat is helpful in timely sowing, reduces cost of production/tillage cost (Erenstein and Laxmi, 2008) and improves soil fertility status (Mohanty *et al.*, 2007). Under ZT, wheat is usually sown using different equipments like zone disk tiller machine, happy seeder, turbo seeder and tine openers (Baker *et al.*, 1996). Lampurlaneas *et al.* (2001) found that ZT facilitated water storage in soil and root growth. For sustaining agronomic production soil physical environment is important. Experimental results showed that soil physical properties are mainly affected by tillage (Rashidi and Keshavarzpour, 2008). Zero tillage changed the physical properties of soil depending on the climate and adaptation factors (Martinez *et al.*, 2008). In zero tillage more drainage, macro pores and higher soil water movement than conventional was recorded. The minimum disturbance of soil results better aggregate stability more infiltration rate of water include bulk density and root penetration resistance compared to conventional system of sowing of wheat with normal ploughing practices (Alvarez and Steinbach, 2009). Similarly, the Karina *et al.* (2009) suggested that ZT reduce damaging effects of soil erosion as compare to the conventional system of tillage.

Therefore, the OM content and pore spaces in the upper soil layers were improved using long term zero tillage system.

Seed selection plays major role in determining the performance and productivity of crop. Moreover, seed quality is affected by many factors *e.g.*, genetic makeup of variety, seed viability and seed vigor, and moisture percentage in grains (Akbari *et al.*, 2004). Hence, seed weight substantially affects the germination, vigor and seedling establishment of crop. Seeds with higher weight may have high germination under field condition (Gorge and Ray, 2005). Size of seed may be positively linked with seed vigor as bold seed produce vigorous seedlings under field conditions.

Effectiveness of seed priming techniques in improving stand establishment and productivity of wheat and many other crops has been reported (Harris *et al.*, 2007; Hussain *et al.*, 2016; Farooq *et al.*, 2017). Seed priming is an economical technology which may improve early growth of crop resulting in better stand establishment and ultimately yield related benefits especially in wheat crop (Rehman *et al.*, 2011; Hussain *et al.*, 2017). In hydropriming seeds are soaked in water and are dried before sowing to complete seed hydration. Aeration may or may not be provided to submerged seeds; whereas, osmopriming refers to soaking of seeds in aerated solutions with low water potential. Then the seeds are re-dried near to original weight after soaking with forced air under shade. It was observed that seeds treated with calcium chloride (CaCl₂) took lesser time for seed germination compared to untreated seeds (Khan *et al.*, 2014, 2015). The CaCl₂ priming also improved total sugars and non-reducing sugars in wheat plants (Afzal *et al.*, 2008; Khan *et al.*, 2014, 2015). Osmopriming with calcium salts is an effective method for enhancing germination rate and better establishment of seedlings in wheat (Hussain *et al.*, 2016; Farooq *et al.*, 2017).

The effect of seed size on the performance of wheat crop in improving yield has been reported earlier (Shahwani *et al.*, 2014; Haider *et al.*, 2016). However, there influence on the productivity of conventional and zero tillage wheat under rice-wheat cropping system are rarely investigated. So keeping this in view the present study was conducted to evaluate the impact of seed size, seed priming techniques along with planting methods on the productivity of wheat sown after puddled and aerobic rice systems.

Materials and Methods

Site, Soil and Climate

Two-year field experiment was carried out at the Adaptive Research Farm, Gujranwala, Punjab (latitude 32°N, longitude 74°E and altitude 226 m) during 2016-2017 and 2017-2018. Prior to experimentation, the soil samples (30 cm depth) were collected from the experimental site to determine the soil properties (Table 1).

Table 1: Post-experimental soil analysis of experimental site

Treatments			pH	EC	OM	N	P	K
			(dS m ⁻¹)		(mgkg ⁻¹)	(mg kg ⁻¹)	(mg kg ⁻¹)	(mgkg ⁻¹)
			7.8	0.93	8.9	0.53	5.0	140
Conventional Method	AS-2002	C	7.1	0.81	8.3	0.40	8.0	129
		H	7.3	0.80	8.5	0.38	9.1	128
		O	8.2	0.90	7.0	0.25	8.2	120
	FSD-2008	C	7.0	0.87	6.9	0.40	8.1	139
		H	6.9	0.87	7.9	0.39	7.3	133
		O	8.1	0.70	8.1	0.49	8.4	129
	Las-2008	C	7.9	0.81	7.2	0.38	6.7	121
		H	8.0	0.91	8.0	0.50	9.7	134
		O	8.2	0.87	8.8	0.49	7.5	110
Happy Seeder	AS-2002	C	7.4	0.80	8.3	0.35	7.9	125
		H	7.0	0.67	7.7	0.43	6.9	132
		O	7.5	0.77	8.0	0.38	7.1	121
	FSD-2008	C	7.8	0.58	8.3	0.50	8.1	134
		H	7.6	0.70	7.2	0.42	7.7	140
		O	8.1	0.70	6.8	0.43	6.9	133
	Las-2008	C	7.7	0.80	7.9	0.47	7.3	125
		H	8.0	0.9	8.0	0.61	6.7	129
		O	8.1	0.87	8.1	0.49	8.0	138
Turbo Seeder	AS-2002	C	8.2	0.70	7.3	0.50	9.1	110
		H	7.8	0.69	7.8	0.38	8.1	133
		O	7.7	0.71	7.6	0.33	8.3	127
	FSD-2008	C	7.9	0.78	8.0	0.45	7.2	134
		H	8.0	0.75	8.1	0.49	7.1	140
		O	8.1	0.81	8.0	0.37	8.2	138
	Las-2008	C	7.7	0.80	8.3	0.50	8.0	133
		H	8.3	0.9	7.9	0.41	7.7	125
		O	8.0	0.87	7.8	0.38	8.0	135

Seed Material and Priming

The experimental treatments were comprised of three seed sizes, *i.e.*, bold seed (AS-2002) (S₁), medium seed (FSD-2008) (S₂) and small seed (Lasani-2008) (S₃), (based on 1000-seed weight), three seed priming treatments, *i.e.*, control (no priming) (P₁), hydropriming (P₂) and osmopriming (P₃). Wheat was planted under different sowing methods including conventional method (plough tillage) (M₁), happy seeder (zero tillage) (M₂) and turbo seeder (zero tillage) (M₃).

Seeds of three different wheat cultivars *i.e.*, AS-2002, Faisalabad-2008, Lasani-2008 were obtained from the Wheat Research Institute, AARI, Faisalabad. For hydropriming, seeds were soaked directly in the distilled water provided with aeration. For osmopriming seeds were soaked in aerated solution of CaCl₂ (1.2%) for 10 h at 25±2°C. The ratio of seeds to solution was 1:5(w/v). After priming, seeds were surface washed with distilled water and re-dried under shade near to their original weight and then sealed in cloth bags and stored in a refrigerator for further use.

Experimental Design and Treatments

The experiment was laid out in randomized complete block design with split-split plot design and four replications, randomizing the sowing method in main plots, seed size in sub plots and seed priming treatments in sub-sub plots. The net plot size was 4×13 m.

Crop Husbandry

Seedbed was prepared according to treatments' requirement. For conventional tillage, the field was cultivated two times with disc harrow followed by two times cultivation along with planking. After land preparation, wheat seeds were broadcast in the field. For zero tillage, after the rice harvesting with combine harvester, wheat seeds were directly sown in field with the help of happy seeder and turbo seeder. Crop was sown on 26th and 27th of November, 2016 and 2017, respectively using 125 kg ha⁻¹ seed rate. Fertilizer was applied at 125-100-60 kg ha⁻¹ nitrogen (N), phosphorus (P) and potassium (K). The whole P and K along with half N were applied at sowing, while remaining N was applied with first irrigation at tillering stage. Totally three irrigations were applied from sowing till maturity. First irrigation was applied 35 days after sowing (DAS), while 2nd and 3rd irrigations were applied in the 4th week of February, and 3rd week of March, respectively. Bromoxynil at 1250 mL ha⁻¹ was applied five days after first irrigation with a knapsack sprayer for the control of broad leaf weeds. Fenoxaprop P-ethyl at 1000 mL ha⁻¹ was applied three days after 2nd irrigation for the control of narrow leaf weeds. The crop was harvested manually at its maturity. The first year trial was harvested on 29th of April, 2017; while the second year trial was harvested on 27th April, 2018.

Observation and Measurements

Ten plants were selected at random from each sub-plot and individual plant height from soil surface to the tip of the ear head was measured with the help of a meter rod and then averaged. Total tillers and productive tillers were counted from unit area (1 m × 1 m) from each plot at harvesting time. Ten spikes were randomly selected and the spike length was recorded with the help of scale and then averaged. These spikes were threshed manually to separate the grains. The threshed grains were counted to record the number of grains per spike. Samples of 1000-grain were taken from each sub plot and then weighed on an electric balance. Total dry biomass (biological yield) of sun-dried samples was recorded for each treatment by using a weighing balance. The crop was threshed manually. Grain yield for each treatment was recorded by an electric balance and was converted into kg/ha and expressed in tons per hectare (t/ha). The values of grain yield were adjusted to 12% moisture level. Harvest index (HI) was calculated as the ratio of grain yield to the biological yield, and was expressed in percentage.

Statistical Analysis

The collected data were analyzed statistically by employing the Fisher's analysis of variance technique (Steel *et al.*, 1997) using Statistix 8.1 (Analytical software, Statistix; Tallahassee, FL, USA, 1985-2003) and honest significance difference (HSD) Tukey's test at 5% probability level was

used to compare the differences among treatments means. None of the interactions were significant hence data only on main effects has been presented.

Results

Germination and Phenological Traits

Among seed size, small seeded cultivar (Lasani-2008) showed the highest germination counts in 2016-17 (Table 2). While during second year, maximum germination count was recorded in small seeded (Lasani-2008) and medium seed sized cultivar (FSD-2008). Among seed priming treatments hydroprimed wheat seed showed highest germination count that was statistically similar with osmopriming during first year; while during 2017-2018, hydropriming was best in this concern. Among wheat sowing methods, wheat planted with turbo seeder had the highest germination count during both years under study (Table 2).

Seed priming significantly affected the different reproductive phases from booting to physiological maturity during both years. Wheat sowing methods significantly affected the days to booting during first years. Among seed priming non-primed seed (control) took less time to reach booting, heading, anthesis and maturity during both years (Table 2).

Morphological, Yield and Yield Related Traits

The highest total tillers were recorded with osmopriming that was statistically similar with hydropriming during first year of study. While during second year osmopriming was best in this concern. Osmoprimed seeds produced maximum productive tillers that were statistically similar with no priming during second year of study. The maximum spike length was recorded with hydropriming that was statistically similar with no priming during both growing seasons. During first year, the maximum grains per spike were recorded with hydropriming (Table 3).

Among seed priming, the maximum biological yield was recorded with hydropriming that was statistically equal with osmopriming during 2016-2017 and 2017-2018. Osmopriming produced maximum biological yield that was statistically equal with hydropriming (Table 4). During first year; while during second year; the maximum harvest index was recorded with osmopriming that was statistically similar with hydropriming. For second year, the maximum harvest index was recorded in non-primed seed (control) that was statistically similar with hydropriming seeds (Table 4).

During first year, wheat planted with turbo seeder produced the highest number of total tillers. During second year, highest total tillers were recorded in wheat planted with turbo seeder that was statistically similar with wheat planted with happy seeder. The highest productive tillers were recorded in wheat planted with turbo seeder that was statistically similar with wheat planted with happy seeder during second year of study (Table 3).

Table 2: Seed germination and phenology of wheat cultivars as affected by seed priming and sowing methods

Treatments	Germination count (%)		Days to booting		Days to heading		Days to anthesis		Days to maturity	
	2016-17	2017-18	2016-17	2017-18	2016-17	2017-18	2016-17	2017-18	2016-17	2017-18
S ₁	222.5 B	221.4 B	97.1	95.5	101.8	100.1	112.5	111.8	138.2	135.5
S ₂	224.3 B	226.5 A	97.1	95.5	101.8	100.1	112.5	111.8	138.2	135.5
S ₃	227.4 A	225.1 A	97.1	95.5	101.8	100.1	112.5	111.8	138.2	135.5
HSD (<i>P</i> <0.05)	1.51	1.66	NS	NS	NS	NS	NS	NS	NS	NS
P ₁	222.08 B	219.1 C	95.3 C	93.7 B	101.0 C	96.4 C	111.6 B	110.8 B	137.5 B	134.7 B
P ₂	228.22 A	228.1 A	97.5 B	95.8 A	102.2 B	100.7 B	112.6 A	111.9 A	138.4 A	135.6 A
P ₃	226.72 A	225.7 B	98.0 A	96.3 A	102.8 A	101.1 A	112.9 A	112.1 A	138.5 A	135.8 A
HSD (<i>P</i> <0.05)	1.52	1.72	0.38	0.57	0.39	0.42	0.33	0.43	0.41	0.34
M ₁	225.2 B	223.9 B	96.8 B	95.2	101.8	97.5	112.3	111.6	138.2	135.4
M ₂	224.3 B	223.1 B	97.3 A	95.6	102.1	100.4	112.4	111.6	138.1	135.3
M ₃	227.4 A	225.9 A	96.7 B	95.1	101.9	100.3	112.4	111.6	138.2	135.5
HSD (<i>P</i> <0.05)	2.72	2.08	0.46	NS	NS	NS	NS	NS	NS	NS

Means not sharing a letter in common differ at 5% probability level by Tukey's HSD test. S₁= Bold seed (AS-2002), S₂= Medium seed (FSD-2008), S₃= Small seed (Lasani-2008); P₁= Control, P₂= Hydropriming, P₃= Osmopriming with CaCl₂ (1.2%), M₁= Conventional method (plough tillage), M₂= Happy seeder (zero tillage), M₃= Turbo seeder (zero tillage)

Table 3: Morphological and yield related traits of wheat cultivars as affected by seed priming and sowing methods

Treatments	Plant height (cm)		Total tillers (m ²)		Productive tillers (m ²)		Spike length (cm)		Spikelets per spike		Grains per spike	
	2016-17	2017-18	2016-17	2017-18	2016-17	2017-18	2016-17	2017-18	2016-17	2017-18	2016-17	2017-18
S ₁	87.0	87.2	314.0 B	303.7 B	314.6	300.6 A	9.04 AB	9.11 A	16	15	42.1 B	42.4
S ₂	89.1	88.6	319.5 AB	307.1 B	320.1	303.8 B	9.14 A	9.13 A	16	16	45.1 A	45.2
S ₃	85.1	86.0	331.3 A	319.6 A	328.6	316.3 A	8.82 B	8.85 B	16	16	45.0 A	46.8
HSD (<i>P</i> <0.05)	ns	Ns	15.5	12.2	Ns	12.45	0.23	0.18	ns	Ns	1.72	1.45
P ₁	86.7	87.0	321.6	310.6	319.3	307.6	9.16	9.14	16	15	44.1	44.5
P ₂	87.4	87.4	321.9	309.0	322.2	305.6	8.88	8.94	16	16	44.1	44.3
P ₃	87.0	87.4	321.3	310.7	321.6	307.5	8.96	9.00	16	16	44.1	45.6
HSD (<i>P</i> <0.05)	ns	Ns	Ns	ns	Ns	Ns	ns	ns	ns	Ns	ns	ns
M ₁	87.2	87.4	296.1 C	283.8 B	293.9 B	280.6 B	8.98	9.06	16	16	44.6	44.3
M ₂	86.9	87.3	323.6 B	314.1 A	326.6 A	310.8 A	9.09	9.05	16	15	43.5	45.6
M ₃	86.9	87.1	345.1 A	332.5 A	342.7 A	329.2 A	8.93	8.98	16	16	43.3	43.7
HSD (<i>P</i> <0.05)	NS	NS	19.8	21.8	19.87	22.13	NS	NS	NS	NS	NS	NS

Means not sharing a letter in common differ at 5% probability level by Tukey's HSD test. S₁= Bold seed (AS-2002), S₂= Medium seed (FSD-2008), S₃= Small seed (Lasani-2008); P₁= Control, P₂= Hydropriming, P₃= Osmopriming with CaCl₂ (1.2%), M₁= Conventional method (plough tillage), M₂= Happy seeder (zero tillage), M₃= Turbo seeder (zero tillage)

During first year the maximum 1000-grain weight was recorded in bold seeded cultivar (AS-2002). While during second year, bold seeded cultivar (AS-2002) had highest 1000-grain weight that was statistically similar with medium seed sized cultivar (FSD-2008). For first year, medium seeded cultivar (FSD-2008) and small seeded cultivar (Lasani-2008) produced maximum grain yield (Table 4). While during second year, bold seeded cultivar (AS-2002) gave highest grain yield that was statistically similar with medium seeded cultivar (Lasani-2008) (Table 4). During first year, small seeded cultivar (Lasani-2008) had highest biological yield that was statistically similar with medium seeded cultivar (FSD-2008), while during second year small seeded cultivar (Lasani-2008) was best in this concern. Similarly, bold seeded cultivar (AS-2002) had maximum harvest index that was statistically similar with medium seeded cultivar (FSD-2008) during both years of study (Table 4).

Among seed priming, biological yield was maximum with hydropriming that was statistically similar with osmopriming during first year. While during second year, osmopriming produced maximum biological yield that was statistically similar with hydropriming (Table 4).

The maximum harvest index was recorded with osmopriming that was statistically similar with hydropriming during first year. During second year, harvest index was maximum with no priming that was statistically similar with hydropriming (Table 4).

Among WSM, during first year, the maximum 1000-grain weight was recorded in wheat planted with happy seeder that was statistically similar with plough tillage wheat. Wheat planted with happy seeder produced maximum grain yield that was statistically similar with wheat planted with turbo seeder during second year. The maximum biological yield was recorded in wheat planted with happy seeder during first year and similar trend was observed during second year. During first year, the maximum harvest index was recorded in wheat planted with plough tillage that was statistically similar with wheat planted with turbo seeder (Table 4).

Economic Analysis

The maximum net income was obtained when medium size seed (FSD-2008) variety was hydroprimed and planted with happy seeder followed by small size seed (Lasani-2008)

Table 4: Yield and related traits of wheat cultivars as affected by seed priming and sowing methods

Treatments	1000-grain weight (g)		Grain yield (t ha ⁻¹)		Biological yield (t ha ⁻¹)		Harvest index (%)	
	2016-17	2017-18	2016-17	2017-18	2016-17	2017-18	2016-17	2017-18
S ₁	43.2 A	39.3 A	3.5 B	3.2 A	8.1 B	8.5 B	43.6 A	38.3 A
S ₂	40.5 B	38.6 A	3.6 A	3.1 AB	8.9 A	7.9 C	42.2 AB	39.1 A
S ₃	36.5 C	34.7 B	3.6 A	3.1 B	9.6 A	9.1 A	49.6 B	33.5 B
HSD (<i>P</i> <0.05)	0.73	1.43	0.15	0.12	0.48	0.34	2.79	2.09
P ₁	39.9	37.4	3.6	3.07	9.5 B	8.0 B	40.1 B	39.02 A
P ₂	39.9	37.5	3.6	3.12	8.8 A	8.7 A	41.6 AB	36.6 AB
P ₃	40.6	37.5	3.6	3.11	8.3 A	8.9 A	43.8 A	35.4 B
HSD (<i>P</i> <0.05)	NS	NS	NS	NS	0.33	0.38	2.22	2.56
M ₁	40.0 AB	36.7	3.6	2.8 B	8.3 B	8.1 B	44.1 A	36.1
M ₂	40.2 A	38.1	3.6	3.3 A	9.4 A	9.2 A	38.9 B	36.6
M ₃	39.6 B	37.7	3.6	3.1 A	8.6 B	8.3 B	42.5 A	38.3
HSD (<i>P</i> <0.05)	0.47	NS	NS	0.22	0.42	0.21	2.93	NS

Means not sharing a letter in common differ at 5% probability level by Tukey's HSD test. S₁= Bold seed (AS-2002), S₂= Medium seed (FSD-2008), S₃= Small seed (Lasani-2008); P₁= Control, P₂= Hydropriming, P₃= Osmopriming with CaCl₂ (1.2%), M₁= Conventional method (plough tillage), M₂= Happy seeder (zero tillage), M₃= Turbo seeder (zero tillage)

Table 5: Economic analysis of potential role of seed size and enhancements in improving productivity of wheat sown by different sowing methods

Treatments	Rabi 2016-2017	Rabi 2017-2018	pooled yield	Income (\$ ha ⁻¹)	Cost of Prod. (\$ ha ⁻¹)	BCR	Net Income (\$ ha ⁻¹)		
Conventional Method	As-2002	Control (non-primed)	3.55	3.04	3.30	1027.5	612.9	1.68	414.6
		Hydro-priming	3.82	2.89	3.35	1045.4	612.9	1.71	432.6
		Osmo-priming	3.46	3.04	3.25	1013.5	612.9	1.65	400.6
	FSD-2008	Control (non-primed)	3.65	2.79	3.22	1003	612.9	1.64	390.1
		Hydro-priming	3.27	2.84	3.05	951.51	612.9	1.55	338.6
		Osmo-priming	3.38	2.76	3.07	956.96	612.9	1.56	344.1
	Lasani-2008	Control (non-primed)	3.48	2.70	3.09	962.81	612.9	1.57	349.9
		Hydro-priming	3.72	2.92	3.32	1034.9	612.9	1.69	422.0
		Osmo-priming	4.10	2.68	3.39	1056.7	612.9	1.72	443.9
Happy seeder	As-2002	Control (non-primed)	3.52	3.25	3.38	1054	505.0	2.09	549.0
		Hydro-priming	3.26	3.44	3.35	1045.4	505.0	2.07	540.4
		Osmo-priming	3.37	3.49	3.43	1068.4	505.0	2.12	563.4
	FSD-2008	Control (non-primed)	3.80	3.38	3.59	1118.3	505.0	2.21	613.3
		Hydro-priming	3.96	3.46	3.71	1155.8	505.0	2.29	650.7
		Osmo-priming	3.76	3.19	3.47	1083.3	505.0	2.15	578.2
	Lasani-2008	Control (non-primed)	3.67	3.10	3.39	1056	505.0	2.09	551.0
		Hydro-priming	3.81	3.13	3.47	1081.3	505.0	2.14	576.3
		Osmo-priming	3.77	3.50	3.63	1132.4	505.0	2.24	627.4
Turbo seeder	As-2002	Control (non-primed)	3.58	3.40	3.49	1088.3	500.3	2.18	588.1
		Hydro-priming	3.64	3.01	3.33	1037.7	500.3	2.07	537.4
		Osmo-priming	3.39	3.31	3.35	1044.7	500.3	2.09	544.4
	FSD-2008	Control (non-primed)	3.80	3.04	3.42	1065.7	500.3	2.13	565.4
		Hydro-priming	4.02	3.19	3.60	1123.8	500.3	2.25	623.5
		Osmo-priming	3.53	3.00	3.26	1017.4	500.3	2.03	517.1
	Lasani-2008	Control (non-primed)	3.59	3.09	3.34	1041.2	500.3	2.08	540.9
		Hydro-priming	3.51	3.23	3.37	1050.9	500.3	2.10	550.6
		Osmo-priming	3.54	3.01	3.27	1020.9	500.3	2.04	520.6

variety using happy seeder combined with osmopriming, medium size seed (FSD-2008) by using turbo seeder with hydropriming method and also medium size seed (FSD-2008) by using happy seeder in conventional method (Table 5).

Discussion

Germination count and yield related traits of wheat cultivars markedly influenced by different seed priming techniques and sowing methods (Tables 2-4) *i.e.*, vigorous seeds germinate quickly and produce plants with more height and more number of tillers per unit area and results in higher grain

yield and production (López-Bellido *et al.*, 2007). All three wheat varieties varied in their performance due to their difference in seed size. The numbers of tillers were maximum in FSD-2008 (medium seed size) due to its genetic character, while number of grains per spike was higher in small seeded cultivar (Lasani-2008) due to less thousand grains weight and size. Growth of seedling depends on the amount of energy stored in the seed. Test weight of bold seed was higher due to increased endosperm size and more accumulation of photo assimilates. The 1000-grain weight of all wheat cultivars (with different seed size) declined during 2018 due to abrupt increase in temperature at grain filling stage which affects the overall grain weight and yield.

Seed size has strong influence on the seedling establishment; seed with more size has more food reserve and often exhibiting rapid emergence resulting in higher grain yield and harvest efficiency (Willenborg *et al.*, 2005). Moreover, germination rate and seedling vigor improve with increase in the seed size (Meyer and Carlson, 2001). Because, in bold seed stored food reserve are the only source of food (Kaya and Day, 2008) unless the seedling start to manufacture food through photosynthesis. Thus, the more food reserve and larger embryo size in bold seeded cultivar enhance the seed germination and improve root growth in different tillage systems. Guillen-Portal *et al.* (2006) reported, large sized seed tend to develop better roots system that enable the seedling to get moisture efficiently from deeper soil layers. As seed size increases, the seed protein and starch contents increases that may lead to improved and faster plant growth (Anuradha *et al.*, 2009).

Wheat sowing methods (conventional and conservation) had no significant impact on phenological stages of wheat. However, the conservation tillage in wheat (turbo seeder and happy seeder) enhanced the seedling emergence, improved the yield related traits and grain yield of wheat (Table 2, 3 and 4). It was reported by Hao *et al.* (2001) and Hemmat and Eskandari (2004) that, under different tillage systems wheat grain yield increased by 27 to 31% over conventional tillage system. The benefits of conservation tillage were previously discussed by Farooq *et al.* (2011). Zero tillage in wheat facilitate timely sowing of wheat, improve soil structure and biological properties (soil microbial biomass carbon), increase input use efficiency that help to maximize the crop productivity (Gupta and Seth, 2007; Mohanty *et al.*, 2007; Liu *et al.*, 2010).

Primed seed meets the optimum growth cycle by diminishing the emergence period (Khan *et al.*, 2014, 2015). Seed priming is a partial hydration practice that metabolizes the seed reserve, make it ready to use by germinating embryo. It speeds up the emergence process and ultimately reduces the days to booting, heading and anthesis of wheat and ultimately results in early physiological maturity (Hussain *et al.*, 2016). However, in this study, wheat crop raised from unprimed seeds showed poor germination with less grain yield as compared to osmoprimed seed sown under conservation tillage systems. Poor crop stand generated by unprimed seeds under conservation tillage may be due to poor seed-soil contact that cause delayed germination (Nawaz *et al.*, 2016). Seed priming with CaCl_2 was effective in improving the stand establishment under zero tillage systems that may be due to decrease in time of germination due to improved germination metabolism. Seed priming with CaCl_2 help achieving the uniform and early crop stand through improving the germination metabolism, quick expansion of embryo and plant tillering capacity resulting in better yield (Pandita *et al.*, 2007; Sarlach *et al.*, 2013). Furthermore, calcium (Ca^{2+}) has vital role in carbohydrate metabolism during stand establishment and initial plant developmental phases; osmopriming with CaCl_2 may

improve the Ca^{2+} contents that are crucial for seed metabolism (Farooq *et al.*, 2010).

Earlier emergence, vigorous seedlings and early flowering of osmoprimed wheat seeds than unprimed seeds might be the reason of better wheat performance under zero tillage system (Haider *et al.*, 2016). Although there is poor soil-seed contact under zero tillage systems, however, primed seed exhibited early germination because of sufficient availability of internal seed moisture that stimulate the radicle protrusion (Nawaz *et al.*, 2016). Maximum net income, benefit to cost ratio and net returns were obtained by planting osmoprimed medium size seeds (Faisalabad-2008) with happy seeder.

Conclusion

Wheat planted under conservation tillage such as using turbo seeder and happy seeder improved the seedling emergence, yield related traits resulting in higher grain yield compared to wheat sown by conventional system. Use of bold grain seeds cultivar produced the higher yield. Moreover, seed priming (hydropriming and osmopriming) further enhanced the performance of wheat sown under conservation tillage.

References

- Afzal, I., S. Rauf, S.M.A. Basra and G. Murtaza, 2008. Halopriming improves vigour, metabolism of reserves and ionic content in wheat seedling under salt stress. *Plant Soil Environ.*, 549: 382–388
- Ahmad, S.M. and C. Meisner, 1996. *Wheat Research and Development in Bangladesh, Bangladesh-Australia Wheat Improvement Project*, p: 201. CIMMYT-Bangladesh, Dhaka, Bangladesh
- Ahmad, T., Ehsanullah and K. Jabran, 2013. Economic efficiency of different cropping systems under Faisalabad conditions. *J. Agric. Res.*, 51: 419–430
- Akbari, G.H.A., M.G. Pirbalouti, M.N. Farahani and M. Shahverdi, 2004. Effect of harvesting time on soybean seed germination and vigor. *J. Agric.*, 6: 9–18
- Alvarez, R. and H.S. Steinbach, 2009. A review of the effects of tillage systems on some soil physical properties, water content, nitrate availability and crops yield in the Argentine, Pampas. *Soil Till. Res.*, 104: 1–15
- Anuradha, R., P. Balamurugan, P. Srimathi and S. Sumathi, 2009. Influence of seed size on seed quality of chick pea (*Cicer arietinum* L.). *Legume Res. Int. J.*, 32: 133–135
- Baker, C.J., K.E. Saxton and W.R. Ritchie, 1996. *No-Tillage Seeding, Science and Practice*, p: 158. CAB International, Wallingford, Oxon, UK
- Erenstein, O. and V. Laxmi, 2008. Zero tillage impacts in India's rice-wheat systems. *Soil Till. Res.*, 100: 1–14
- Erenstein, O., K. Sayre, P. Wall, J. Dixon and J. Hellin, 2008. Adapting no-tillage agriculture to the conditions of smallholder maize and wheat farmers in the tropics and sub-tropics. In: *No-till Farming Systems, Special Publication 3. World Association of Soil and Water Conservation (WASWC)*, pp: 253–278. Goddard, T., M. Zoenbisch, Y. Gan, W. Ellis, A. Watson, S. Sombatpanit, (eds.). Bangkok
- Farooq, M., K.C. Flower, K. Jabran, A. Wahid and K.H.M. Siddique, 2011. Crop yield and weed management in rainfed conservation agriculture. *Soil Till. Res.*, 117: 172–183
- Farooq, M., S.M.A. Basra, A. Wahid and N. Ahmad, 2010. Changes in nutrient-homeostasis and reserves metabolism during rice seed priming: consequences for germination and seedling growth. *Agric. Sci. Chin.*, 9: 101–108

- Farooq, S., M. Hussain, K. Jabran, W. Hassan, M.S. Rizwan and T.A. Yasir, 2017. Osmopriming with CaCl₂ improves wheat (*Triticum aestivum* L.) production under water-limited environments. *Environ. Sci. Pollut. Res.*, 24: 13638–13649
- Fugelsnes, E. and E. Lie, 2011. *Heavy Agricultural Machinery can Damage the Soil*. Available online at <http://www.sciencedaily.com/releases/2011/05/110505083737.htm> (Accessed on March 03, 2015)
- Gorge, M.H.A. and R.T. Ray, 2005. Germination characterization of guayule seed by morphology, mass and, X-ray analysis. *Ind. Crop Prod.*, 23: 59–63
- Guillen-Portal, F.R., R.N. Stouggaard, Q. Xue and K.M. Eskridge, 2006. Compensatory mechanism associated with the effect of spring wheat seed size on wild oat competition. *Crop Sci.*, 46: 935–945
- Gupta, R. and A. Seth, 2007. A review of resource conserving technologies for sustainable management of the rice-wheat cropping systems of the Indo-Gangetic plains (IGP). *Crop Prot.*, 26: 436–447
- Haider, M.U., M. Hussain, M.B. Khan, M. Ijaz, A. Sattar, M. Akram and W. Hassan, 2016. Influence of seed priming and seed size on wheat performance under different tillage systems. *Int. J. Agric. Biol.*, 18: 858–864
- Hao, X., C. Chang, R.L. Conner and P. Bergen, 2001. Effect of minimum tillage and crop sequence on crop yield and quality under irrigation in Southern Alberta clay loam soil. *Soil Till. Res.*, 59: 45–55
- Harris, D., D. Rashid, G. Miraj, M. Arif and H. Shah, 2007. 'On farm' seed priming with zinc sulphate solution- A cost-effective way to increase the maize yields of resource- poor farmers. *Field Crop Res.*, 102: 119–127
- Hemmat, A. and I. Eskandari, 2004. Conservation tillage practices for winter wheat-fallow farming in the temperate continental climate of northwestern Iran. *Soil Till. Res.*, 89: 123–133
- Hussain, M., M. Waqas-Ul-Haq, S. Farooq, K. Jabran and M. Farooq, 2016. The impact of seed priming and row spacing on the productivity of different cultivars of irrigated wheat under early season drought. *Exp. Agric.*, 52: 477–490
- Hussain, M., M.B. Khan, M. Shahzad, A. Ullah, A. Sher and A. Sattar, 2017. Influence of priming on emergence, weed infestation, growth and yield of wheat sown under different tillage practices. *Int. J. Agric. Biol.*, 19: 367–373
- Jabran, K., Ehsanullah, M. Farooq, M. Hussain, N. Haider and B.S. Chauhan, 2015. Water saving, water productivity and yield outputs of fine-grain rice cultivars under conventional and water-saving rice production systems. *Exp. Agric.*, 51: 567–581
- Karina, M.V.C., A.P.D. Silva, C.A. Tormena, T.P. Leaˆo, A.R. Dexter and I.H. Kansson, 2009. Long-term effects of no-tillage on dynamic soil physical properties in a Rhodic Ferrasol in Paranaˆ, Brazil. *Soil Till. Res.*, 103: 158–164
- Kaya, M.D. and S. Day, 2008. Relationship between seed size and NaCl on germination, seed vigor and early seedling growth of sunflower (*Helianthus annuus* L.). *Afr. J. Agric. Res.*, 3: 787–791
- Khan, M.B., M. Hussain, A. Raza, S. Farooq and K. Jabran, 2015. Seed priming with CaCl₂ and ridge planting improves drought resistance of maize. *Turk. J. Agric. For.*, 39: 193–203
- Khan, M.B., M.A. Gurchani, S. Freed, K. Jabran and M. Hussain, 2014. Osmopriming improves the emergence, growth, nutrient uptake and soluble sugar contents of wheat seedlings. *Soil Environ.*, 33: 142–148
- Lampurlaneas, J., P. Angaˆs and C. Cantero-Martˆoˆnez, 2001. Root growth, soil water content and yield of barley under different tillage systems on two soils in semiarid conditions. *Field Crop Res.*, 69: 27–40
- Liu, E.K., B.Q. Zhao, X.R. Mei, H.B. So, J. Li and X.Y. Li, 2010. Effect of no tillage management on soil biochemical characteristics in northern China. *Chin. J. Agric. Sci.*, 148: 217–223
- Lˆopez-Bellido, R.J., L. Lˆopez-Bellido, J. Benitez-Vega and F.J. Lˆopez-Bellido, 2007. Tillage system, preceding crop, and nitrogen fertilizer in wheat crop: I. Soil water content. *Agron. J.*, 99: 59–65
- Martinez, E., J. Fuentes, P. Silva, S. Valle and E. Acevedo, 2008. Soil physical properties and wheat root growth as affected by no-tillage and conventional tillage systems in a Mediterranean environment of Chile. *Soil Till. Res.*, 99: 232–244
- Meyer, S.E. and S.L. Carlson, 2001. Achene mass variation in *Ericamerian auseosus* (Asteraceae) in relation to dispersal ability and seedling fitness. *Funct. Ecol.*, 15: 274–281
- Mohanty, M., D.K. Painuli, A.K. Misra and P.K. Ghosh, 2007. Soil quality effects of tillage and residue under rice-wheat cropping on a Vertisol in India. *Soil Till. Res.*, 92: 243–250
- Nawaz, A., M. Farooq, R. Ahmad, S.M.A. Basra and R. Lal, 2016. Seed priming improves stand establishment and productivity of no till wheat grown after direct seeded aerobic and transplanted flooded rice. *Eur. J. Agron.*, 76: 130–137
- Pandita, V.K., A. Anand and S. Nagarajan, 2007. Enhancement of seed germination in hot pepper following pre-sowing treatments. *Seed Sci. Technol.*, 35: 282–290
- Rashidi, M. and F. Keshavarzpour, 2008. Effect of different tillage methods on soil physical properties and crop yield of melon (*Cucumis melo*). *J. Agric. Environ. Sci.*, 3: 31–36
- Rehman, H., S.M.A. Basra, M. Farooq, N. Ahmed and I. Afzal, 2011. Seed priming with CaCl₂ improves the stand establishment, yield and quality attributes in direct seeded rice (*Oryza sativa* L.). *Int. J. Agric. Biol.*, 113: 786–790
- Sarlach, R.S., A. Sharma and N.S. Bains, 2013. Seed priming in wheat: effect on seed germination, yield parameters and grain yield. *J. Progressive Res.*, 8: 109–112
- Shahwani, A.R., S.U. Baloch, S.K. Baloch, B. Mengal, W. Bashir, H.N. Baloch and A. Ahmed, 2014. Influence of seed size on germination ability and grain yield of wheat (*Triticum aestivum* L.) varieties. *J. Nat. Sci. Res.*, 4: 23
- Shahzad, M., M. Hussain, M. Farooq, S. Farooq, K. Jabran and A. Nawaz, 2017. Economic assessment of conventional and conservation tillage practices in different wheat-based cropping systems of Punjab, Pakistan. *Environ. Sci. Pollut. Res.*, 24: 24634–24643
- Steel, R.G.D., J.H. Torrie and D.A. Dickny, 1997. *Principles and Procedures of Statistics, A Biometrical Approach*, pp. 352–358, 3rd edition. McGraw Hill Book Co., Inc., New York, USA
- Willenborg, C.J., J.C. Wildeman, A.K. Miller, B.G. Rossnaged and S.J. Shirliffe, 2005. Oat germination characteristics differ among genotypes, seed sizes and osmotic potentials. *Crop Sci.*, 45: 2023–2029

(Received 03 November 2018; Accepted 26 November 2018)