



## Review Article

# Research Advances on Cuticular Waxes Biosynthesis in Crops: A Review

Md Shaheenuzzamn, Tianxiang Liu, Shandang Shi, Hongqi Wu and Zhonghua Wang\*

College of Agronomy, Northwest A&F University, Yangling, Shaanxi 712100, China

\*For correspondence: zhonghuawang@nwafu.edu.cn

## Abstract

Cuticular waxes are the hydrocarbon consisting of very long chain primary alcohols, aldehydes, fatty acids, alkane and esters. They are hydrophobic layer which protect aerial plant organs and help plant species for adaptation in different environments. Wax deposition and chemical composition vary considerably among crop species. Cuticular waxes play a significant role against major abiotic stresses in plants such as drought, high salinity and cold. So, it draws close attention to molecular processes of cuticular wax biosynthesis under stress factors. Here, we briefly summarized the existing knowledge on the cuticular waxes properties, diversity, morphological changes in leaf surface wax crystals and cuticular wax amount and compositions of cuticular wax. We also provide information about wax biosynthesis genes in crops. Recently, due to progress of plant genome sequence, numerous genes involved in biosynthesis of cuticular waxes have been characterized both for model plant (*Arabidopsis*) and crops such as rapeseed (*Brassica napus*), *Camelina spp*, potato (*Solanum tuberosum*), eggplant (*S. macrocarpon*), tomato (*S. lycopersicum*), barley (*Hordeum vulgare*), rice (*Oryza sativa*), maize (*Zea mays*), wheat (*Triticum aestivum*), broccoli (*B. oleracea*), sesame (*Sesamum indicum*), tobacco (*Nicotiana tabacum*), cucumber (*Cucumis sativus*), cabbage (*B. oleracea*) etc. Basic compositions of cuticular wax are alcohols, branched alkanes, alkenes, aldehydes, fatty acids, esters, ketones, triterpenoids and sterols in crops. However, they vary from one crop species to the other. Cuticular wax biosynthesis is organ-specific and depends upon developmental stages of crops, and induced by environmental stimuli. The genetic factors also control wax biosynthesis and composition. However, cuticular wax also acts as a photoprotector layer during photosynthesis and protect from UV light radiation. It is also linked to gas exchange and plant development. In this review, we have summarized the cuticular wax deposition and composition, and also identified genes involved in cuticular wax biosynthesis in several crop species. This knowledge may be helpful in potential applications for selection of crop for agricultural sustainability. © 2019 Friends Science Publishers

**Keywords:** Crops; Cuticular wax; Environmental stimuli; Wax biosynthesis; Wax diversity

## Introduction

In growth and development stages, plants have to face enormous environmental stresses like drought, salinity, cold, heat, UV light, high radiation, insect or fungal and pathogens. The cuticular waxes are a surface layer of the plant which provide defense against pests and pathogens (Wink, 1988; Holmes and Keiller, 2002; Bargel *et al.*, 2004; Yeats and Rose, 2013). The cuticle is composed of two distinct layers which chemically separate compounds including a lipophilic cutin polymer matrix and waxes (Holloway, 1982; Jeffree, 1996; Kunst *et al.*, 2005). Cuticular waxes are most important elements which prevent uncontrolled evaporation of water at the leaves surfaces (Jetter and Riederer, 2000; Knoche *et al.*, 2000). The cuticular wax is composed of very long-chain fatty acid compounds (VLCFAs; C20 to C34). These VLCFAs compounds consist of Branched alkanes, primary alcohols, alkenes, aldehydes, secondary alcohols,  $\beta$ - and OH- $\beta$ -diketones, esters and often triterpenoids and flavonoids

(Jetter *et al.*, 2006; Samuels *et al.*, 2008). The genetic and environmental factors influence on the deposition and composition of cuticular waxes (Bianchi, 1995; Post-Beittenmiller, 1996). Cuticular wax biosynthetic pathways have been studied extensively in *Arabidopsis* (Hannoufa *et al.*, 1993; McNevin *et al.*, 1993; Jenks *et al.*, 1995; Suh *et al.*, 2005; Kunst and Samuels, 2009; Nawrath *et al.*, 2013). Several cuticular waxes genes from *Arabidopsis* were identified such as *FATB*, *LCASILACS2*, *LCAS4*, *ACC1*, *KCS1*, *KCS2/DAISY*, *KCS6/CER6/CUT1*, *KCS9*, *KCS20*, *KCR1*, *HCD/PAS2*, *ECR/ECR10*, *CER2*, *CER2-LIKE1*, *CER2-LIKE2*, *CER1*, *RST1*, *CYTB5-B*, *CYTB5-C*, *CYTB-D*, *CYTB5-E*, *MAH1*, *FAR3/CER4* and *WSD1* (Lee and Suh, 2015). Number of cuticular wax genes has already been identified in crop species (Table 1). Transcription, mRNA and post-translational modification are controlled by these genes in waxy and waxless plants (Von Wettstein-Knowles, 1995; Pu *et al.*, 2013; Lee *et al.*, 2015). However, genetic mechanism related to deposition of cuticular waxes in crop species is still elusive and subject to further investigations.

Cuticular wax composition also depends on leaf color, insect-plant interaction and plant development. The quantity of plant cuticular wax largely depends upon environment conditions. Researchers have great interest to comprehend the detail genetic behavior of wax biosynthesis genes in crop species. In this review, we summarized the amount and contents of cuticular waxes in the different crop species and focused on recent progress about the molecular and biological function of genes engaged in biosynthesis of cuticular wax. We also briefly discussed the properties of cuticular waxes and diversities of cuticular waxes in crop species.

### Properties of Cuticular Wax

**Plant response and adaptation to abiotic and biotic stresses:** Plant transpiration depends on two factors. Basically plant transpiration take place through stomata and a non-stomatal component is also there. Bernard and Joubès (2013) reported that there is significance correlation between the lipid cuticle layer and transpiration which was first proof about the role of cuticle for non-stomatal water loss (Stiles, 1994). Stomata remains close during water stress or night time, and provide space for significant cuticle transpiration. Several plant studies such as on tobacco and sesame also showed that wax biosynthesis was increased during water stress, and played an important role in preventing the cuticular desiccation (Cameron *et al.*, 2006; Kim *et al.*, 2007). In Arabidopsis, water and osmotic stresses increased wax deposition that in turn were associated with a resistance to water stress, and cuticle formation could be a part of mechanism to acquire tolerance to water stress (Kosma *et al.*, 2009). Amount of wax and resistance to water flow depends on the cuticle biosynthesis enzymes (Aharoni *et al.*, 2004; Bourdenx *et al.*, 2011; Seo *et al.*, 2011). The regulation of cuticular permeability mainly depends on the wax deposition mechanisms during water stress. Several studies have reported that increase in cuticular permeability reduces wax load and vice versa (Chen *et al.*, 2003; Zhang *et al.*, 2005; Kosma *et al.*, 2009; Lü *et al.*, 2009, 2012; Bourdenx *et al.*, 2011; Seo *et al.*, 2011). Higher cuticular permeability depends on increased amount of cutin and waxes depositions.

**Cuticular waxes need for plant development:** Beside their major contribution to stress tolerance, cuticular waxes are active players in the growth and developmental processes in plant. Kurdyukov *et al.* (2006) reported that organ fusion phenotypes were frequently associated with severe imperfections in either cuticular wax or cutin biosynthesis, as noticed in, *bodyguard*, *Wddlehead*, *cer3/wax2* and numerous other mutants. Other studies on Arabidopsis also indicated that *lacs1lacs2* double-mutant plants demonstrated pleiotropic phenotypes such as organ fusion, unusual flower development and decreased seed set (Weng *et al.*, 2010). The wax-deficient *cer1* mutant of Arabidopsis had a conditional male-sterile phenotype that

reduced pollen viability contributed to the low seed yield (Aarts *et al.*, 1995). These studies indicated that a deficiency in cuticular waxes biosynthesis or cutin synthesis has a effects on cuticular barrier, water movement, defend against drought stress and protect organ fusion. Arabidopsis mutant's analyses help us to understand of changes in cuticular waxes amount and composition during different developmental stages and add conception to the action of cuticular waxes in plant physiology.

### Diversity of Cuticular Wax

Despite our relatively advanced understanding of wax compound structure and biosynthesis in Arabidopsis, crucial questions remain unanswered about how chemical composition determines the physical properties of the cuticular wax mixture. Before addressing these questions, a thorough understanding of the major dimensions of cuticular wax diversity is needed, in particular, the diversity in the chemical structures and diversity in the waxes covering on different biological organs. It will move us closer to a fundamental understanding of the relationships between structure and function in the plant cuticular wax diversity.

**Structural diversities of wax compound:** Structural diversity may be evident from the aliphatic tail (e.g., number of unsaturations, aliphatic branches, TCN, etc) or functional groups (e.g., number of functional groups, positions on the aliphatic tail, oxidation state, etc) in the wax molecules. Some of those compounds were found in large amount in different plants and crops, indicating that they play a vital role in the properties of cuticular waxes and alter cuticular wax mixture. Moreover, their biosynthesis mechanisms of converting branched wax precursors into branched wax compounds are still not clear. It is needed to characterize branched wax compounds biosynthesis pathway to remove uncertainty of model species. Fatty acids, primary alcohols, alkenes, wax esters, aldehydes and branched alkanes are major cuticular wax components. However, wax profiles in different plant organs also revealed that some plants accumulate secondary functional groups with major cuticular wax compounds (Gunthardt-Goerg, 1986; Wen *et al.*, 2006). Ketones, ketoalcohols, and ketoaldehydes found on the surfaces of the fern *Osmunda re-alias* (Jetter and Riederer, 2000), while  $\beta$ -diketones are present on the surface of wheat and barley (Tulloch and Weenink, 1969; Jackson, 1971; Han-Avivi *et al.*, 2016; Schneider *et al.*, 2016; Huang *et al.*, 2017). It was proved that the true diversity of cuticular waxes presents in different plants. Thus, our knowledge about branched wax compound and biosynthesis is still nascent stage.

**Biological variability in wax coverage and composition:** Aerial plant organs are covered with waxes throughout their developmental stages. The quantity and composition wax in plants depend upon the stresses in an age-dependent manner. Indeed, previous studies had reported that wax

**Table 1:** Genes known to be involved in cuticular wax biosynthesis in crop species

Mechanism	Species	Organs	Protein family name	Abbreviation	Reference
Cuticular wax biosynthesis	Maize	Seedling leaf	Arabidopsis <i>KCS6</i> homolog	<i>GL4</i>	Avato <i>et al.</i> , 1987; Liu <i>et al.</i> , 2009
		Seedling leaf	Arabidopsis <i>KCR</i> homolog	<i>GL8a</i>	Xu <i>et al.</i> , 1997; Dietrich <i>et al.</i> , 2005
		Seedling leaf	Arabidopsis <i>KCR</i> homolog	<i>GL8b</i>	Dietrich <i>et al.</i> , 2005
		Seedling leaf	Arabidopsis <i>CER3</i> homolog	<i>GL1</i>	Hansen <i>et al.</i> , 1997; Sturaro <i>et al.</i> , 2005
		Seedling leaf	Arabidopsis <i>CER2</i> homolog	<i>GL2</i>	Lemieux, 1996; Velasco <i>et al.</i> , 2002
		Leaf	KCS	<i>WSL1</i>	Yu <i>et al.</i> , 2008
		Shoot	KCS	<i>ONH1</i>	Ito <i>et al.</i> , 2011
		Anther	Arabidopsis <i>CER1</i> homolog	<i>WDA1</i>	Jung <i>et al.</i> , 2006
		Leaf	Arabidopsis <i>CER1</i> homolog	<i>OsGL1-6</i>	Zhou <i>et al.</i> , 2013
		Leaf	Arabidopsis <i>CER3</i> , maize <i>GL1</i> homolog	<i>OsGL1-2</i>	Islam <i>et al.</i> , 2009
		Leaf	Arabidopsis <i>CER3</i> homolog	<i>OsGL1-1/WSL2</i>	Qin <i>et al.</i> , 2011; Mao <i>et al.</i> , 2012
		Leaf	AMP-binding domain contained	<i>DWA1</i>	Zhu and Xiong, 2013
		Leaf	Arabidopsis <i>CER3</i> homolog/ Maize <i>GL1</i> homolog	<i>OsGL1-3</i>	Zhou <i>et al.</i> , 2015
		Leaf	A homolog of the <i>MBOAT</i> transferase family	<i>OsWS1</i>	Xia <i>et al.</i> , 2015
		Leaf	<i>NAD<sup>+</sup>/NADP<sup>+</sup></i> -dependent sterol dehydrogenase	<i>OsHSD1</i>	Zhanget <i>et al.</i> , 2016
		Leaf	<i>KCR</i>	<i>WSL3</i>	Gan <i>et al.</i> , 2016
		Leaf	<i>KCS</i>	<i>WSL3</i>	Hong-bing <i>et al.</i> , 2017
		Leaf	Arabidopsis <i>CER6</i> homolog	<i>WSL4</i>	Wang <i>et al.</i> , 2017
		Leaf	<i>CER6-like KCS</i>	<i>LeCER6</i>	Vogg <i>et al.</i> , 2004;
		Tomato	Fruit	B-Amyrin synthesis	<i>SITTS1</i>
	Tomato	Fruit	Oxidosqualene cyclase	<i>SITTS2</i>	Wang <i>et al.</i> , 2011
	Tomato	Leaf	Unknown	<i>BnaA.GL</i>	Pu <i>et al.</i> , 2013
	<i>B. napus</i>	Leaf	lipid transfer proteins	<i>BraLTP1</i>	Liu <i>et al.</i> , 2014
	<i>Brassica rapa</i>	Leaf	Arabidopsis <i>CER2</i> homolog	<i>BrWax1</i>	Zhang <i>et al.</i> , 2013a
		Leaf	Arabidopsis <i>KCS2</i> homolog	<i>CsKCS2</i>	
		Leaf	Arabidopsis <i>KCS6</i> homolog	<i>CsKCS6</i>	
		Leaf	Arabidopsis <i>KCR 1</i> homolog	<i>CsKCR1-1</i>	Lee <i>et al.</i> , 2014
	<i>C. sativa</i>	Leaf	Arabidopsis <i>KCR 1</i> homolog	<i>CsKCR1-2</i>	
		Leaf	Arabidopsis <i>ECR</i> homolog	<i>CsECR</i>	
	Cucumber	Leaf	Arabidopsis <i>KCS2</i> homolog	<i>CsMAH1</i>	
		Leaf	Arabidopsis <i>WAX2</i> homolog	<i>CsWAX2</i>	Wang <i>et al.</i> , 2015a
		Leaf	Arabidopsis <i>CER1</i> homolog	<i>CsCER1</i>	Wang <i>et al.</i> , 2015b
		leaf	lipid transfer proteins	<i>BoLTP2</i>	
	Cabbage	Leaf	Arabidopsis <i>CER3</i> homolog	<i>BoCER3</i>	
		Leaf	Arabidopsis <i>KCS1</i> homolog	<i>BoKCS1</i>	
		Leaf	Arabidopsis <i>KCR1</i> homolog	<i>BoKCR1</i>	Laila <i>et al.</i> , 2017
		Leaf	Arabidopsis <i>LACS1</i> homolog	<i>BoLACS1</i>	
		Leaf	alkane hydroxylase <i>CYP96A15</i>	<i>BoMAH1</i>	
		Leaf	Arabidopsis <i>CER4</i> homolog	<i>BoFAR3</i>	
		Leaf	Arabidopsis <i>WSD1</i> -like family	<i>BoWSD1</i>	
Leaf		Arabidopsis <i>CER4-6</i> homolog	<i>W1W2</i>	Zhang <i>et al.</i> , 2013b	
Leaf		<i>CER1</i> and <i>CER3</i> homologs	<i>W3</i>	Zhang <i>et al.</i> , 2015	
Leaf		Arabidopsis <i>CER4</i> homolog	<i>TaFAR1</i>	Wang <i>et al.</i> , 2015b	
Wheat	Leaf	Arabidopsis <i>CER4</i> homolog	<i>TaFAR5</i>	Wang <i>et al.</i> , 2015c	
	Leaf	Arabidopsis <i>CER4</i> homolog	<i>TaFAR2, TaFAR3, TaFAR4,</i>	Wang <i>et al.</i> , 2016	
	Spike	miRNA (MIRNA)	<i>W1-COE and / or W2-COE</i>	Huang <i>et al.</i> , 2017	
	Leaf	Arabidopsis <i>CER4</i> homolog	<i>Ae.tFAR1, Ae.tFAR2,</i>	Wang <i>et al.</i> , 2017	
			<i>Ae.tFAR3, Ae.tFAR4, Ae.tFAR6</i>		
			<i>TaFAR6, TaFAR7, TaFAR8</i>	Chai <i>et al.</i> , 2018	
Barley	Spike	PKS (DMP), Hydrolase (DMH), CYP450 (DMC)	<i>Cer-cqu</i>	Hen-Avivi <i>et al.</i> , 2016; Schneider <i>et al.</i> , 2016	
	Leaf	Protein phosphatase 2C family protein	<i>Cer-b</i>	Zhou <i>et al.</i> , 2017	

amount and compositions vary between plant surfaces of different ages (Atkin and Hamilton, 1950; Gülz *et al.*, 1992; Viougeas *et al.*, 1995). Largely, the understanding of cuticular wax deposition on plant surfaces needs further studies (Suh *et al.*, 2005). Plant surface development depends on cuticular wax deposition at different environmental conditions. It is needed to investigate the

dimensions of biological variability and its relationship with the developmental biology of the plants. Thus, Cuticular waxes display structural diversity in their aliphatic tails and functional group (s) and biological variability depending on surfaces of different species, surfaces of different plant organs, and organ surfaces at different ages.


## Cuticular Wax Morphology

The wax morphology was examined on both the adaxial and abaxial sides of leaf by scanning electron microscopy (SEM) to achieve insight into epicuticular wax crystals (Fig. 1). Leaf blades were collected from D genome (*Aegilops tauschii*) at three plant development stages (seedling, heading and filling stages). Two forms of wax crystals: platelets and tubules (A–L Figs) were found in the wheat leaf. Barthlott (1998) classified the epicuticular waxes. Some platelets shape wax crystals were joined to their adjacent crystals making a dense network. The length of platelet shaped wax crystals was between 0.3 and 0.7  $\mu\text{m}$  and height between 0.3 and 0.5  $\mu\text{m}$ . The platelet shaped wax crystals had irregular margins, and were present at different angles with respect to each other (Fig. 1 A–L). So, the wax morphology changed during their development stages. It is concluded that as the plant ages, the cuticular wax morphology changes on the leaf surface (Wang *et al.*, 2015a).

## Cuticular Waxes in Crop Species

**Amounts and contents of cuticular waxes in crop:** It is very well known that cuticular waxes deposition varies across crop species and from organ to organ (Barthlott *et al.*, 1998; Kosma *et al.*, 2010; Buschhaus and Jetter, 2011; Bernard and Joubès, 2013). Cuticular wax contents were measured in different organs of Arabidopsis, rapeseed (*Brassica napus*), *Camelina* spp, potato (*S. tuberosum*), eggplant (*S. macrocarpon*), tomato (*S. lycopersicum*), barley (*Hordeum vulgare*), maize (*Zea mays*), rice (*Oryza sativa*), wheat (*Triticum aestivum*), broccoli (*B. oleracea*), sesame (*Sesamum indicum*), tobacco (*Nicotiana tabacum*), cucumber (*Cucumis sativus*) and cabbage (*B. oleracea*) (Table 2). Previous studies have shown that in Arabidopsis ecotype Columbia-0, the wax was 0.7–1.5, 13–24, 13  $\mu\text{g}/\text{cm}^2$  in leaves, stems and siliques, while it was 23–82 and 36–170  $\mu\text{g}/\text{g}$  in flowers and seed coat, respectively. Alkane contributed up to 50% of total wax loads and represented the most dominant wax compound in all organs of Arabidopsis. In stems, silique walls, flowers, and seed coats, secondary alcohols and ketones were present but their very low amounts was noticed on leaf (Jenks *et al.*, 1995; Bernard and Joubès, 2013; Lee and Shu, 2015).

In rapeseed, cuticular wax amount in leaves is considerably higher than Arabidopsis leaves, while other wax compounds were similar (Pu *et al.*, 2013). In rapeseed breeding line 6-3476, the amount of wax in leaves (687–2255  $\mu\text{g}/\text{g}$ ) was similar to flower of Arabidopsis (2382  $\mu\text{g}/\text{g}$ ) (Tassone *et al.*, 2016). In *Camelina sativum* var Celina, the amount of wax in leaves, stem, flower and seed coat was 6.24, 164, 264, 0.24  $\mu\text{g}/\text{cm}^2$ , respectively. Wax esters (74%) in leaf and triterpenoids/sterols (53%) in stems were observed, but the primary alcohols (65%) were detected in seed coats (Razeq *et al.*, 2014).



**Fig. 1:** Epicuticular wax crystals patterns on the adaxial and abaxial leaf surfaces of the wheat D genome (*Aegilops tauschii*) detected by SEM at three stages of plant development. A, D are the adaxial surface of leaves during seedling stage; B, E are the adaxial surface of leaves during heading stage; C, F are the adaxial surface of leaves during filling stage; G, J are the abaxial of leaves during seedling stage; H, K are the abaxial surface of leaves during seedling stage; I, L are the abaxial surface of leaves during Filling stage. The micrographs are at a resolution of 10 000 $\times$  and 30 000 $\times$ , and the bars indicate 1  $\mu\text{m}$  and 0.5  $\mu\text{m}$ , respectively

Furthermore, in leaf of *Camelina* MYB96 transgenic line, the amount of wax was 2.9  $\mu\text{g}/\text{cm}^2$  while in the leaf of Robinson variety it was 0.72  $\mu\text{g}/\text{cm}^2$  (Tomasi *et al.*, 2017). In another report, the wax compositions of cultivar *C. sativum* var Celina was similar to that reported by Razeq *et al.* (2014). Furthermore, the amount and composition in leaves of *C. sativa* 1.37  $\mu\text{g}/\text{cm}^2$ , *C. rumelica* 2.01  $\mu\text{g}/\text{cm}^2$ , *C. hispida* 0.85  $\mu\text{g}/\text{cm}^2$  and *C. microcarpa* 0.84  $\mu\text{g}/\text{cm}^2$  were reported but total wax loads were lower than younger leaves. In all *Camelina* species, the primary alcohols and alkanes were dominant components followed by wax esters, fatty acids and aldehydes. Interestingly, *C. sativa* var MYB96 synthesized higher levels of primary alcohols. It indicates that MYB96 might be an effective gene involved in the primary alcohols biosynthesis (Tomasi *et al.*, 2017). In potato, the amount of wax in the leaves was 5  $\mu\text{g}/\text{cm}^2$  in Perkoz, 6  $\mu\text{g}/\text{cm}^2$  in Aster and Maryna, and 7  $\mu\text{g}/\text{cm}^2$  in Ibis. Alkanes were dominant compound and primary alcohols were second major class in all potato varieties (Szafranek and Synak, 2006). In Gboma eggplant plant, the wax in cultivar UVPP was 2.7  $\mu\text{g}/\text{cm}^2$  and in Urafiki it was 2.3  $\mu\text{g}/\text{cm}^2$  were observed in leaves. Alkanes, primary alcohols, fatty acids, sterols, and triterpenols were also found. Alkanes (47–56%) were the dominant component in cuticular wax in Gboma Eggplant. Sterols content were observed much higher than triterpenes, consisting of 19 and 32% of the total waxes in Urafiki and UVPP cultivars, respectively (Halinski *et al.*, 2012).

**Table 2:** Cuticular wax amounts and contents in different organs of crop species

Species	Organs	Total Loads	Components (relatives %)													Reference	
			Fatty acids	Aldehydes	Alkanes	Primary alcohols	Sec. alcohols	Ketons	Wax esters	Iso-alkanes	Anteiso-alkanes	Alkenols	Alkenes	Triterpenoids	Ferulate & Esters/Phenol		
Arabidopsis	Leaf	0.7–1.5 <sup>m</sup>	1	14	73	8	0	0	4	0	0	0	0	0	0	0	Bernard and Joubès, 2013; Li-Beisson <i>et al.</i> , 2013
	Stem <sup>a</sup>	13–24 <sup>m</sup>	1	7	44	12	9	22	5	0	0	0	0	0	0	0	Bernard and Joubès, 2013; Li-Beisson <i>et al.</i> , 2013
	Silique <sup>a</sup>	13 <sup>m</sup>		1	4	50	14	9	22	0	0	0	0	0	0	0	Kim <i>et al.</i> , 2012
	Flower <sup>a</sup>	2382 <sup>n</sup>	0	2	62	5	15	14	2	0	0	0	0	0	0	0	Bernard and Joubès, 2013
	Seed coat <sup>a</sup>	36–170 <sup>n</sup>	0	9	51	13	18	9	0	0	0	0	0	0	0	0	Li <i>et al.</i> 2007; Bernard and Joubès, 2013
Rapeseed	Leaf	29.4 <sup>m</sup>	1	4	55	2	10	27	2	0	0	0	0	0	0	0	Pu <i>et al.</i> , 2013
	Leaf <sup>b</sup>	687–2255 <sup>n</sup>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	Tassone <i>et al.</i> , 2016
<i>C. sativa</i>	Leaf <sup>c</sup>	6.2 <sup>m</sup>	3	0	3	20	0	0	74	0	0	0	0	0	0	0	Razeq <i>et al.</i> , 2014
	Steam <sup>c</sup>	16 <sup>m</sup>	6	0	28	13	0	0	0	0	0	0	0	53	0	0	
	Flower <sup>c</sup>	264 <sup>n</sup>	14	0	64	1	0	0	0	0	0	0	0	21	0	0	
	Seed coat <sup>c</sup>	0.2 <sup>m</sup>	6	0	29	65	0	0	0	0	0	0	0	0	0	0	
<i>C. sativa</i>	Leaf <sup>d</sup>	2.9 <sup>m</sup>	16	2	35	42	0	0	3	0	0	0	0	0	0	0	Tomasi <i>et al.</i> , 2017
<i>C. sativa</i>	Leaf <sup>e</sup>	0.72 <sup>m</sup>	9	0	9	49	0	0	17	0	0	0	0	0	0	0	
<i>C. sativa</i>	Leaf <sup>e</sup>	0.83 <sup>m</sup>	7	0	17	50	0	0	25	0	0	0	0	0	0	0	
<i>C. sativum</i>	leaf <sup>e</sup>	1.37 <sup>m</sup>	12	1	22	43	0	0	14	0	0	0	0	0	0	0	
<i>C. rumelica</i>	Leaf	2.01 <sup>m</sup>	6	1	35	40	0	0	17	0	0	0	0	0	0	0	
<i>C. hispida</i>	Leaf	0.85 <sup>m</sup>	7	0	20	27	0	0	40	0	0	0	0	0	0	0	
<i>C. microcarpa</i>	Leaf	0.84 <sup>m</sup>	4	0	45	44	0	0	0	0	0	0	0	0	0	0	
Potato	Leaf <sup>f</sup>	6 <sup>m</sup>	9	0	61	12	1	1	3	0	0	0	0	2	0	0	Szafranek and Synak, 2006
	Leaf <sup>g</sup>	7 <sup>m</sup>	9	0	65	10	0	1	5	0	0	0	0	1	0	0	
	Leaf <sup>h</sup>	6 <sup>m</sup>	10	0	61	11	1	2	2	0	0	0	0	1	0	0	
	Leaf <sup>i</sup>	5 <sup>m</sup>	5	0	61	7	1	2	2	0	0	0	0	0	1	0	
Egg Plant	Leaf <sup>j</sup>	2.7 <sup>m</sup>	3	0	47	16	0	0	0	0	0	0	5	19	0	0	Halinski <i>et al.</i> , 2012
	Leaf <sup>k</sup>	2.3 <sup>m</sup>	5	0	56	18	0	0	0	0	0	0	5	32	0	0	Wang <i>et al.</i> , 2011
	Leaf <sup>l</sup>	6.7 <sup>m</sup>	0	0	74	3	0	0	0	18	0	0	0	5	0	0	
	Anther <sup>l</sup>	2155 <sup>n</sup>	21	0	33	0	0	0	0	33	6	0	0	7	0	0	Smirnova <i>et al.</i> , 2013
Tomato	Fruit <sup>m</sup>	8.4 <sup>m</sup>	7	4	37	6	0	0	0	5	1	4	16	21	0	0	Isaacson <i>et al.</i> , 2009
	Fruit <sup>n</sup>	10.2 <sup>m</sup>	3	2	49	6	0	0	0	2	0	5	9	24	0	0	Kosma <i>et al.</i> , 2010
	Leaf <sup>L1</sup>	2094 <sup>n</sup>	2	0	72	1	0	0	0	0	0	0	0	10	0	0	Halinski <i>et al.</i> , 2015
	Leaf <sup>L2</sup>	2118 <sup>n</sup>	1	0	75	1	0	0	0	0	0	0	0	7	0	0	
	Leaf <sup>L3</sup>	1686 <sup>n</sup>	1	0	73	1	0	0	0	0	0	0	0	12	0	0	
	Leaf <sup>L4</sup>	1131 <sup>n</sup>	4	0	73	2	0	0	0	0	0	0	0	7	0	0	
	Leaf <sup>L5</sup>	1019 <sup>n</sup>	3	0	77	1	0	0	0	0	0	0	0	7	0	0	
	Leaf <sup>Pen1</sup>	2351 <sup>n</sup>	3	0	38	1	0	0	0	0	0	0	0	2	0	0	
	Leaf <sup>Pen2</sup>	3933 <sup>n</sup>	4	0	53	2	0	0	0	0	0	0	0	7	0	0	
	Leaf <sup>Pen3</sup>	3738 <sup>n</sup>	11	0	35	3	0	0	0	0	0	0	0	19	0	0	
	Leaf <sup>Pen4</sup>	3971 <sup>n</sup>	1	0	71	1	0	0	0	0	0	0	0	1	0	0	
	Leaf <sup>Pen5</sup>	3116 <sup>n</sup>	45	0	34	0	0	0	0	0	0	0	0	11	0	0	
	Leaf <sup>LxPen</sup>	2657 <sup>n</sup>	1	0	62	1	0	0	0	0	0	0	0	4	0	0	
	Leaf <sup>Pim1</sup>	1342 <sup>n</sup>	1	0	71	2	0	0	0	0	0	0	0	5	0	0	
Leaf <sup>LxPim</sup>	1069 <sup>n</sup>	2	0	71	1	0	0	0	0	0	0	0	5	0	0		
Leaf <sup>P</sup>	14.7 <sup>m</sup>	9	4	1	75	0	0	0	11	0	0	0	0	0	0	Von Wettstein-Knowles, 1971; Avato <i>et al.</i> , 1982	
Barley	Leaf <sup>P</sup>	14.6 <sup>m</sup>	2	1	1	6	0	59	0	0	0	0	0	0	0	0	Yu <i>et al.</i> , 2008

Table 2: Continued

Table 2: Continued

Rice	Leaf <sup>4</sup>	7.57 <sup>m</sup>	18	37	2	32	0	0	11	0	0	0	0	0	0	Mao <i>et al.</i> , 2012
	Blade															
	Leaf	5.8 <sup>m</sup>	14	32	3	36	0	0	16	0	0	0	0	0	0	
	Sheath <sup>4</sup>															
	Leaf	8.4 <sup>m</sup>	34	31	7	24	0	0	4	0	0	0	0	0	0	
Maize <sup>1</sup>	Blade															Javelle <i>et al.</i> , 2010
	Leaf	4.5 <sup>m</sup>	43	24	9	23	0	0	0	0	0	0	0	0	0	
	Sheath <sup>f</sup>															
	Leaf	6.2 <sup>m</sup>	58	0	19	22	0	0	0	0	0	0	0	0	0	
	Blade <sup>5</sup>															
Wheat	Leaf	8.2 <sup>m</sup>	0	25	4	69	0	0	2	0	0	0	0	0	0	Wang <i>et al.</i> , 2015a
	Blade															
	Leaf	4.5 <sup>m</sup>	0	42	14	39	0	5	0	0	0	0	0	0	0	
	Sheath															
	Leaf <sup>1</sup>	3.4 <sup>m</sup>	6	3	14	71	0	3	2	0	0	0	0	0	0	
	Leaf <sup>2</sup>	3.6 <sup>m</sup>	4	3	10	79	0	1	2	0	0	0	0	0	0	
	Leaf <sup>3</sup>	3.9 <sup>m</sup>	5	4	15	68	0	6	3	0	0	0	0	0	0	
	Leaf <sup>4</sup>	3.6 <sup>m</sup>	5	3	11	68	0	12	2	0	0	0	0	0	0	
	Spike <sup>1</sup>	4.5 <sup>m</sup>	13	1	40	18	0	37	0	0	0	0	0	0	0	
	Spike <sup>2</sup>	5.8 <sup>m</sup>	11	1	31	30	0	27	0	0	0	0	0	0	0	
Broccoli	Spike <sup>3</sup>	7.7 <sup>m</sup>	10	1	19	7	0	63	0	0	0	0	0	0	0	Zhang <i>et al.</i> , 2015
	Spike <sup>4</sup>	6.6 <sup>m</sup>	10	1	31	11	0	47	0	0	0	0	0	0	0	
	Leaf <sup>5</sup>		1	0	34	1	0	63	1	0	0	0	0	0	0	
	Leaf <sup>6</sup>	16 <sup>n</sup>	1	3	9	0	0	14	9	0	0	55	0	0	0	
	Peduncle <sup>7</sup>	4.9 <sup>m</sup>	1	1	7	0	0	81	2	0	0	2	0	0	0	
	Seedling leaf <sup>7</sup>	5.4 <sup>m</sup>	1	5	2	84	0	0	1	0	0	0	0	0	0	
	Flag leaf <sup>7</sup>	8.4 <sup>m</sup>	2	9	6	77	0	1	1	0	0	0	0	0	0	
	Leaf sheaths <sup>7</sup>	3.5 <sup>m</sup>	2	4	15	57	0	15	3	0	0	0	0	0	0	
	Peduncles <sup>7</sup>	2.9 <sup>m</sup>	3	13	37	23	0	36	2	0	0	0	0	0	0	
	Glumes <sup>7</sup>	1.0 <sup>m</sup>	2	7	10	30	0	10	2	0	0	0	0	0	0	
Sesame	Anthers <sup>7</sup>	0.3 <sup>m</sup>	54	1	27	3	0	27	1	0	0	0	0	0	0	Lee <i>et al.</i> , 2015
	Leaf <sup>18</sup>	1929 n	13	20	39	3	5	19	0	0	0	0	0	0	0	
	Leaf <sup>19</sup>	3733 n	15	13	40	3	7	22	0	0	0	0	0	0	0	
Tobacco	Leaf <sup>10</sup>	7.69 m	0	11	68	0	0	0	0	0	0	0	0	0	0	Kim <i>et al.</i> , 2007
	Leaf <sup>11</sup>	13.9m	7.8	0	74	7	0	0	0	0	0	0	0	0	0	
Cucumber	Friut <sup>12</sup>	1.3m	21	22	46	0	3	0	0	2	0	0	2	0	3	Wang <i>et al.</i> , 2015a
	Stem <sup>12</sup>	1.6m	6	8	82	0	10	0	0	1	0	0	1	0	3	
	Leaf <sup>12</sup>	1.8m	8	15	62	0	8	0	0	5	0	0	0	0	2	
Cabbage	Leaf <sup>13</sup>	0	0	6	34	6	14	31	0	0	0	0	0	0	0	Laila <i>et al.</i> , 2017

Note: number 0 indicates that trace or undetectable amounts were audited. aecotype col-0, bbreeding line 6-3476, ccultivar *cemelina sativum* var celina, dc. *sativum* var myb 96, ec. *sativum* var robinson, fpotato cultivar aster, gpotato cultivar ibis, hpotato cultivar maryna, ipotato cultivar perkoz, jegg plant cultivar uvpp, kgboma egg plant cultivar urafiki, lcultivar micro tom, mcultivar tomato m82, ncultivar tomato ailsa craig, 11tomato cultivar vf-36, 12tomato cultivar wild (ecuador), 13tomato cultivar wild (ecuador), 14tomato cultivar nagcarlang, 15tomato cultivar wild (usa), pen1tomato cultivar wild (peru), pen2tomato cultivar wild (peru), pen3tomato cultivar wild (peru), pen4 tomatocultivar wild (peru), pen5 tomato cultivar wild (peru), lxpentomato cultivar hybrid, pim1tomato cultivar wild (ecuador), lxpintomato cultivar hybrid, obarley cultivar bonus, pbarley cultivar bowman, qrice cultivar nipponbare, rice cultivar japonica, smaize inbred line a188, 1wheat cultivar a14, 2wheat cultivar jing 2001, 3wheat cultivar fanmai 5,4wheat cultivar shanken 99, 5wheat cultivar bob white, 6wheat cultivar bethlehem, 7wild grass *ae. tauschii*. 8broccoli cultivar mc 117,9broccoli cultivar mc91, 10sesame cultivar various, 11 tobacco cultivar graha, 12cucumber cultivar wild type, 13cabbage cultivar *b. oleracea*, m unit =  $\mu\text{g}/\text{cm}^2$ , n unit =  $\mu\text{g}/\text{g}$

In tomato cultivar Micro-Tom, amount and composition of cuticular waxes in leaf and anther were 6.7  $\mu\text{g}/\text{cm}^2$  and 2155  $\mu\text{g}/\text{g}$ , respectively, and the most abundant components were alkanes (74%) in leaves (Wang *et al.*, 2011). Tomato cultivar M82 and cultivar Ailsa Craig in fruits contained 8.4  $\mu\text{g}/\text{cm}^2$  and 10.2  $\mu\text{g}/\text{cm}^2$  wax compounds, respectively while alkanes and alkenes were the dominant wax component (Isaacson *et al.*, 2009; Kosma *et al.*, 2010). However, the most dominant waxes components of these varieties were alkanes ranged from 34 to 71% (Halinski *et al.*, 2015). In tomato leaves, fruits and anthers

too branched alkanes, alkenes, and cyclic compounds were detected (Isaacson *et al.*, 2009; Wang *et al.*, 2011; Smirnova *et al.*, 2013; Halinski *et al.*, 2015). This finding was in the agreement with the other researchers that cuticular waxes in tomato leaf were consisted by hydrocarbons (Zygadlo *et al.*, 1994; Smith *et al.*, 1996; Vogg *et al.*, 2004). Primary alcohols, aldehydes and fatty acid were found to be principal components of cuticular wax, where alkanes occupied less than 15% of the total wax loads in leaves of barley, rice and maize (Von Wettstein-Knowles, 1971; Avato *et al.*, 1982; Javelle *et al.*, 2010; Mao *et al.*, 2012)



whereas ketones were the most abundant component in leaves of barley cultivar Bowman (Yu *et al.*, 2008).

In wheat, primary alcohols were the major components in leaves cuticular wax followed by alkanes, esters, aldehydes and fatty acids. Ketones deposition variations were observed in wheat leaves. In wheat leaves, ketones were found in trace amount the cuticular wax. However, in spike, a traceable amount of ketones was identified (Wang *et al.*, 2015a, 2017). In addition, there were large differences in the amounts of ketones among different wheat varieties (Wang *et al.*, 2015a; Zhang *et al.*, 2015). Wheat cultivar Bethlehem revealed wax coverage of 16  $\mu\text{g}/\text{cm}^2$  in leaf and 49  $\mu\text{g}/\text{cm}^2$  in peduncles (Zhang *et al.*, 2015). Furthermore, in wheat cultivar Bethlehem, alkanols were 55% of total wax loads in leaf and  $\beta$ -diketone and hydroxy- $\beta$ -diketones collectively comprised 81% of the total wax loads in peduncle. This happened due to discrepancy in the regulation of the acyl-reduction and  $\beta$ -diketone biosynthetic pathways in the two examined organs (Racovita *et al.*, 2016).

In broccoli cultivars MC117 and MC91, fatty acids (13 and 15%, respectively), aldehydes (20 and 13%, respectively), alkanes (38 and 39%, respectively) and ketones (19 and 21%, respectively) were found in the total wax loads in leaf (Lee *et al.*, 2015). In sesame, major components of waxes in leaves were alkanes (68% of total wax) and aldehydes (11% of total wax) (Kim *et al.*, 2007). In tobacco, the primary component of cuticular wax was alkanes, which constituted 75% of the total wax load fully expanded leaves while fatty acids and alcohols occupied smaller proportion of total wax loads (Cameron *et al.*, 2006). In cucumber, alkenes, primary alcohols, branched alkanes, phenols, esters, and aldehydes, phenols were major compounds in the cuticular waxes (Wang *et al.*, 2015a). In cabbage leaf, alkane contributed 34% followed by ketones 31% of total wax loads (Laila *et al.*, 2017).

In spite of a vast variation in cuticular wax loads and contents depending upon crop species and organs, the prime reasons that responsible for these variations are not identified. Nevertheless, the present information may be used as a genetic source for determination of new genes associated with cuticular wax biosynthesis.

**Biosynthesis of cuticular wax genes in crop species:** In crop species, the genes responsible for biosynthesis of cuticular wax are not well characterized such as in *Arabidopsis*, though, recently notable advances have been made in this respect (Table 1). Identification and functional expression of the glossy (*gl*) mutants provided an opportunity to understand about wax biosynthesis in maize. Mutants of *GL4* and *GL8* are homolog of *AtKCS6* and *AtKCR*, respectively. They displayed a spectacular decrease in the amount of alkanes, alcohols and aldehydes in the leaves of wild type seedling (Avato *et al.*, 1987; Dietrich *et al.*, 2005; Liu *et al.*, 2009). In another study on maize, it was found that *gl1* and *gl2* mutants which were homolog to *AtCER3* and *AtCER2*, respectively, declined wax deposition

in seedling leaves, especially the aldehydes levels significantly decreased or could not be identified. Contrarily, the *gl1gl2* double mutant enhanced the levels of wax esters (Bianchi *et al.*, 1979; Lemieux, 1996; Hansen *et al.*, 1997; Velasco *et al.*, 2002; Sturaro *et al.*, 2005). It was also found that *GL13*, an ABC transporter, was associated with cuticular wax deposition (Li *et al.*, 2013).

In rice, *crystal-spires leaf1* (*wsl1*) mutant, is a *KCS* gene which has a lesion of *WSL1* gene, catalyzed the creation of C20–C24 VLCFA precursors of leaf waxes. It reduced growth, fertility, leaf fusion and increased drought sensitivity due to wax-deficiency. This is indicated that *WSL1* might be involved in the deposition of other lipids associated with growth and development of the plant (Yu *et al.*, 2008). *ONIONI* (*ONII*) is another homolog to rice *KCS* protein. It is accountable for synthesis of C20 and C22 saturated VLCFAs, which are important for development of shoot (Ito *et al.*, 2011). From functional expression of wax-deficient *anther1* (*wda1*) mutant, it was observed that a *WDA1* protein, which is homolog to *AtCER1*, was responsible for synthesis of VLC alkenes and alkanes in pollen and anthers (Jung *et al.*, 2006). Functional characterization of *OsGLI-6*, which is also homolog to *AtCER1*, reported that it is needed for wax biosynthesis on leaf blades. It was also found that reduced expression of the *OsGLI-6* gene was linked with notable decreases in total wax loads and enhanced drought sensitivity (Zhou *et al.*, 2013). Characterization of rice mutants, *gl1-2* and *gl1-1/wsl2*, revealed that these mutants reduced overall cuticular wax loads and increased sensitivity to drought stress (Islam *et al.*, 2009; Qin *et al.*, 2011; Mao *et al.*, 2012). Over expression of *Drought-Induced Wax Accumulation 1* (*DWA1*) held an AMP-binding domain, which increased VLCFA synthesis and enhanced drought resistance (Zhu and Xiong, 2013). *OsGLI-3* is homologous to maize *GL1* and *Arabidopsis* *WAX2/YRE/CER3/FLP*, which significantly increased biosynthesis of cuticular wax and enhanced tolerance to water stress (Zhou *et al.*, 2015). *OsWS1* belongs to the membrane-bound O-acyl transferase gene family, and is associated with wax biosynthesis in rice (Xia *et al.*, 2015). *OsHSD1* belongs to the short-chain dehydrogenase reductase family, which enhanced VLCFAs biosynthesis and soluble fatty acids in the leaves of the *oshsd1* mutant (Zhang *et al.*, 2016). *WSL3* encodes *KCR* in rice, which contributes to VLCFA biosynthesis and wax depositions in leaf. On the other hand in rice mutant, *wax crystal-sparse leaf 3* (*wsl3*) gene reduced epicuticular wax crystals and wax composition on the leaf surface (Gan *et al.*, 2016). *ORF4* is homologous to the *KCS6* family of *KCS*, which is similar to *WSL3* gene of rice and regulates cuticular wax formation (Hong-bing *et al.*, 2017). *WSL4* encodes a *KCS*, a homolog of *AtCER6*, which increased the cuticular wax load in rice leaves (Wang *et al.*, 2017).

In tomato fruit cuticle, the *lecer6* mutant belongs to *CER6-like KCS* (*LeCER6*), decreased accumulation of alkanes and aldehydes but on the other hand, amounts of

triterpenoid increased in the total wax loads (Vogg *et al.*, 2004). In tomato fruits, the over expression of *SITTS1* and *SITTS2* enhanced biosynthesis of terpenoid (Wang *et al.*, 2011). However, alkane is the dominant wax component in the tomato wax but genes/proteins have not yet been identified for the biosynthesis of alkanes. In *B. napus*, glossy mutant *BnaA.GL* was characterized, which reduced wax biosynthesis and increased sensitivity to drought stress (Pu *et al.*, 2013). *BraLTP1* belongs to non-specific lipid transfer proteins (*nsLTPs*), which decreased wax deposition in leaves (Liu *et al.*, 2014). In *B. rapa*, the *BrWax1* gene is found on linkage group A01, which is involved in cuticular wax biosynthesis in leaves (Zhang *et al.*, 2013a). In *Camelina sativa*, few wax biosynthesis genes have been detected recently but their characterization is not complete yet (Lee *et al.*, 2014). In Cucumber, *CsWAX2* is homolog of *AtWAX2*, which performs fundamental functions in wax biosynthesis (Wang *et al.*, 2015a) and *CsCER1*, a homolog of *AtCER1*, played an important role in wax VLC alkanes biosynthesis (Wang *et al.*, 2015b). In Cabbage, *BoLTP2* gene is a non-specific *lipid-transfer protein1*, involved in the transformation of ketones to lipid which reduced wax deposition. *BoCER3* gene is homologous to *AtCER3* protein, which is related to wax deposition through converting aldehydes to alkanes. *BoKCS1* and *BoKCR1* are homologs to *AtKCS* and *AtKCR1* protein, respectively, which decreased acyl-CoAs synthesis and eventually influence total wax loads. Likewise, expressions of *BoLACS1* influenced wax depositions. Besides, the *BoMAH1* gene engaged in synthesis of secondary alcohols and ketones (Laila *et al.*, 2017).

In wheat, *WIW2* is a homolog of Arabidopsis *CER4-6* proteins. It produced hydroxyl- $\beta$ -diketones, which enhanced drought tolerance through reducing cuticle permeability (Zhang *et al.*, 2013b). *W3*, a homolog of Arabidopsis *CER1* and *CER3*, is essential for  $\beta$ -diketone biosynthesis but suppresses its hydroxylation (Zhang *et al.*, 2015). *TaFAR1*, a homolog of Arabidopsis *CER4*, is an active acyl-CoA reductase. It produced primary alcohols, and as a result augmented total wax loads on wheat leaf blades (Wang *et al.*, 2015b). *TaFAR5*, a homolog to Arabidopsis *CER4*, is an alcohol-forming fatty acyl-coenzyme A reductase (*FAR*), which contributes significantly to produce primary alcohols in wheat leaf blade (Wang *et al.*, 2015c). *TaFAR2*, *TaFAR3*, and *TaFAR4* genes are a homolog of Arabidopsis *CER4* protein, which produced primary alcohols in cuticular wax (Wang *et al.*, 2016). From characterization of the *Iw* genes, it was found that *Iw* genes regulatory mechanism control *WCOE* expression and  $\beta$ -diketone formation (Huang *et al.*, 2017). In Dgenome (*Ae. Tauschii*), *Ae.tFAR1*, *Ae.tFAR2*, *Ae.tFAR3*, *Ae.tFAR4*, and *Ae.tFAR6* are homolog to *AtCER4*, which principally accountable for deposition of primary alcohols (Wang *et al.*, 2017). In barley, the *Cer-cqu* gene cluster is involved in  $\beta$ -diketons biosynthesis which consists of several proteins families including type-III polyketide synthases, hydrolases, and cytochrome P450s

(Hen-Avivi *et al.*, 2016; Schneider *et al.*, 2016). The barley *eceriferum-b.2* (*cer-b.2*) mutant produces  $\beta$ -diketons, which makes glossy leaf sheaths and deficient in the cuticular wax component, 14, 16-hentriacontanedione (Zhou *et al.*, 2017).

## Conclusion

This review has revealed that wax compositional differences exist among different crop species even organ to organ. High levels of structural diversities associates with cuticular wax deposition in crop species is largely influenced by different genes. Several genes have been already identified, which are related to cuticular wax biosynthesis. Cuticular wax components are produced by two different complex pathways due to the influence of biotic and abiotic stress, which allow adaptive mechanism at the time of crop-environment interactions. Moreover, specific single wax components yet remain unknown during development and growth stages of different crop species. Very few studies have focused on primary alcohols and ketons in wheat and barley, but most of the factors are still unidentified. Additionally, genome sequencing technologies have been progressed enormously, allowed identifying new race of cuticular waxes biosynthesis genes in crop species. Knowledge on the wax biosynthesis mechanisms in different plant species will be helpful to breed new crops cultivars better tolerant to environmental stresses.

## References

- Aarts, M.G., C.J. Keijzer, W.J. Stiekema and A. Pereira, 1995. Molecular characterization of the *CER1* gene of arabidopsis involved in epicuticular wax biosynthesis and pollen fertility. *Plant Cell*, 7: 2115–2127
- Aharoni, A., S. Dixit, R. Jetter, E. Thoenes, G.V. Arkel and A. Pereira, 2004. The *SHINE* clade of AP2 domain transcription factors activates wax biosynthesis, alters cuticle properties, and confers drought tolerance when overexpressed in *Arabidopsis*. *Plant Cell*, 16: 2463–2480
- Atkin, D.S.J. and R.J. Hamilton, 1950. The changes with age in the epicuticular wax of sorghum bicolor. *J. Nat. Prod.*, 45: 697–703
- Avato, P., G. Bianchi, A. Nayak, F. Salamini and E. Gentinetta, 1987. Epicuticular waxes of maize as affected by the interaction of mutant *gl8* with *gl3*, *gl4* and *gl15*. *Lipids*, 22: 11–16
- Avato, P., J.D. Mikkelsen and P. von Wettstein-Knowles, 1982. Synthesis of epicuticular primary alcohols and intracellular fatty acids by tissue slices from *cer-j59* barley leaves. *Carlsberg Res. Commun.*, 47: 377–390
- Bargel, H., W. Barthlott, K. Koch, L. Schreiber and C. Neinhuis, 2004. 10- Plant cuticles: multifunctional interfaces between plant and environment. In: *Evolution Plant Physiology*, pp: 171–194. A.R. Hemsley and I. Poole (eds.). Academic Press, London
- Barthlott, W., C. Neinhuis, D. Cutler, F. Ditsch, I. Muesel, I. Theisen and H. Wilhelm, 1998. Classification and terminology of plant epicuticular waxes. *Bot. J. Linn. Soc.*, 126: 237–260
- Bernard, A. and J. Joubès, 2013. Arabidopsis cuticular waxes: advances in synthesis, export and regulation. *Prog. Lipid Res.*, 52: 110–129
- Bianchi, G., 1995. Plant waxes. In: *Waxes: Chemistry, Molecular Biology and Functions*. pp: 175–222. Hamilton, R.J. (ed.). The Oily Press, Dundee, Scotland, UK
- Bianchi, G., P. Avato and F. Salamini, 1979. The absence of fatty acid aldehydes in *gl1 gl2* waxes. *Maize Gen Coop Newsl.*, 53: 103



- Bourdenx, B., A. Bernard, F. Domergue, S. Pascal, A. Léger, D. Roby, M. Pervent, D. Vile, R.P. Haslam, J.A. Napier, R. Lessire and J. Joubès, 2011. Overexpression of Arabidopsis *ECERIFERUM1* promotes wax very-long-chain alkane biosynthesis and influences plant response to biotic and abiotic stresses. *Plant Physiol.*, 156: 29–45
- Buschhaus, C. and R. Jetter, 2011. Composition differences between epicuticular and intracuticular wax substructures: how do plants seal their epidermal surfaces. *J. Exp. Bot.*, 62: 841–853
- Cameron, M.A., Teece and L.B. Smart, 2006. Increased accumulation of cuticular wax and expression of lipid transfer protein in response to periodic drying events in leaves of tree tobacco. *Plant Physiol.*, 140: 176–183
- Chai, G., C. Li, F. Xu, Y. Li, X. Shi, Y. Wang and Z. Wang, 2018. Three endoplasmic reticulum-associated fatty acyl-coenzyme A reductases were involved in the production of primaryalcohols in hexaploid wheat (*Triticum aestivum* L.). *BMC Plant Biol.*, 18: 41
- Chen, X., S.M. Goodwin, V.L. Boroff, X. Liu and M.A. Jenks, 2003. Cloning and characterization of the *WAX2* gene of Arabidopsis involved in cuticle membrane and wax production. *Plant Cell*, 15: 1170–1185
- Dietrich, C.R., M.A.D.N. Perera, D.M. Yandean-Nelson, R.B. Meeley and B.J. Nikolau, 2005. Characterization of two *GL8* paralogs reveals that the 3-ketoacyl reductase component of fatty acid elongase is essential for maize (*Zea mays* L.) development. *Plant J.*, 42: 844–861
- Gan, L., X. Wang, Z. Cheng, L. Liu, J. Wang, Z. Zhang, Y. Ren, C. Lei, Z. Zhao, S. Zhu, Q. Lin, F. Wu, X. Gao, J. Wang, X. Zhang and J. Wan, 2016. *Wax crystal-sparse leaf 3* encoding a  $\beta$ -ketoacyl-CoA reductase is involved in cuticular wax biosynthesis in rice. *Plant Cell Rep.*, 35: 1687–1698
- Gülz, P.G., E. Müller, K. Schmitz, F.J. Mamer and S. Güth. 1992. Chemical composition and surface structures of epicuticular leaf waxes of *Ginkgo biloba*, *Magnolia grandiflora* and *Liriodendron tulipifera*. *Z Naturforsch C.*, 47: 516–526
- Gunthardt-Goerg, M.S., 1986. Epicuticular wax of needles of *Pinus cembra*, *Pinus sylvestris* and *Picea abies*. *Eur. J. For. Pathol.*, 16: 400–408
- Halinski, L.P., M. Kalkowska, M. Kalkowski, J. Piorunowska, A. Topolewska, P. Stepnowski, 2015. Cuticular wax variation in the tomato (*Solanum lycopersicum* L.) related wild species and their interspecific hybrids. *Biochem. Syst. Ecol.*, 60: 215–224
- Halinski, L.P., M. Paszkiewicz, M.G. Biowski and P. Stepnowski, 2012. The chemical composition of cuticular waxes from leaves of the gboma eggplant (*Solanum macrocarpon* L.). *J. Food Comp. Anal.*, 25: 74–78
- Hannoufa, A., J. McNeven and B. Lemieux, 1993. Epicuticular waxes of eceriferum mutants of *Arabidopsis thaliana*. *Phytochemistry*, 33: 851–855
- Hansen, J.D., J. Pyee, Y. Xia, T.J. Wen, D.S. Robertson, P.E. Kolattukudy, B.J. Nikolau and P.S. Schnable, 1997. The *glossy1* locus of maize and an epidermis-specific cDNA from *Klenia odora* define a class of receptor-like proteins required for the normal accumulation of cuticular waxes. *Plant Physiol.*, 113: 1091–1100
- Hen-Avivi, S., O. Savin, R.C. Racovita, W.S. Lee, N.M. Adamski, S. Malitsky, E. Almekias-Siegl, M. Levy, S. Vautrin, H. Bergès, G. Friedlander, E. Kartvelishvily, G. Ben-Zvi, N. Alkan, C. Uauy, K. Kanyuka, R. Jetter, A. Distelfeld and A. Aharoni, 2016. A metabolic gene cluster in the wheat *W1* and the barley *Cer-cqu* loci determines  $\beta$ -diketone biosynthesis and glaucousness. *Plant Cell*, 28: 1440–1460
- Holloway, P.J., 1982. Structure and histochemistry of plant cuticular membranes: An overview. In: *The Plant Cuticle*, pp: 1–32. D.F. Cutler, K.L. Alvin and C.E. Price (eds.). Academic Press, London
- Holmes, M.G. and D.R. Keiller, 2002. Effects of pubescence and waxes on the reflectance of leaves in the ultraviolet and photosynthetic wavebands: a comparison of a range of species. *Plant Cell Environ.*, 25: 85–93
- Hong-bing, G., Z. Sheng-yuan, X. Xiang, Z. Ya-fang, C. Zong-xiang, Z. Shi-min, L. Chuang, L. Tian-zi, J. De-dao, Y. Bo, Q. Hua-fei, P. Xue-biao and S. Sheng-lan, 2017. Fine mapping of a novel wax *crystal-sparse leaf3* gene in rice. *J. Integr. Agric.*, 16: 497–502
- Huang, D., J.A. Feurtadao, M.A. Smitha, L.K. Flatmana, C. Koha and A.J. Cutler, 2017. Long noncoding miRNA gene represses wheat  $\beta$ -diketone waxes. *Proc. Natl. Acad. Sci. USA*, 114: E3149–E3158
- Isaacson, T., D.K. Kosma, A.J. Matas, G.J. Buda, Y. He, B. Yu, A. Pravitari, J.D. Batteas, R.E. Stark, M.A. Jenks and J.K.C. Rose, 2009. Cutin deficiency in the tomato fruit cuticle consistently affects resistance to microbial infection and biomechanical properties, but not transpirational water loss. *Plant J.*, 60: 363–377
- Islam, M.A., H. Du, J. Ning, H. Ye and L. Xiong, 2009. Characterization of *Glossy1* homologous genes in rice involved in leaf wax accumulation and drought resistance. *Plant Mol. Biol.*, 70: 443–456
- Ito, Y., F. Kimura, K. Hirakata, K. Tsuda and T. Takasugi, 2011. Fatty acid elongase is required for shoot development in rice. *Plant J.*, 66: 680–688
- Jackson, L.L., 1971. Beta-diketone, alcohol and hydrocarbons of barley surface lipids. *Phytochemistry*, 10: 487–490
- Javelle, M., V. Vernoud, N. Depège-Fargeix, C. Arnould, D. Oursel, F. Domergue, X. Sarda and P.M. Rogowsky, 2010. Overexpression of the epidermis-specific homeodomain-leucine zipper IV transcription factor Outer Cell Layer1 in maize identifies target genes involved in lipid metabolism and cuticle biosynthesis. *Plant Physiol.*, 154: 273–286
- Javelle, M., V. Vernoud, P.M. Rogowsky and G.C. Ingram, 2011. Epidermis: the formation and functions of a fundamental plant tissue. *New Phytol.*, 189: 17–39
- Jeffree, C.E., 1996. Structure and ontogeny of plant cuticles. In: *Plant Cuticles: An Integrated Functional Approach*, pp: 33–82. Kersteins, G. (ed.). BIOS Scientific Publishers Ltd, Oxford, UK
- Jenks, M.A., H.A. Tuttle, S.D. Eigenbrode and K.A. Feldmann, 1995. Leaf epicuticular waxes of the eceriferum mutants in Arabidopsis. *Plant Physiol.*, 108: 369–377
- Jetter, R. and M. Riederer, 2000. Composition of cuticular waxes on *Osmunda regalis* fronds. *J. Chem. Ecol.*, 26: 399–412
- Jetter, R., L. Kunst and A.L. Samuels, 2006. Composition of plant cuticular waxes. In: *Biology of the Plant Cuticle*, Vol. 23, pp: 145–181. M. Riederer and C. Müller (eds.). Blackwell, Oxford, UK
- Jung, K.H., M.J. Han, D.y. Lee, Y.S. Lee, L. Schreiber, R. Franke, A. Faust, A. Yephremov, H. Saedler, Y.W. Kim, I. Hwang and G. An, 2006. *Wax-deficient anther1* is involved in cuticle and wax production in rice anther walls and is required for pollen development. *Plant Cell*, 18: 3015–3032
- Kim, H., S.B. Lee, H.J. Kim, M.K. Min, I. Hwang and M.C. Suh, 2012. Characterization of Glycosyl phosphatidylinositol-Anchored Lipid Transfer Protein2 (*LTPG2*) and overlapping function between *LTPG/LTPG1* and *LTPG2* in cuticular wax export or accumulation in *Arabidopsis thaliana*. *Plant Cell Physiol.*, 53: 1391–1403
- Kim, K.S., S.H. Park and M.A. Jenks, 2007. Changes in leaf cuticular waxes of sesame (*Sesamum indicum* L.) plants exposed to water deficit. *J. Plant Physiol.*, 164: 1134–1143
- Knoche, M., S. Peschel, M. Hinz and M.J. Bukovac, 2000. Studies on water transport through the sweet cherry fruit surface: characterizing conductance of the cuticular membrane using pericarp segments. *Planta*, 212: 127–135
- Kosma, D.K., B. Bourdenx, A. Bernard, E.P. Parsons, S. Lu, J. Joubès and M.A. Jenks, 2009. The impact of water deficiency on leaf cuticle lipids of Arabidopsis. *Plant Physiol.*, 151: 1918–1929
- Kosma, D.K., E.P. Parsons, T. Isaacson, S. Lü, J.K. Rose, M.A. Jenks, 2010. Fruit cuticle lipid composition during development in tomato ripening mutants. *Physiol. Plant.*, 139: 107–117
- Kunst, L. and L. Samuels, 2009. Plant cuticles shine: advances in wax biosynthesis and export. *Curr. Opin. Plant Biol.*, 12: 721–727
- Kunst, L., A.L. Samuels and R. Jetter, 2005. The plant cuticle: formation and structure of epidermal surfaces. In: *Plant Lipids—Biology, Utilisation and Manipulation*, pp: 270–302. D. Murphy (ed.). Blackwell, Oxford, UK
- Kurdyukov, S., A. Faust, C. Nawrath, S. Bär, D. Voisin, N. Efremova, R. Franke, L. Schreiber, H. Saedler, J.P. Métraux and A. Yephremov, 2006. The epidermis-specific extracellular *BODYGUARD* controls cuticle development and morphogenesis in Arabidopsis. *Plant Cell*, 18: 321–339

- Laila, R., A.H. Robin, K. Yang, J.I. Park, M.C. Suh, J. Kim and I.S. Nou, 2017. Developmental and genotypic variation in leaf wax content and composition, and in expression of wax biosynthetic genes in *Brassica oleracea* var. capitata. *Front. Plant Sci.*, 7: 1972
- Lee, J., K. Yang, M. Lee, S. Kim, J. Kim, S. Lim, G.H. Kang, S.R. Min, S.J. Kim, S.U. Park, Y.S. Jang, S.S. Lim and H.R. Kim, 2015. Differentiated cuticular wax content and expression patterns of cuticular wax biosynthetic genes in bloomed and bloomless broccoli (*Brassica oleracea* var. italica). *Process Biochem.*, 50: 456–462
- Lee, S.B. and M.C. Suh, 2015. Advances in the understanding of cuticular waxes in *Arabidopsis thaliana* and crop species. *Plant Cell Rep.*, 34: 557–572
- Lee, S.B., H. Kim, R.J. Kim and M.C. Suh, 2014. Overexpression of *Arabidopsis* MYB96 confers drought resistance in *Camelina sativa* via cuticular wax accumulation. *Plant Cell Rep.*, 33: 1535–1546
- Lemieux, B., 1996. Molecular genetics of epicuticular waxes biosynthesis. *Trends Plant Sci.*, 1: 312–318
- Li, L., D. Li, S. Liu, X. Ma, C.R. Dietrich, H.C. Hu, G. Zhang, Z. Liu, J. Zheng, G. Wang and P.S. Schnable, 2013. The maize *glossy13* gene, cloned via BSR-Seq and Seq-Walking encodes a putative ABC transporter required for the normal accumulation of epicuticular waxes. *PLoS One*, 8: e82333
- Li-Beisson, Y., B. Shorrosh, F. Beisson, F.X. Andersson, V. Arondel, P.D. Bates, S. Baud, D. Bird, A. DeBono, T.P. Durrett, R.B. Franke, I.A. Graham, K. Katayama, A.A. Kelly, T. Larson, J.E. Markham, M. Miquel, I. Molina, I. Nishida, O. Rowland, L. Samuels, K.M. Schmid, H. Wada, R. Welti, C. Xu, R. Zallot and J. Ohlrogge, 2013. Acyl-lipid metabolism. *Arabidopsis Book*, 11: e0161
- Liu, F., X. Xiong, L. Wu, D. Fu, A. Hayward, X. Zeng, Y. Cao, Y. Wu, Y. Li and G. Wu, 2014. *BraLTP1*, a lipid transfer protein gene involved in epicuticular wax deposition, cell proliferation and flower development in *Brassica napus*. *PLoS One*, 9: e110272
- Liu, S., C.R. Dietrich and P.S. Schnable, 2009. DLA-based strategies for cloning insertion mutants: cloning the *gl4* locus of maize using Mu transposon tagged alleles. *Genetics*, 183: 1215–1225
- Lü, S., H.Y. Zhao, E.P. Parsons, C.C. Xu, D.K. Kosma, X. Xu, D. Chao, G. Lohrey, D.K. Bangarusamy, G. Wang, R.A. Bressan and M.A. Jenks, 2012. The glossyhead1 allele of *ACC1* reveals a principal role for multidomain acetyl-coenzyme A carboxylase in the biosynthesis of cuticular waxes by *Arabidopsis*. *Plant Physiol.*, 157: 1079–1092
- Lü, S., T. Song, D.K. Kosma, E.P. Parsons, O. Rowland and M.A. Jenks, 2009. *Arabidopsis* *CER8* encodes long-chain acyl CoA synthetase 1 (*LACS1*) that has overlapping functions with *LACS2* in plant wax and cutin synthesis. *Plant J.*, 59: 553–564
- Mao, B., Z. Cheng, C. Lei, F. Xu, S. Gao, Y. Ren, J. Wang, X. Zhang, J. Wang, F. Wu, X. Guo, X. Liu, C. Wu, H. Wang and J. Wan, 2012. *Wax crystal-sparse leaf2*, a rice homologue of *WAX2/GL1*, is involved in synthesis of leaf cuticular wax. *Planta*, 235: 39–52
- McNevin, J.P., W. Woodward, A. Hannoufa, K.A. Feldmann and B. Lemieux, 1993. Isolation and characterization of *eceriferum* (*cer*) mutants induced by T-DNA insertions in *Arabidopsis thaliana*. *Genome*, 36: 610–618
- Nawrath, C., L. Schreiber, R.B. Franke, N. Geldner, J.J. Reina-Pinto and L. Kunst, 2013. Apoplastic diffusion barriers in *Arabidopsis*. *Arabidopsis Book*, 11: e0167
- Post-Beittenmiller, D., 1996. Biochemistry and molecular biology of wax production in plants. *Annu. Rev. Plant Biol.*, 47: 405–430
- Pu, Y., J. Gao, Y. Guo, T. Liu, L. Zhu, P. Xu, B. Yi, J. Wen, J. Tu, C. Ma, T. Fu, J. Zou and J. Shen, 2013. A novel dominant glossy mutation causes suppression of wax biosynthesis pathway and deficiency of cuticular wax in *Brassica napus*. *BMC Plant Biol.*, 13: 1
- Qin, B., D. Tang, J. Huang, M. Li, X. Wu and L. Lu, 2011. Rice *OGLI-1* is involved in leaf cuticular wax and cuticle membrane. *Mol. Plant*, 4: 985–995
- Racovita, R.C., S. Hen-Avivi, J.P. Fernandez-Moreno, A. Granel, A. Aharoni, R. Jetter, 2016. Composition of cuticular waxes coating flag leaf blades and peduncles of *Triticum aestivum* cv. Bethlehem. *Phytochemistry*, 130: 182–192
- Razeq, F.M., D.K. Kosma, O. Rowland and I. Molina, 2014. Extracellular lipids of *Camelina sativa*: characterization of chloroform-extractable waxes from aerial and subterranean surfaces. *Phytochemistry*, 106: 188–196
- Samuels, L., L. Kunst and R. Jetter, 2008. Sealing plant surfaces: cuticular wax formation by epidermal cells. *Plant Biol.*, 59: 683–707
- Schneider, L.M., N.M. Adamski, C.E. Christensen, D.B. Stuart, S. Vautrin, M. Hansson, C. Uauy and P. von Wettstein-Knowles, 2016. The *Cer-cqu* gene cluster determines three key players in a  $\beta$ -diketone synthase polyketide pathway synthesizing aliphatics in epicuticular waxes. *J. Exp. Bot.*, 67: 2715–2730
- Seo, P.J., S.B. Lee, M.C. Suh, M.J. Park, Y.S. Go and C.M. Park, 2011. The *MYB96* transcription factor regulates cuticular wax biosynthesis under drought conditions in *Arabidopsis*. *Plant Cell*, 23: 1138–1152
- Smimova, A., J. Leide and M. Riederer, 2013. Deficiency in a very-long-chain fatty acid  $\beta$ -ketoacyl-CoA synthase (*SICER6*) of tomato impairs micro gametogenesis and causes floral organ fusion. *Plant Physiol.*, 161: 196–209
- Smith, R.M., J.A. Marshall, M.R. Davey, K.C. Lowe, J.B. Power, 1996. Comparison of volatiles and waxes in leaves of genetically engineered tomatoes. *Photochemistry*, 43: 753–758
- Stiles, W., 1994. *Principles of Plant Physiology*. Discovery Publishing House, New Delhi, India
- Sturaro, M., H. Hartings, E. Schmelzer, R. Velasco, F. Salamini and M. Motto, 2005. Cloning and characterization of *GLOSSY1*, a maize gene involved in cuticle membrane and wax production. *Plant Physiol.*, 138: 478–489
- Suh, M.C., A.L. Samuels, R. Jetter, L. Kunst, M. Pollard, J. Ohlrogge and F. Beisson, 2005. Cuticular lipid composition, surface structure, and gene expression in *Arabidopsis* stem epidermis. *Plant Physiol.*, 139: 1649–1665
- Szafrańek, B.M. and E.E. Synak, 2006. Cuticular waxes from potato (*Solanum tuberosum*) leaves. *Phytochemistry*, 67: 80–90
- Tassone, E.E., A.E. Lipka, P. Tomasi, G.T. Lohrey, W. Qian, J.M. Dyer, M.A. Gore and M.A. Jenks, 2016. Chemical variation for leaf cuticular waxes and their levels revealed in a diverse panel of *Brassica napus* L. *Ind. Crop Prod.*, 79: 77–83
- Tomasi, P., H. Wang, G.T. Lohrey, S. Park, J.M. Dyer, M.A. Jenks, H. Abdel-Haleem, 2017. Characterization of leaf cuticular waxes and cutin monomers of *Camelina sativa* and closely-related *Camelina* species. *Ind. Crop Prod.*, 98: 130–138
- Tulloch, A.P. and R.O. Weenink, 1969. Composition of the leaf wax of Little Club wheat. *Can J. Chem.*, 47: 3119–3126
- Velasco, R., C. Korfhage, A. Salamini, E. Tacke, J. Schmitz, M. Motto, F. Salamini, H.P. Doring, 2002. Expression of the *glossy2* gene of maize during plant development. *Maydica*, 47: 71–81
- Viougeas, M.A., R. Rohr and A. Chamel, 1995. Structural changes and permeability of ivy (*Hedera helix* L.) leaf cuticles in relation to leaf development and after selective chemical treatments. *New Phytol.*, 130: 337–348
- Vogg, G., S. Fischer, J. Leide, E. Emmanuel, R. Jetter, A.A. Levy and M. Riederer, 2004. Tomato fruit cuticular waxes and their effects on transpiration barrier properties: functional characterization of a mutant deficient in a very-long-chain fatty acid  $\beta$ -ketoacyl-CoA synthase. *J. Exp. Bot.*, 55: 1401–1410
- von Wettstein-Knowles, P., 1971. The molecular phenotypes of the *eceriferum* mutants. In: *Barley genetics II*. pp: 146–193. Nilan, R.A. (ed.). Washington State University Press, Pullman, Washington, USA
- von Wettstein-Knowles, P., 1995. Biosynthesis and genetics of waxes. In: *Waxes: Chemistry, Molecular Biology and Functions*, pp: 91–130. Hamilton, R.J. (ed.). Oily Press, Dundee, UK
- Wang, M., H. Wu, J. Xu, C. Li, Y. Wang and Z. Wang, 2017. Five Fatty Acyl-Coenzyme A Reductases Are Involved in the Biosynthesis of Primary Alcohols in *Aegilops tauschii* leaves. *Front. Plant Sci.*, 8: 1012
- Wang, W., X. Liu, X. Gai, J. Ren, X. Liu, Y. Cai, Q. Wang and H. Ren, 2015a. *Cucumis sativus* L. *WAX2* Plays a Pivotal Role in Wax Biosynthesis, Influencing Pollen Fertility and Plant Biotic and Abiotic Stress Responses. *Plant Cell Physiol.*, 56: 1339–1354

- Wang, X., Y. Guan, D. Zhang, X. Dong, L. Tian and Q. LQ, 2017. A  $\beta$ -Ketoacyl-CoA Synthase Is Involved in Rice Leaf Cuticular Wax Synthesis and Requires a CER2-LIKE Protein as a Cofactor1. *Plant Physiol.*, 173: 944–955
- Wang, M., Y. Wang, H. Wu, J. Xu, T. Li, D. Hegebarth, R. Jetter, L. Chen and Z. Wang, 2016. Three *TaFAR* genes function in the biosynthesis of primary alcohols and the response to abiotic stresses in *Triticum aestivum*. *Sci. Rep.*, 6: 25008
- Wang, W., Y. Zhang, C. Xu, J. Ren, X. Liu, K. Black, X. Gai, Q. Wang and H. Ren, 2015b. Cucumber *ECERIFERUM1* (*CsCER1*), which influences the cuticle properties and drought tolerance of cucumber, plays a key role in VLC alkanes biosynthesis. *Plant Mol. Biol.*, 87: 219
- Wang, Y., J. Wang, G. Chai, C. Li, Y. Hu, X. Chen and Z. Wang, 2015a. Developmental changes in composition and morphology of cuticular waxes on leaves and spikes of glossy and glaucous wheat (*Triticum aestivum* L.). *PLoS One*, 10: e0141239
- Wang, Y., M.L. Wang, Y.L. Sun, D. Hegebarth, T.T. Li, R. Jetter and Z. Wang, 2015b. Molecular characterization of *TaFAR1* involved in primary alcohol biosynthesis of cuticular wax in hexaploid wheat. *Plant Cell Physiol.*, 56: 1944–1961
- Wang, Y., M.L. Wang, Y.L. Sun, Y.T. Wang, T.T. Li, G.Q. Chai, W. Jiang, L. Shan, C. Li, E. Xiao and Z. Wang, 2015c. *FAR5*, a fatty acyl-coenzyme A reductase, is involved in primary alcohol biosynthesis of the leaf blade cuticular wax in wheat (*Triticum aestivum* L.). *J. Exp. Bot.*, 66: 1165–1178
- Wang, Z., O. Guhling, R. Yao, F. Li, T.H. Yeats, J.K.C. Rose and R. Jetter, 2011. Two oxidosqualene cyclases responsible for biosynthesis of tomato fruit cuticular triterpenoids. *Plant Physiol.*, 155: 540–552
- Wen, M., C. Buschhaus and R. Jetter, 2006. Nanotubules on plant surfaces: Chemical composition of epicuticular wax crystals on needles of *Taxusbaccata* L. *Phytochemistry*, 67: 1808–1817
- Weng, H., I. Molina, J. Shockey and J. Browse, 2010. Organ fusion and defective cuticle function in a *lacs1lacs2* double mutant of Arabidopsis. *PLANTA*, 231: 1089–1100
- Wink, M., 1988. Plant breeding: importance of plant secondary metabolites for protection against pathogens and herbivores. *Theor. Appl. Genet.*, 75: 225–233
- Xia, K., X. Ou, C. Gao, H. Tang, Y. Jia, R. Deng, X. Xu and M. Zhang, 2015. *OsWS1* involved in cuticular wax biosynthesis is regulated by *osa-miR1848*. *Plant Cell Environ.*, 38: 2662–2673
- Xu, X., C.R. Dietrich, M. Delledonne, Y. Xia, T.J. Wen, D.S. Robertson, B.J. Nikolau and P.S. Schnable, 1997. Sequence analysis of the cloned *GLOSSY8* gene of maize suggests that it may code for a  $\beta$ -ketoacyl reductase required for the biosynthesis of cuticular waxes. *Plant Physiol.*, 115: 501–510
- Yeats, T.H. and J.K. Rose, 2013. The formation and function of plant cuticles. *Plant Physiol.*, 163: 5–20
- Yu, D., K. Ranathunge, H. Huang, Z. Pei, R. Franke, L. Schreiber and C. He, 2008. *Wax Crystal-Sparse Leaf1* encodes a  $\beta$ -ketoacyl CoA synthase involved in biosynthesis of cuticular waxes on rice leaf. *PLANTA*, 228: 675–685
- Zhang, H., O. Rowland and L. Kunst, 2005. Disruptions of the Arabidopsis enoyl-CoA reductase gene reveal an essential role for very-long-chain fatty acid synthesis in cell expansion during plant morphogenesis. *Plant Cell*, 17: 1467–1481
- Zhang, X., Z. Liu, Q. Wang, S. Yang and H. Feng, 2013a. Fine mapping of *BrWax1*, a gene controlling cuticular wax biosynthesis in Chinese cabbage (*Brassica rapa* L. ssp. *pekinensis*). *Mol. Breeding*, 32: 867–874
- Zhang, Z., W. Wang and W. Li, 2013b. Genetic interactions underlying the biosynthesis and inhibition of  $\beta$ -diketones in wheat and their impact on glaucousness and cuticle permeability. *PLoS One*, 8: e54129
- Zhang, Z., W. Wei, H. Zhu, G.S. Challa, C. Bi, H.N. Trick and W. Li, 2015. *W3* Is a New Wax Locus That Is Essential for Biosynthesis of  $\beta$ -Diketone, Development of Glaucousness, and Reduction of Cuticle Permeability in Common Wheat. *PLoS One*, 10: e0140524
- Zhang, Z., Z.J. Chen, L. Gan, H. Zhang, F.Q. Wu, Q.B. Lin, J.L. Wang, J. Wang, X.P. Guo, X. Zhang, Z.C. Zhao, C.L. Lei, S.S. Zhu, C.M. Wang and J.M. Wan, 2016. *OsHSD1*, a hydroxysteroid dehydrogenase, is involved in cuticle formation and lipid homeostasis in rice. *Plant Sci.*, 249: 35–45
- Zhou, L.Y., E.D. Ni, J.W. Yang, H. Zhou, H. Liang, J. Li, D. Jiang, Z. Wang, Z. Liu and C. Zhuang, 2013. Rice *OsGLI-6* is involved in leaf cuticular wax accumulation and drought resistance. *PLoS One*, 8: e65139
- Zhou, Q., C. Li, K. Mishina, J. Zhao, J. Zhang, R. Duan, X. Ma, A. Wang, Q. Meng, T. Komatsuda and G. Chen, 2017. Characterization and genetic mapping of the  $\beta$ -diketone deficient *eceriferum-b* barley mutant. *Theor Appl. Genet.*, 130: 1169
- Zhou, X., L. Li, J. Xiang, G. Gao, F. Xu, A. Liu, X. Zhang, Y. Peng, X. Chen and X. Wan, 2015. *OsGLI-3* is involved in Cuticular Wax Biosynthesis and Tolerance to Water Deficit in Rice. *PLoS One*, 10: e116676
- Zhu, X. and L. Xiong, 2013. Putative megaenzyme *DWA1* plays essential roles in drought resistance by regulating stress-induced wax deposition in rice. *Proc. Natl. Acad. Sci. USA*, 110: 17790–17795
- Zygadlo, J.A., D.M. Maestri and N.R. Grosso, 1994. Alkane distribution in epicuticular wax of some Solanaceae species. *Biochem. Syst. Ecol.*, 22: 203–209

(Received 24 October 2018; Accepted 24 November 2018)