



Full Length Article

Genetic Diversity between and within *Lilium* Species and Hybrids by ISSR Markers

Ming-hong Chen^{1*}, Xin She¹, Duo Liu¹, Yu Li¹ and Cheng-yun Li²

¹School of Ethnic Medicine, Yunnan Minzu University, Kunming 650500, China

²Key Laboratory of Agro-Biodiversity and Pest Management of Education Ministry of China, Yunnan Agricultural University, Kunming 650201, China

*For correspondence: cmhkm99@163.com

Abstract

Genetic variation and relationships among 22 *Lilium* accessions including 14 wild species and 8 hybrids were examined by using inter simple sequence repeat (ISSR) markers. Among 100 primers used for PCR amplification of genomic DNA, 10 primers (10%) generated polymorphic DNA fragments. A total of 148 ISSR fragments were generated from 10 primers, 97.97% of which were polymorphic among the 22 *Lilium* accessions. The genetic variation revealed by ISSR markers indicated that 22 *Lilium* accessions showed high polymorphism, abundant genetic variation, and the genetic diversity within wild species was higher than that within hybrids. Cluster analysis based on ISSR data by Unweighted Pair Group Method with Arithmetic Means (UPGMA) showed that 22 *Lilium* accessions could be placed in two groups with a similarity ranging from 0.4459 to 0.9392. This revealed that wild species were genetically distant from hybrids, and was basically consistent with the results of traditional classification based on morphological characters in *Lilium*. The results of this study revealed that molecular techniques might be helpful or even necessary to the precise classification of *Lilium* and the necessity of further study on the genetic relationships between wild *Lilium* species and hybrids. At the same time, the results also revealed that *L. wenshanense* might be of great value to new variety breeding of *Lilium*. This study will provide the basis for the systematic study and the rational utilization of *Lilium* germplasm resources. © 2019 Friends Science Publishers

Keywords: *Lilium*; Germplasm resources; Genetic diversity; Relationship; ISSR; Molecular marker

Introduction

The genus *Lilium* contains around 110 accepted species native to cold and temperate regions of the Northern Hemisphere between 10 and 60° latitudes (Bakhshaie *et al.*, 2016). Lilies are herbaceous perennials that form underground scaly bulbs (Lee *et al.*, 2011b), and have unsurpassed beauty and great commercial significance (Van Tuyl *et al.*, 2011). Lilies have been cultivated as food and medicine in China for at least 2,000 years (Haw, 1986). Many hybrids have been developed through conventional breeding by utilizing several *Lilium* species, and hybrids showing hetero genetic characteristics in flower color, shape of flowers, and various morphological characters (Lee *et al.*, 1996). At present, *Lilium* hybrids are one of the most important cut flowers and potted plants, wild *Lilium* species and hybrids are also used as large, showy flower garden plants in the world (Van Tuyl *et al.*, 2011).

Discrimination of *Lilium* germplasm has been done by using morphological markers, isozyme markers and molecular markers (Yamagishi, 1995; Bakhshaie *et al.*, 2016; Kumari *et al.*, 2018). Among them, morphological markers are unstable under the influence of environment,

and the use of isozyme markers is limited by the number of informative markers (Yamagishi, 1995). However, molecular markers may be useful tools for these purposes because of circumventing the above-mentioned disadvantages. Researches on genetic diversity and relationships in *Lilium* by using molecular markers are of great theoretical significance and practical value in molecular identification, genetic improvement, germplasm conservation, construction of core germplasm, gene location of important agronomic traits, molecular marker assisted breeding of *Lilium* germplasm resources, etc.

As a microsatellite-based technique, ISSR is a widely used PCR-based molecular marker system with the advantages of easy scoring, high cost-effectiveness, high sensitivity and good reproducibility (Tang *et al.*, 2014). It combines the universality of random amplified polymorphic DNA (RAPD) with most of the advantages of simple sequence repeat (SSR) and amplified fragment length polymorphism (AFLP) etc. (Bornet and Branchard, 2001) It is now broadly used by researchers in various fields of plant biology (Godwin *et al.*, 1997; Abe *et al.*, 2002; Reddy *et al.*, 2002; Arzate-Fernández *et al.*, 2005; Wu *et al.*, 2015; Chaudhary *et al.*, 2018).

In the present study, we presented the use of ISSR as genetic markers for 22 *Lilium* accessions including 14 wild species and 8 hybrids. This study aimed to analyze the genetic variation and relationships of *Lilium*, thus providing the basis for the systematic study and the rational utilization of *Lilium* germplasm resources.

Materials and Methods

Plant Materials

Fourteen *Lilium* species (*L. lancifolium*, *L. pumilum*, *L. davidii*, *L. davidii* var. *unicolor*, *L. regale*, *L. lophophorum*, *L. dauricum*, *L. sargentiae*, *L. henryi*, *L. taliense*, *L. leucanthum*, *L. distichum* and *L. auratum*) and eight hybrids (Tigerwoods *L. wenshanense*, Siberia, Starfighter, Sorbonne, Robina, Royal Trinity, Conca D'or and Yelloween) were used as experimental materials in the present study (Table 1). They were collected from Yunnan, Sichuan, Gansu, Hubei, Heilongjiang, Jilin province. All samples were authenticated based on the morphological characters (Wang and Tang, 1980; Long *et al.*, 1999), and stored in the molecular biology laboratory, School of Ethnic Medicine, Yunnan Minzu University, China.

DNA Isolation and PCR Amplification

DNA samples were isolated from silica-gel dried leaf tissue using a modified CTAB procedure (Yamagishi *et al.*, 2002). PCR was carried out in a 25 μ L reaction mixture containing 40 ng of template DNA, 2.0 mM MgCl₂, 1.5 units of *Taq* DNA polymerase, 200 μ M each of dCTP, dGTP, dATP and dTTP, 800nM of primer (with reference to the ISSR primer sequences published on www.UBC.ca.com) and to Yamagishi's 3A ISSR primer sequences (Yamagishi *et al.*, 2002), primers synthesised by Sangon Biotech (Shanghai) Co., Ltd., 1 \times reaction buffer. DNA amplification was performed by a programmed temperature control system (Eppendorf AG22331 Hamburg, Germany). The thermal cycle program consisted of three steps: 94°C for 5 min; 35 cycles of 94°C for 1 min, 51.8°C for 1 min and 72°C for 2 min; 72°C for 8 min, followed by maintaining the mixture at 4°C. Amplification products were analyzed by electrophoresis on 1.5% agarose gels using 0.5 \times TBE buffer at a constant 50V and then visualized by staining with ethidium bromide. Total genomic DNA extraction and PCR reactions were carried out three times, and only reproducible and well-defined bands were scored (Pérez de la Torre *et al.*, 2012).

Statistical Analysis

For ISSR analysis, the banding patterns were recorded by a gel imaging system (Syngene, GeneGenius, Cambridge, UK). Bands with the same mobility and molecular weight

were treated as identical fragments. In the data matrices, presence or absence of each ISSR fragment was coded as "1" or "0" for each of the primer pairs (Huang and Sun, 2000). With the amplification data of all informational primers, the similarity was estimated according to the number of shared amplification products (Nei, 1978) by using POPGEN software (Version1.32) (Yeh *et al.*, 2002), and a dendrogram based on similarity coefficients was generated by using NTSYSpc software (Version2.1) (Rohlf, 2000) according to UPGMA.

Results

Primer Screening and PCR Amplification

One hundred primers were screened using the template DNA from *L. lancifolium*, *L. davidii*, Siberia and Sorbonne to identify primers that generated informative arrays of PCR products, and 10 informative primers were finally selected for further analysis in this study. The selected 10 primers generated from 11 to 23 bands with an average of 14.8 bands per primer, while the proportion of polymorphic bands varied from 85.71 to 100% (Table 2). Two examples of PCR amplification of DNA samples from the 22 *Lilium* accessions with primers UBC835 and 3A25 were showed respectively (Fig. 1–2).

Genetic Diversity Analysis

ISSR analysis using the 10 informative primers generated a total of 148 DNA bands in 22 *Lilium* accessions, and the percentage of polymorphic loci (*PPL*) was 97.97%. On the species level, the mean observed allele (*Ao*) was 1.9797, the mean effective allele (*Ae*) was 1.6688, the Nei's gene diversity (*H*) was 0.3786, the shannon's information index (*I*) was 0.5558, indicating that all 22 *Lilium* accessions showed high polymorphism, abundant genetic variation. At the same time, the genetic variation within 14 wild species was compared with that within 8 hybrids, and all index values of genetic diversity indicated that the genetic diversity within wild species was higher than that within hybrids (Table 3).



Genetic Similarity Analysis

The similarity matrix obtained by multivariate analysis using Nei's coefficient is shown in Table 4. From the similarity matrix, the similarity coefficients ranged from 0.4459 to 0.9392 among the samples. The results indicated that *L. lophophorum* was the most closely related to *L. dauricum* due to both belonging to Sect. *Lophophorum*. Among wild species, *L. lophophorum* was the most closely related to *L. dauricum*. Among hybrids, Tigerwoods was the most closely related to Siberia, similarity coefficient between them was 0.8581.

Table 1: The experimental materials used for the ISSR analysis

No.	Name	Lines	No.	Name	Lines
1	<i>L. lancifolium</i>	W	12	<i>L. leucanthum</i>	W
2	<i>L. pumilum</i>	W	13	<i>L. distichum</i>	W
3	<i>L. davidii</i>	W	14	<i>L. auratum</i>	W
4	<i>L. davidii</i> var. <i>unicolor</i>	W	15	Tigerwoods	O
5	<i>L. regale</i>	W	16	Siberia	O
6	<i>L. lophophorum</i>	W	17	Starfighter	O
7	<i>L. dauricum</i>	W	18	Sorbonne	O
8	<i>L. sargentiae</i>	W	19	Robina	O/T
9	<i>L. henryi</i>	W	20	Royal Trinity	L/A
10	<i>L. wenshanense</i>	W	21	Conca D'or	O/T
11	<i>L. taliense</i>	W	22	Yelloween	O/T

Note: W: Wild species; O: Oriental hybrids; L/A: *Longiflorum*×Asiatic hybrids; O/T: Oriental × Trumpet hybrids

**Fig. 1:** Electrophoresis of ISSR-PCR with primer UBC835
M: Marker DL2000; 1-22: 22 *Lilium* accessions**Fig. 2:** Electrophoresis of ISSR-PCR with primer 3A25
M: Marker DL2000; 1-22: 22 *Lilium* accessions

Clustering Analysis

The similarity matrix above was used to generate a dendrogram (Fig. 3) by UPGMA analysis in order to determine the grouping of different accessions. The result illustrated that the 22 *Lilium* accessions could be placed in two groups (A and B) at a threshold value of 0.57. The first group (A) contained all wild species and could be further separated into three subgroups at a threshold value of 0.59. The first subgroup (A₁) contained five species in which *L. lancifolium*, *L. davidii*, *L. davidii* var. *unicolor* and *L. henryi* belonging to Sect. *Sinomartagon*. The second subgroup (A₂) contained two species (*L. regale* and *L. sargentiae*) both belonging to Sect. *Lilium*. The third subgroup (A₃)

contained seven species in which *L. pumilum* and *L. taliense* belonging to Sect. *Sinomartagon*, *L. lophophorum* and *L. dauricum* belonging to Sect. *Lophophorum*, *L. leucanthum* belonging to Sect. *Lilium* and *L. distichum* belonging to Sect. *Martagon*, respectively. The second group (B) contained all hybrids and could be further separated into two subgroups at a threshold value of 0.627. The first subgroup (B₁) contained all the Oriental hybrids (Tigerwoods, Siberia, Starfighter and Sorbonne). The second subgroup (B₂) contained all the Oriental×Trumpet hybrids (Robina, Conca D'or and Yelloween) and Royal Trinity belonging to *Longiflorum*×Asiatic hybrids. In 22 *Lilium* accessions, all wild species and all hybrids were placed in two different groups, indicating that wild species were genetically distant from hybrids.

Discussion

Breeders have realized that a large amount of genetic variation within crop plants exists unexploited in wild germplasm as a result of domestication (Acosta-Gallegos *et al.*, 2007), these wild relatives can play important roles on the *Lilium* breeding. In China, *Lilium* hybrids are mainly introduced from abroad. As a result of the artificial hybridization that has been going on for nearly 100 years, genetic relationships within hybrids are getting closer and closer, and genetic distances between hybrids and their original wild ancestors are getting larger and larger. Therefore, the genetic background of hybrids is very complicated. In the present study, the genetic variation revealed by ISSR markers indicated that the genetic diversity within wild species was higher than that within hybrids, and cluster analysis based on ISSR data indicated that wild species were genetically distant from hybrids. The results of this study revealed the necessity of further study on the genetic relationships between wild *Lilium* species and hybrids.

For a long time, there have been so many arguments in classification, origin, evolution and other issues of subordinate taxa in *Lilium*. Nowadays, classification of *Lilium* according to Comber (1949) is universally acceptable that the species of *Lilium* are classified into seven sections, ie, *Lilium*, *Martagon*, *Pseudolirium*, *Archelirion*, *Sinomartagon*, *Leucolirion* and *Oxypetala* on basis of their 15 morphological characteristics. Since then, numerous researchers have explored and evaluated the classification methods of *Lilium* (Haruki *et al.*, 1997; Dubouzet and Shinoda, 1999; Hayashi and Kawano, 2000; Nishikawa *et al.*, 1999, 2001; Lee *et al.*, 2011a). On the basis of morphological characteristics such as flower shape, leaf arrangement, the existence form and curvature of stamens, the species of *Lilium* in China could be classified into four sections, i.e., Sect. *Lilium*, Sect. *Lophophorum*, Sect. *Sinomartagon* and Sect. *Martagon* (Wang and Tang, 1980). In the wild species that *L. lancifolium*, *L. davidii*, *L. davidii* var. *unicolor* and *L. henryi*

Table 2: The names and sequences of primers and amplified results

Primer	Primer sequence(5'→3')	Total band number	Polymorphic band number	Percentage of polymorphic band (%)
UBC811	GAGAGAGAGAGAGAGAC	15	15	100
UBC825	ACACACACACACACT	12	12	100
UBC835	AGAGAGAGAGAGAGAYC	15	15	100
UBC842	GAGAGAGAGAGAGAYG	12	11	91.67
UBC895	AGAGTTGGTAGCTCTGATC	14	12	85.71
UBC899	CATGGTGTGGTCAATTGTCCA	11	11	100
3A13	CACACACACACAGTC	17	17	100
3A25	CTCTCTCTCTCTGAC	15	15	100
3A37	CACACACACACATGA	23	23	100
3A39	AGAGAGAGAGAGGCTT	14	14	100
Total		148	145	

Table 3: Genetic variation of 22 *Lilium* accessions detected by ISSR (standard deviation in parentheses)

Taxon	NP	PPL(%)	A_o	A_e	H	I
Wild species	137	96.48	1.9648(0.1850)	1.6475(0.2948)	0.3699(0.1327)	0.5445(0.1677)
Hybrids	104	86.67	1.8667(0.3414)	1.5898(0.3513)	0.3349(0.1680)	0.4928(0.2269)
At species level	145	97.97	1.9797(0.1414)	1.6688(0.2936)	0.3786(0.1286)	0.5558(0.1584)

NP: The number of polymorphic loci; A_o : Mean observed number of alleles; A_e : Mean effective number of alleles; H: Nei's genetic diversity; I: Shannon's information index; PPL: The percentage of polymorphic loci

Table 4: Similarity matrix for Nei's coefficient of 22 *Lilium* accessions obtained from ISSR marker analysis

No.	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
1	0.601	0.678	0.655	0.595	0.554	0.561	0.568	0.601	0.628	0.568	0.541	0.588	0.561	0.588	0.581	0.622	0.588	0.581	0.574	0.520	0.581
2		0.655	0.595	0.507	0.710	0.649	0.547	0.622	0.581	0.682	0.750	0.676	0.676	0.608	0.655	0.574	0.595	0.547	0.581	0.541	0.534
3			0.642	0.595	0.595	0.588	0.554	0.601	0.574	0.541	0.568	0.574	0.534	0.493	0.554	0.527	0.520	0.487	0.493	0.507	0.527
4				0.615	0.628	0.635	0.588	0.608	0.581	0.547	0.628	0.622	0.622	0.569	0.588	0.601	0.581	0.507	0.473	0.487	0.547
5					0.595	0.547	0.703	0.628	0.574	0.568	0.635	0.628	0.601	0.547	0.622	0.595	0.615	0.554	0.601	0.534	0.568
6						0.939	0.541	0.642	0.547	0.662	0.716	0.669	0.655	0.534	0.595	0.554	0.601	0.500	0.534	0.466	0.527
7							0.507	0.581	0.514	0.601	0.669	0.622	0.622	0.527	0.574	0.561	0.595	0.466	0.487	0.446	0.510
8								0.615	0.547	0.554	0.608	0.520	0.561	0.561	0.581	0.541	0.601	0.554	0.520	0.507	0.595
9									0.676	0.628	0.615	0.649	0.554	0.527	0.574	0.574	0.581	0.574	0.568	0.541	0.574
10										0.601	0.547	0.595	0.568	0.595	0.574	0.601	0.635	0.574	0.568	0.554	0.601
11											0.811	0.737	0.819	0.723	0.716	0.608	0.655	0.581	0.642	0.534	0.527
12												0.710	0.804	0.642	0.757	0.595	0.669	0.554	0.601	0.534	0.487
13													0.757	0.568	0.615	0.561	0.595	0.561	0.541	0.541	0.507
14														0.710	0.737	0.628	0.676	0.601	0.595	0.554	0.520
15															0.858	0.831	0.770	0.642	0.662	0.595	0.642
16																0.743	0.804	0.662	0.669	0.615	0.595
17																	0.737	0.581	0.655	0.574	0.635
18																		0.628	0.622	0.581	0.642
19																			0.845	0.750	0.784
20																				0.770	0.710
21																					0.710

belonging to Sect. *Sinomartagon* were clustered into the first subgroup, *L. regale* and *L. sargentiae* belonging to Sect. *Lilium* were clustered into the second subgroup, *L. lophophorum* and *L. dauricum* belonging to Sect. *Lophophorum* were closely related to each other and their similarity coefficient was 0.939 indicated that the result of cluster analysis was basically consistent with that of morphological classification in this study. Although some species are distantly related to each other, but its morphological traits especially flower type are very similar, so it is difficult to classify *Lilium* only based on morphological characteristics (Nishikawa et al., 2001). To some extent, this view was confirmed that the cluster forms of *L. pumilum*, *L. taliense*, *L. auratum*, *L. leucanthum* and *L. distichum* were different from traditional morphological

classification results. That *L. pumilum* and *L. taliense* were distantly related to *L. lancifolium*, *L. davidii*, *L. davidii* var. *unicolor* and *L. henryi*, and were clustered into different subgroups indicated that Sect. *Sinomartagon* was very complicated with strong differentiation within section, and might be diverse origins which were consistent with the previous reports (Nishikawa et al., 1999, 2001; Lee et al., 2011a). The results that the similarity coefficient between *L. sargentiae* and *L. regale* was 0.7027, that between *L. regale* and *L. leucanthum* was 0.6351 and that between *L. sargentiae* and *L. leucanthum* was 0.6081 were basically consistent with the view that *L. sargentiae*, *L. regale* and *L. leucanthum* belonging to Sect. *Lilium* had close relationships and the former two were more closely related to each other (Nishikawa et al., 1999).




Fig. 3: Dendrogram of 22 *Lilium* accessions generated from ISSR data using UPGMA analysis. The scale is based on Nei's coefficients of similarity

At present, there are few reports about *L. wenshanense* which is native to the grassy slope at altitudes ranging from 1,000 to 2,200 m in Wenshan of Yunnan, China. The results that *L. wenshanense* were distantly related to the tested hybrids indicated that *L. wenshanense* might be of great value to new variety breeding of *Lilium* in the future according to its excellent characters such as flower shape and flower color.

Conclusion

All 22 *Lilium* accessions showed high polymorphism, abundant genetic variation, and the results of cluster analysis were basically consistent with those of traditional classification based on morphological characters in *Lilium*. However, the relationships among several species belonging to Sect. *Sinomartagon* were far away, and their clustering in different subgroups indicated that the origin of this Sect. was very complicated. In cluster analysis, seven species belonging to different sections were finally got together in a subgroup, indicating that the precise classification of *Lilium* should be combined with morphological and molecular techniques. At the same time, the genetic variation revealed by ISSR markers indicated that the genetic diversity within wild species was higher than that within hybrids, and cluster analysis based on ISSR data indicated that wild species were genetically distant from hybrids. Results further revealed the necessity of study on the genetic relationships between wild *Lilium* species and hybrids. In addition, *L. wenshanense*, a rarely reported elite wild germplasm, was distantly related to the tested hybrids revealed that it might be of great value to new variety breeding of *Lilium*. This study will provide the basis for the systematic study and the rational utilization of *Lilium* germplasm resources.

Acknowledgements

This research was funded by the National Natural Science Foundation of China (Grant No. 30860161) and the Scientific Research Foundation of the Education Department of Yunnan Province (Grant No. 2013Z040).

References

- Abe, H., M. Nakano, A. Nakatsuka, M. Nakayama, M. Koshioka and M. Yamagishi, 2002. Genetic analysis of floral anthocyanin pigmentation traits in Asiatic hybrid lily using molecular linkagemaps. *Theor. Appl. Genet.*, 105: 1175–1182
- Acosta-Gallegos, J.A., J.D. Kelly and P. Gepts, 2007. Prebreeding in common bean and use of genetic diversity from wild germplasm. *Crop Sci.*, 47: 44–59
- Arzate-Fernández, A.M., M. Miwa, T. Shimada, T. Yonekura and K. Ogawa, 2005. Genetic diversity of Miyamasukashi-yuri (*Lilium maculatum* Thunb. var. *bukosanense*), an endemic and endangered species at Mount Buko, Saitama, Japan. *Plant Spec. Biol.*, 20: 57–65
- Bakhshaie, M., S. Khosravi, P. Azadi, H. Bagheri and J.M. Van Tuyl, 2016. Biotechnological advances in *Lilium*. *Plant Cell Rep.*, 35: 1799–1826
- Bornet, B. and M. Branchard, 2001. Nonanchored inter simple sequence repeat (ISSR) markers: reproducible and specific tools for genome fingerprinting. *Plant Mol. Biol. Rep.*, 19: 209–215
- Chaudhary, V., M. Kumar, S. Sharma, N. Kumar, V. Kumar, H.K. Yadav, S. Sharma and U. Sirohi, 2018. Assessment of genetic diversity and population structure in gladiolus (*Gladiolus hybridus* Hort.) by ISSR markers. *Physiol. Mol. Biol. Plants*, 24: 493–501
- Comber, H.F., 1949. A new classification of the genus *Lilium*. In: *Lily Yearbook*, pp: 86–105. Royal Hort. Soc. (eds.) London, UK
- Dubouzet, J.G. and K. Shinoda, 1999. Phylogenetic analysis of the internal transcribed spacer region of Japanese *Lilium* species. *Theor. Appl. Genet.*, 98: 954–960
- Godwin, I.D., E.A.B. Aitken and L.W. Smith, 1997. Application of inter-simple sequence repeat (ISSR) markers to plant genetics. *Electrophoresis*, 18: 1524–1528
- Haruki, K., T. Hosoki, Y. Nako and K. Ohta, 1997. Possibility of classification in some species of *Lilium* by PCR-RFLP of ribulose-1, 5-bisphosphate carboxylase large subunit (*rbcL*) gene and ribosomal RNA gene. *J. Jpn. Soc. Hortic. Sci.*, 66: 189–192
- Haw, S.G., 1986. *The Lilies of China*. Timber Press, Portland, USA
- Hayashi, K. and S. Kawano, 2000. Molecular systematics of *Lilium* and allied genera (Liliaceae): Phylogenetic relationships among *Lilium* and related genera based on the *rbcL* and *matK* gene sequence data. *Plant Spec. Biol.*, 15: 73–93
- Huang, J.C. and M. Sun, 2000. Genetic diversity and relationships of sweet potato and its wild relatives in Ipomoea series Batatas (Convolvulaceae) as revealed by inter-simple sequence repeat (ISSR) and restriction analysis of chloroplast DNA. *Theor. Appl. Genet.*, 100: 1050–1060
- Kumari, S., B.K. Kanth, Y. Jeon, J.Y. Jang, H.S. Kim and G.J. Lee, 2018. Internal transcribed spacer-based CAPS marker development for *Lilium hansonii* identification from wild *Lilium* native to Korea. *Sci. Hortic.*, 236: 52–59
- Lee, C.S., S.C. Kim, S.H. Yeau and N.S. Lee, 2011a. Major lineages of the genus *Lilium* (Liliaceae) based on nrDNA ITS sequences, with special emphasis on the Korean species. *J. Plant Biol.*, 54: 159–171
- Lee, J.S., P.O. Lee, Y.P. Lim, E.M. Shin, S.Y. Park and M.S. Roh, 1996. Classification of lilies using random amplified polymorphic DNA (RAPD) analysis. *Acta. Hort.*, 414: 137–144
- Lee, S.I., K.C. Park, Y.S. Song, J.H. Son, S.J. Kwon, J.K. Na, J.H. Kim and N.S. Kim, 2011b. Development of expressed sequence tag derived-simple sequence repeats in the genus *Lilium*. *Genes Genom.*, 33: 727–733
- Long, Y.Y., J.Z. Zhang and L.N. Zhang, 1999. *Lily – the King of Bulbous Flowers*. Jindun Publishing House, Beijing, China

- Nei, M., 1978. Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics*, 89: 583–590
- Nishikawa, T., K. Okazaki, K. Arakawa and T. Nagamine, 2001. Phylogenetic analysis of Section *Sinomartagon* in genus *Lilium* using sequences of the internal transcribed spacer region in nuclear ribosomal DNA. *Breed. Sci.*, 51: 39–46
- Nishikawa, T., K. Okazaki, T. Uchino, K. Arakawa and T. Nagamine, 1999. A molecular phylogeny of *Lilium* in the internal transcribed spacer region in nuclear ribosomal DNA. *J. Mol. Evol.*, 49: 238–249
- Pérez de la Torre, M., M. García, R. Heinz and A. Escandón, 2012. Analysis of genetic variability by ISSR markers in *Calibrachoa caesia*. *Electron. J. Biotechnol.*, 15. Available from: <http://dx.doi.org/10.2225/vol15-issue5-fulltext-8>
- Reddy, M.P., N. Sarla and E.A. Siddiq, 2002. Inter simple sequence repeat (ISSR) polymorphism and its application in plant breeding. *Euphytica*, 128: 9–17
- Rohlf, F.J., 2000. *NTSYS-pc: Numerical Taxonomy and Multivariate Analysis System*. Version 2.1. Exceter Software, New York, USA
- Tang, N., G.C. Mo, J.M. Van Tuyl, P. Arens, J.J. Liu and D.C. Tang, 2014. Genetic diversity and structure of *Lilium pumilum* DC. in southeast of Qinghai–Tibet plateau. *Plant Syst. Evol.*, 300: 1453–1464
- Van Tuyl, J.M., P. Arens, M.S. Ramanna, A. Shahin, N. Khan, S. Xie, A. Marasek–Ciolkowska, K.B. Lim and R. Barba–Gonzalez, 2011. *Lilium*. In: *Wild Crop Relatives: Genomic and Breeding Resources*, pp: 161–183. Kole, C. (ed.). Springer, Berlin, Heidelberg, Germany
- Wang, F.Z. and J. Tang, 1980. *Flora of China*. Science Press, Beijing, China
- Wu, Z.H., J. Shi, M.L. Xi, F.X. Jiang, M.W. Deng and S. Dayanandan, 2015. Inter-simple sequence repeat data reveals high genetic diversity in wild populations of the narrowly distributed endemic *Lilium regale* in the Minjiang River Valley of China. *PLoS One*, 10: e0118831
- Yamagishi, M., 1995. Detection of section-specific random amplified polymorphic DNA (RAPD) markers in *Lilium*. *Theor. Appl. Genet.*, 91: 830–835
- Yamagishi, M., H. Abe, M. Nakano and A. Nakatsuka, 2002. PCR-based molecular markers in Asiatic hybrid lily. *Sci. Hortic.*, 96: 225–234
- Yeh, F.C., R. Yang, T.J. Boyle, Z. Ye and J.X. Mao, 2002. *Popgen 32, microsoftware Windows Based Freeware for Population Genetic Analysis*. Molecular Biology and Biotechnology Center, University of Alberta, Edmonton, Canada

(Received 01 December 2018; Accepted 09 January 2019)