



Full Length Article

Isolation, Screening and Fermentation Optimization of *Monascus* Strains with High Monacolin K Yield and the Cholesterol Lowering Effect of Red Yeast Rice

Zhiting Chen^{1,2}, Yihan Zhu³, Xiaohua Cao³, Qinyou Wen^{1,2}, Zixiao Xiong^{1,4}, Zuxin Cheng³, Chuannan Long⁵, Shiqiang Lin³, Xinying Huang³, Jianghong Liu³ and Zhiwei Huang^{1,2,4,*}

¹College of Food Science, Fujian Agriculture and Forestry University, Fuzhou 350002, China

²Fujian Provincial Key Laboratory of Quality Science and Processing Technology in Special Starch, Fujian Agriculture and Forestry University, Fuzhou 350002, China

³Key Laboratory of Crop Biotechnology (Fujian Agriculture and Forestry University), Fujian Province University, Fuzhou 350002, China

⁴China-Ireland International Cooperation Centre for Food Material Science and Structure Design, Fujian Agriculture and Forestry University, Fuzhou 350002, China

⁵Jiangxi Key Laboratory of Bioprocess Engineering, Jiangxi Science and Technology Normal University, Nanchang 330013, China

*For correspondence: hzwfau@163.com

Abstract

Monacolin K (MK), one of the main active ingredients of red yeast rice, has cholesterol-lowering effect. This study aimed to improve MK content in red yeast rice through three aspects including strain and rice material screening and optimization of fermentation conditions. The cholesterol-lowering effect of high MK red yeast rice in mice was also studied. The results showed that: *Monascus* strain FG-2 has the highest MK yields, about 2.3 times of model strain ACCC 30501. Molecular identification reveals the strain as *Monascus purpureus*. When fermented with polished rice of variety "1a825" as substrate, strain FG-2 and the highest MK yield, about 2.7 times of commercially available rice "Dulimi". The optimum fermentation conditions for strain FG-2 were determined by single factor optimization and orthogonal test: pre-fermentation temperature at 30°C, seed solution pH at 5.0, rice amount in fermentation-flask 40 g, fermentation time 17 d. MK yield under this fermentation condition was 4.64 mg/g, 28.5% higher than before optimization, which was about 8.3 times of commercially available Gutian red yeast rice. The high MK red yeast rice prepared herein and commercially available Gutian red yeast rice have significant cholesterol lowering effects, with the former superior to the latter. © 2019 Friends Science Publishers

Keywords: *Monascus*; Monacolin K; Strain screening; Rice raw material; Fermentation condition; Cholesterol lowering

Introduction

Red yeast rice, described as a traditional food and medicine in the Chinese Pharmacopoeia, originated in China thousands of years ago (Erdoğan and Azirak, 2004). As purple-red rice koji made from rice as raw material and fermented by *Monascus*, red yeast rice is a natural food additive benefiting human health (Zlová *et al.*, 1996).

Endo (1979) first extracted an active substance that inhibits cholesterol synthesis in the body from culture of *Monascus ruber* and named it as Monacolin K (MK, also known as lovastatin, mevastatin). Alberts *et al.* (1980) extracted lovastatin from the culture of *Aspergillus terreus*, which has the same substance as MK. MK, an active ingredient in red yeast rice, lowers blood cholesterol (Barrios-Gonzalez *et al.*, 2008), antibiosis (Zhou *et al.*, 2018), colon cancer (Hong *et al.*, 2008), breast cancer (Zhu *et al.*, 2012), anti-oxidation (Dhale *et al.*, 2007; Awad *et al.*, 2017),

lowering prostate (Shannon *et al.*, 2005), liver protection (Wei and Popovich, 2013) and accelerating nerve regeneration (Ghayour *et al.*, 2017). MK can effectively reduce cholesterol in dogs (Alberts *et al.*, 1980), reduce fibrinogen content in blood of high-fat model mice, increase antithrombin III and protein C levels, and reduce aortic lipid plaques (Lee *et al.*, 2013). In clinical trials, MK significantly reduced total cholesterol and low-density lipoprotein cholesterol levels, suggesting that MK also has lipid-lowering effect on humans (Ruscica *et al.*, 2014). MK has been widely used as an effective drug to lower blood cholesterol (Panda *et al.*, 2010), arousing the attention to red yeast rice in medical field. Therefore, how to improve MK content in red yeast rice is still a research hotspot in the field of hypolipidemic drugs.

MK has a highly effective and specific inhibitory effect on rate-limiting enzyme HMG-CoA reductase (HMGR) of the cholesterol synthesis pathway, thereby lowering

cholesterol synthesis (Singgih *et al.*, 2014). Natural MK has two different structures, namely open loop β -hydroxy acid form and closed-loop lactone form (Kennedy *et al.*, 1999; Huang *et al.*, 2010). MK mainly exists in an ester-closed loop structure in naturally fermented red yeast rice, but only MK in open-loop acid structure can bind with HMGR to exert an inhibitory effect. II Gum *et al.* (2017) found that red yeast rice fermented with *Bacillus subtilis* significantly increased acid ester proportion in lovastatin and antioxidant activity (II Gum *et al.*, 2017).

The study on the biosynthesis pathway of MK and lovastatin began with *A. terreus*. In 1999, there were reports on the genes related to lovastatin synthesis in *A. terreus* (Kennedy *et al.*, 1999) and the synthetic pathway of lovastatin in *A. terreus* has been well studied (Guo and Wang, 2014; Yin *et al.*, 2016). Later, the genes of MK synthesis in *Monascus* were also studied. MK is a secondary metabolite synthesized by polyketide synthase (PKS) in *Monascus*. Chen *et al.* (2008) cloned the genome sequence of MK biosynthetic gene cluster from *Monascus pilosus* (GenBank: DQ176595.1), composed of 9 genes *mok A - mok I*, which has similarity of over 54% with the lovastatin biosynthetic gene cluster from *A. terreus*. Then, the core of *mok A* in the wild-type *Monascus pilosus* strain was replaced through homologous recombination, forming strain lacking *mok A*. Such strain did not produce MK, indicating that *mok A* encodes polyketide synthase responsible for MK synthesis in *Monascus pilosus* (Chen *et al.*, 2008). The *mok B* is located at the end of the gene cluster and knockout strain failed to produce MK, but accumulated the intermediate product of MK synthesis pathway, Monacolin J (MJ), indicating that *mok B* encodes polyketide synthase and is responsible for MK side chain diketone synthesis (Sakai *et al.*, 2009). The *mok H* encodes Zn (II)₂ cys₆ dinuclear DNA binding protein as a catalyst for MK synthesis. The *mok H* can play a positive regulatory role in the transcription of MK synthesis-related genes, thereby increasing MK yield (Chen *et al.*, 2010). Lin *et al.* (2018) overexpressed *mok E* in *Monascus fuliginosus* and showed that *mok E* expression level of the transformed strain was 4.8 times higher than the wild type strain and MK yield was increased by 2.5 times. In recent years, high-throughput sequencing technology has also been applied to the study of lovastatin and MK synthesis related genes. Savitha *et al.* (2016) reported whole genome sequence of *A. terreus* (KM017963), including complete sequence of lovastatin biosynthesis gene cluster. Zhang *et al.* (2017) performed de novo RNA sequencing and transcriptome analysis on *Monascus purpureus* cultured for different days, demonstrating that *mok F* plays a regulatory role in MK synthesis, and 9 genes (*mok A - mok I*) are related to MK synthesis.

In addition to association with the genotype of microorganisms, the production of microbial secondary metabolites is largely influenced by environmental conditions. Studies have shown that *A. terreus* uses glycerol and lactose as carbon sources, which is more conducive to the synthesis of lovastatin. However, when only glycerol is used

as a carbon source, geodin will be preferably synthesized (Hasan *et al.*, 2018). This is consistent in *Monascus*. The carbon and nitrogen source types, pH, culture temperature and light of the culture medium will all affect its MK yield (Calvo *et al.*, 2002; Li *et al.*, 2010). Lin *et al.* (2017) studied the effect of culture temperature on the yield of *Monascus fuliginosus* MK, finding that constant temperature culture could promote mycelial growth of *Monascus*, but did not facilitate MK production. However, under the condition of varying temperature, *Monascus* MK synthesis related genes had expression level higher than those of constant temperature culture, with MK yield 16 times higher than constant temperature culture.

There are mainly two ways to increase MK yield of *Monascus*. One is to select the strain with high MK yield, and the other is to screen the optimum fermentation conditions for MK production, such as culture temperature, pH, inoculum size, stirring/ventilation condition, fermentation time, carbon nitrogen ratio (Mulder *et al.*, 2015). In the liquid fermentation process, hyphae tend to accumulate into mycelium pellet and appropriate volume of mycelium pellet promotes MK formation. However, as the fermentation proceeds, the mycelium pellet increases volume and fermentation liquid viscosity, hindering the transportation of nutrients and oxygen so that the interior of the mycelium pellet cannot be replenished in time, causing autolysis and "hollowness" of the interior mycelium pellet, which in turn affects MK generation. Through stirring, the hyphae can be prevented from being too dense, the mycelium pellet volume can be reduced, and the dissolved oxygen can be increased to improve the transportation of nutrients and oxygen in the liquid (Mulder *et al.*, 2015). Studies have shown that red yeast rice produced by solid state fermentation has higher MK yield than liquid fermentation (Zhang *et al.*, 2013, 2015). At present, some studies focus on how to improve MK content in red yeast rice, mainly by transforming the strain, such as: engineering its key genes by molecular means, or performing mutagenesis by physical, chemical methods (Mulder *et al.*, 2015) and even metabolically engineering fungus that could perform de novo synthesis of MJ and MK with methanol as a precursor (Liu *et al.*, 2018). However, there are rare reports on the differences in MK production by *Monascus* fermentation using different varieties of rice materials. In this study, the optimal combination of MK high-yield strains and rice raw materials was screened, fermentation process in MK production was optimized, cholesterol-lowering effect of red yeast rice was studied via mice test to provide experimental basis for preparation and application of high MK red yeast.

Materials and Methods

Experimental Materials

Red yeast powder: 7 kinds in total were purchased from Hebei, Beijing, Gansu, Guangdong and Fujian Province in China.

Monascus strains: 13 strains in total were purchased from Agricultural Culture Collection of China (ACCC), China Center of Industrial Culture Collection (CICC) and Guangdong Microbiology Culture Center (GDMCC) in China.

Rice varieties: a total of 26 varieties were obtained from Key Laboratory of Crop Biotechnology (Fujian Agriculture and Forestry University), Fujian Province University, China.

Test animals: KM strain male mice (clean grade, weight 15-20 g/mice), were purchased from Laboratory Animal Center, Fujian Medical University in China.

Rice Planting and Rice Raw Material Preparation

All the rice varieties were planted in the field of Fujian Agriculture and Forestry University, in Minhou, Fuzhou. The field management was conducted the same as common rice varieties. After the maturation of rice grain, three samples were randomly taken from each variety. The samples were then grained to milled rice and brown rice for later use. The market bought common rice "Dulimi" and "Gutian red yeast rice" were used as controls.

Separation of *Monascus* Strains

The 5 g of commercially available red yeast powder was transferred into a 250 mL flask containing 45 mL sterile physiological saline, and shaken at 150 rpm for 30 min to prepare diluents with a concentration gradient of 10^{-2} , 10^{-3} , 10^{-4} , 10^{-5} and 10^{-6} . The diluents were spread on PDA medium plates and incubated at 28°C for 3-5 d. All single colonies with *Monascus* characteristics were selected and arranged in a "Z" shape on a new PDA plate, cultured at 28°C for 3-5 d and transferred to a new PDA plate. After multiple isolation and purification, it was transferred to PDA slant medium, cultured at 28°C for 5 d, and stored in a 4°C refrigerator for later use.

Preparation and Inoculation of Culture Medium

Preparation of seed solution medium: glucose 50 g, peptone 5 g, yeast extract 1 g, KH_2PO_4 1 g, $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ 0.01 g, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.5 g. Add water to a constant volume of 1000 mL. Each 50 mL of seed culture medium was dispensed into a 250 mL flask, added with eight glass beads and autoclaved at 121°C for 20 min.

Preparation of rice fermentation medium: Wash the rice, soak in water for three hours, steam for 20 min, then each 50 g of rice was dispensed into 300 mL plastic fermentation bottle and autoclaved at 121°C for 20 min.

After the *Monascus* strains were activated and cultured on the PDA plate, the fungus cake was taken with a 5 mm diameter puncher and 5 pieces of the cake were added to a 250 mL flask containing 50 mL of the seed liquid medium (each bottle was put with eight small glass beads) and cultured at 30°C, 150 rpm for 72 h. The seed solution was diluted to a spore concentration of about 1×10^7

cell/mL, added to a fermentation flask containing rice fermentation medium at 10% inoculum size, and cultured in a 28°C incubator for 16 d. There were three repetitions for each treatment.

HPLC Detection of MK Content in Red Yeast Rice

Pretreatment of the sample to be tested: The solid fermentation product was dried at 50°C to constant weight, and ground into powder. The 0.5 g sample to be tested was added to a 10 mL brown volumetric flask and 5 mL of 0.2 mol/L NaOH solution was added and volume was adjusted to 10 mL with methanol and sonicated at 50°C for 1 h. After centrifugation at 8000 g for 10 min at 4°C, 5 mL of supernatant was transferred to the brown volumetric flask and adjusted to 25 mL with methanol. After full shaking, 1 mL of the diluents was taken and passed through 0.45 μm filter to be injected.

HPLC chromatographic conditions: chromatographic column Vision HT C18HL (5 μm , 250 \times 4.6 mm), Grace; mobile phase, acetonitrile: 0.18% phosphoric acid solution = 50:50 (v/v); flow rate 1.0 mL/min; detector wavelength 238 nm; column temperature 28°C; injection volume 10 μL .

Molecular Identification of High-yield MK *Monascus* Strains

Genomic DNA of high-yield MK *Monascus* mycelium was extracted using genomic DNA rapid extraction kit (Beijing Ding Guo Changsheng Biotechnology Company, Ltd., China) and its ITS sequence was amplified with 2 \times Eco Taq PCR SuperMix (TransGen Biotech, AS151). The primers were ITS1 (5'-TCCGTAGGTGAACCTGCGG-3') and ITS4 (5'-TCCTCCGCTTATTGATATGC-3'). The PCR reaction system had a total volume of 50 μL containing 25.0 μL 2 \times Eco Taq PCR SuperMix, 1.0 μL each of the forward and reverse primers (20 μM), 4.0 μL DNA template and 19.0 μL ddH₂O. The PCR reaction conditions were 94°C for 5 min; followed by 35 cycles of 94°C for 30 s, 58°C for 30 s, 72°C for 45 s and a final extension at 72°C for 10 min. The amplified ITS fragment was recovered and sent to BioSune Biotechnology (Shanghai) Co., Ltd. for sequencing with primers ITS1 and ITS4, and the sequence obtained was aligned with the reported ITS sequence in the GenBank database using BLASTN to obtain species information of the high-yield MK *Monascus* strain.

In vivo Cholesterol-lowering Test of High MK Red Yeast Rice

KM mice were divided into five groups: basal feed control group A, hyperlipidemia model control group B, Gutian red yeast powder test group C, homemade red yeast powder test group D, Xuezhikang drug test group E, with 10 mice in each group. At the start of the experiment, mice were grouped by weight to make overall weight of each group as balanced as possible. Each group was fed with basal feed every day and

was free to drink. Except group A, each other group was intraperitoneally injected with 0.2 mL of 75% fresh egg yolk emulsion for 7 consecutive days to establish experimental hyperlipidemia mouse model. Blood was drawn from the mice tail in each group, centrifuged at 3000 g for 15 min to separate serum after standing for 15 min. The levels of TC and TG in the serum were measured using a total cholesterol (TC) kit and a triglyceride (TG) kit (Nanjing Jian Cheng Biotechnology Company, Ltd., China), and various cholesterol indexes of the initial serum were recorded.

Groups C, D and E were fed daily with feed mixed with equal amounts (0.2 g/group) of Gutian red yeast powder, homemade red yeast powder and Xuezhikang drug respectively for 3 weeks. Groups A and B were fed with basal feed. After the last administration, each group of mice was fasted (without water fasting) for 12 h and then the levels of TC and TG in the serum were determined for each group with reference to the above method.

The experimental data in this study were all statistically analyzed using DPS data processing system.

Results

Separation and Screening Results of High-yield MK *Monascus* Strains

A total of 27 *Monascus* strains with good mycelial growth were isolated and purified from 7 commercially available samples of red yeast powder, and 13 strains were purchased from the Culture Collection. The solid fermentation products of 40 strains were pretreated, with their MK content determined by HPLC. The results showed that (Table S1): the MK yields of strains FG-2, APS 012, FG-1, FG-4, HB-3 and BG-2 were significantly higher than the model strain ACCC 30501. The strain with the highest MK yield was FG-2, reaching 2.46 mg/g, about 2.3 times that of the control strain ACCC 30501.

Molecular Identification Results of High-yield MK *Monascus* Strains

The ITS sequence amplification product of strain FG-2 showed a clear specific band between 500-600 bp (Fig. 1). The results of sequencing and alignment analysis showed that the strain has the highest ITS sequence similarity to the *Monascus purpureus* strain Mp-41, reaching 99% (Fig. S1). Therefore, strain FG-2 was identified as *Monascus purpureus*.

Effects of Different Rice Raw Materials on MK Yield by *Monascus* Fermentation

The solid fermentation test of strain FG-2 was carried out using 13 varieties of rice as raw materials. Each rice variety was divided into polished and brown rice, with a total of 26

rice raw materials. With the commercially available high-quality rice "Dulimi" as the control CK1 and the commercially available Gutian red yeast rice as the control CK2, comparison was made on the effects of different rice raw materials on MK yield. The results show that (Table S2): rice raw materials in different varieties have significant effects on MK yield of *Monascus*. Glutinous rice was not suitable as rice raw material for *Monascus* fermentation. MK yield was higher when polished rice was used as the raw material compared to brown rice of the same variety. When the rice variety "1a825" (polished rice) was used as raw material, the MK yield was the highest, reaching 3.61 mg/g, about 2.7 times that of "Dulimi" and 6.5 times that of the commercially available Gutian red yeast rice. Therefore, in the subsequent experiments of this study, the rice variety "1a825" (polished rice) was used as the raw material for *Monascus* fermentation.

Single Factor Optimization Results of Solid State Fermentation Conditions

Effect of rice amount in fermentation bottle on MK yield in fermentation products: The 20 g, 30 g, 40 g, 50 g, 60 g of rice steamed until zero white core were weighed and placed in a plastic fermentation bottle. Refer to 1.2.2 for other conditions. The results showed that (Fig. 2): the rice amount of fermentation bottle has a great influence on MK yield. When the rice amount is too small, the medium tends to lose moisture and the nutrients will dry quickly, unable to provide the nutrients necessary for normal *Monascus* growth; when the rice amount is too much, the oxygen in the fermentation bottle is insufficient, which is not conducive to *Monascus* growth and in turn affects MK formation. MK yield was the highest when the rice amount was 30 g, reaching 4.67 mg/g.

Effect of pre-fermentation temperature on MK yield in fermentation products: The red yeast fermentation bottles were placed in an incubator at 26°C, 28°C, 30°C and 32°C, respectively. On the 8th day, all were transferred to a 26°C incubator for further fermentation. Refer to 1.2.2 for the other conditions. The results showed that (Fig. 3): pre-fermentation temperature also has a great influence on MK yield. Temperature can directly affect the growth rate of mycelium. When the pre-fermentation temperature is too low, the mycelium grows slowly and the amount of fermentation products decreases. When the pre-fermentation temperature is too high, the mycelium is too exuberant with nutrients consumed in advance, which is also not beneficial to the accumulation of fermentation products in the later stage. Therefore, in the early stage of fermentation (1-7d), it is appropriate for mycelium to grow at a higher temperature; and the later (after 8d) low temperature environment is helpful to the formation of secondary metabolites. MK yield was the highest when the pre-fermentation temperature was 30°C, reaching 4.30 mg/g.




Fig. 1: Amplification of ITS sequence from *Monascus* strain FG-2. (M) DNA Marker DL 2000. (1) amplification of ITS from *Monascus* strain FG-2




Fig. 2: Effects of rice weight loaded in a fermentation flask on the production of MK. Different letters in different treatment indicate significant differences ($P < 0.05$), the same below




Fig. 3: Effects of temperature of fermentation prophase on the production of MK




Fig. 4: Effects of pH of seed liquid on the production of MK

Effect of seed solution pH on MK yield in fermentation products:

Five pH levels of the seed solution were selected: 4.0, 4.5, 5.0, 5.5 and 6.0. Refer to 1.2.2 for other conditions. The results showed (Fig. 4) that seed solution pH had a significant effect on MK yield. As *Monascus* is acidophilic and suitable for growth in an acidic environment, seed solution pH can be adjusted with acetic acid to increase MK yield. When seed solution pH is below 5.0, MK yield increases with pH increase; when the seed solution pH is over 5.0, MK yield decreases significantly with pH increase. MK yield was the highest when seed solution pH was 5.0, reaching 4.18 mg/g.

Effect of *Monascus* inoculum concentration on MK yield in fermentation products:

Five inoculum sizes were selected: 6%, 8%, 10%, 12%, 14% (v/m). Refer to 1.2.2 for other conditions. The results showed that (Fig. 5): the inoculum size of seed solution had a significant effect on MK yield. When the inoculum size is too small, with excessive culture medium nutrient, slowed mycelium growth and prolonged fermentation cycle, MK yield is low; when the inoculum size is excessive, with exuberant mycelium growth and insufficient supply of medium nutrient and oxygen, MK yield is limited. MK yield was the highest when the inoculum size was 12%, reaching 4.42 mg/g.

Effect of fermentation bottle permeability on the yield of MK in fermentation products:

The fermentation bottle permeability was adjusted by changing the number of air holes of the fermentation bottle. Five air hole levels of 1, 2, 3, 4 and 5 were selected. Refer to 1.2.2 for other conditions. The results showed that (Fig. 6): fermentation bottle permeability has a great influence on MK yield. MK yield increases with the gas permeability increase. *Monascus* is an aerobic microorganism, which has a high oxygen demand in the fermentation process. Better fermentation bottle permeability is favorable for the growth and fermentation of *Monascus* under the premise of no contamination by infectious microbe, leading to increased MK yield. Accordingly, MK yield of *Monascus* can be increased by appropriately increasing fermentation bottle permeability.

Effect of fermentation time on MK yield in fermentation products:

During the fermentation, samples were taken at 5 d, 8 d, 11 d, 14 d, 17 d and 20 d respectively to determine MK content. The results show that (Fig. 7): MK is not detected in the fermentation products at the initial fermentation stage (≤ 5 d), indicating that *Monascus* is still in the process of mycelium growth in this stage, and no MK is formed; in the fermentation metaphase (8-17 d), mycelium growth is basically completed, MK gradually forms; when fermented to 17 d, MK yield is the highest; when fermented to 20 d, MK content decreases. During the fermentation of *Monascus* strain FG-2, MK yield reached its maximum at 17 d.

Effect of additional carbon source types on MK yield in fermentation products:

Glucose, sucrose, α -lactose and soluble starch were added to the rice fermentation medium as additional carbon sources. The addition amount




Fig. 5: Effects of inoculum concentration of *Monascus* on the production of MK




Fig. 6: Effects of ventilation property of fermentation flask on the production of MK




Fig. 7: Effects of fermentation time on the production of MK




Fig. 8: Effects of additional carbon sources on the production of MK

was 3% (w/w) and that without additional carbon source was taken as the control group. Refer to 1.2. 2 for other conditions. The results show that (Fig. 8): addition of soluble

starch and glucose can significantly increase MK yield of the fermentation product, while the addition of sucrose and α -lactase has no significant effect on MK yield. Soluble starch is more expensive than glucose, so glucose can be selected as an additional carbon source to increase MK yield in consideration of the production cost.

Effect of additional nitrogen source types on MK yield in fermentation products: Yeast extract, peptone, sodium nitrate and ammonium sulfate were added to the rice fermentation medium as additional nitrogen sources. The addition amount was 4% (w/w) and that without additional nitrogen source was taken as the control group. Refer to 1.2.2 for other conditions. The results show that (Fig. 9): addition of sodium nitrate can greatly increase MK yield, followed by the effect of peptone and ammonium sulfate, and yeast has least effect. Therefore, sodium nitrate can be selected as an additional nitrogen source to increase MK yield.

Effect of rice soaking time on MK yield in fermentation products: Five rice soaking time was selected: 0h, 3h, 6h, 9h and 12h. The rice was drained after soaking. Refer to 1.2.2 for other conditions. The results show that (Fig. 10): MK yield gradually decreases with the increase of rice soaking time. It is possible that the rice variety "1a825" is suitable for direct cooking without soaking. Hence, the rice is directly cooked after cleaning, which is more favorable for MK production in *Monascus* fermentation.

Effect of Additional Precursor Substance on MK Yield in Fermentation Products: It can be known from the synthetic pathway of MK that, acetic acid is a precursor substance for MK synthesis, but acetic acid is volatile and has some toxicity to the thallus growth. Therefore, sodium acetate is usually used instead of acetic acid as a precursor substance of MK (Mulder *et al.*, 2015).



The 1% (w/w) sodium acetate was added to the rice fermentation medium and that without sodium acetate was taken as the control group. Refer to 1.2.2 for other conditions. The results show (Fig. 11) that addition of 1% sodium acetate can increase MK yield by 44.8%.

Orthogonal Optimization Results of Solid State Fermentation Conditions

According to the above single factor optimization experiment results, three levels of the four factors were selected for orthogonal optimization of solid state fermentation conditions of strain FG-2: pre-fermentation temperature (28°C, 30°C, 32°C), seed solution pH (4.5, 5.0, 5.5), rice amount of fermentation bottle (20 g, 30 g, 40 g) and fermentation time (14 d, 17 d, 20 d). The results show that (Table 1), the primary and secondary factors affecting MK yield are in the order of: B>A>D>C, *i.e.*: initial pH > temperature > fermentation time > rice amount. Further analysis of variance and significance test showed that factors A (temperature) and B (pH) had extremely significant effects on MK yield ($p < 0.01$). By comparing the magnitude of K, the optimal levels of the four factors

Table 1: The results and analysis of L_9 (3^4) orthogonal test

test number	factors				MK yield (mg/g)
	A (temperature of fermentation prophase / °C)	B (pH of seed liquid)	C (weigh of rice / g)	D (fermentation time / d)	
1	28	4.5	20	14	3.062
2	28	5.0	30	17	4.375
3	28	5.5	40	20	3.584
4	30	4.5	30	20	3.719
5	30	5.0	40	14	4.510
6	30	5.5	20	17	3.825
7	32	4.5	40	17	3.277
8	32	5.0	20	20	3.616
9	32	5.5	30	14	3.028
K1	3.674	3.353	3.501	3.533	
K2	4.018	4.167	3.707	3.826	
K3	3.307	3.479	3.790	3.640	
R	0.711	0.814	0.289	0.293	


Fig. 9: Effects of additional nitrogen sources on the production of MK

Fig. 10: Effects of rice soaking time on the production of MK

were determined as A2, B2, C3 and D2, that is, pre-fermentation temperature 30°C, seed solution pH 5.0, rice amount in the fermentation bottle 40 g and fermentation time 17 d. The MK yield under the fermentation conditions was 4.64 mg/g (Note: the inoculum size was 10%, the number of air holes was adjusted to 5 wells, no additional carbon source, nitrogen source and sodium acetate were added), which was 28.5% higher than before optimization, and about 8.3 times that of commercially available Gutian red yeast rice.

Study on Cholesterol-lowering Effect of High MK Red Yeast Rice

The results show that (Fig. 12 and 13): before administration, the TC values of the serum of B, C, D, E groups are 42.6% higher than that of A group ($P < 0.05$); the TG value is about 38.5% higher than that of A group ($P < 0.01$). This indicates that the experimental hyperlipidemia model of KM mice is successfully modeled. After three weeks of administration, the TC values of C, D and E groups were 29.5%, 34.1% and 37.0% lower than those of group B ($P < 0.01$); The TG values were 3.7%, 4.9% and 7.9% lower than those of group B, respectively, without significant difference. This shows that high MK red yeast rice prepared herein and the commercially available Gutian red yeast rice can significantly reduce the total cholesterol level in mice.

Discussions

Studies had shown that different substrates, such as rice, millet, wheat and barley, sorghum, cassava, sweet potato and potato, have an important influence on MK yield of *Monascus* fermentation (Lee *et al.*, 2006; Subhagar *et al.*, 2009; Venkateswaran and Vijayalakshmi, 2010; Zhang *et al.*, 2018). Priatni *et al.* (2014) used tofu, rice and *Dioscorea hispida* respectively as the substrate to carry out solid fermentation of *Monascus*. The results showed that MK yield was the highest in rice 4.6 times of *Dioscorea hispida* and 4.3 times of tofu. Panda *et al.* (2010) found that the predicted MK yield was 2.83 mg/g and the actual MK yield was 2.80 mg/g when rice used as the substrate, indicating that the rice-based fermentation efficiency was as high as 98.93%. Therefore, rice may be an excellent substrate for the fermentation of *Monascus*.

Pengnoi *et al.* (2017) used different varieties of rice as the matrix for *Monascus* fermentation. The results showed that MK yield (13.48 mg/kg) of rice variety Doi Muser as a matrix was significantly higher than other varieties of rice. This indicated that different rice varieties also had a significant effect on MK yield in *Monascus* fermentation.




Fig. 11: Effects of additional precursors on the production of MK




Fig. 12: TC levels in the serum of mice in each test group before administration and three weeks after administration. * indicates significant difference between B, C, D, E groups and A group before the administration ($P < 0.05$); ** indicates extremely significant difference between B group and A group after administration ($P < 0.01$)




Fig. 13: TG levels in the serum of mice in each test group before administration and three weeks after administration. ** indicates extremely significant difference between B, C, D, E groups and A group before the administration ($P < 0.01$)

In addition, the viscosity of rice has a great influence on MK yield. Because of its relatively high viscosity and easy agglomeration, glutinous rice is not conducive to *Monascus* growth (Pattanagul et al., 2007). This study also showed that MK yield of the fermentation product is extremely low, even without detection when the rice (all glutinous rice) numbered

"1a816", "1a814", "n84", "axn", "1A814", "1A816", "N84", "AXN" are used as the raw material (Table S2). This suggests that the glutinous rice is not suitable as a substrate for *Monascus* fermentation. It is possible that glutinous rice is bound into clusters after being cooked, the poor gas permeability hinders the mycelial growth of *Monascus*, thereby further reducing MK yield and even producing no MK. In addition, it can be seen from Table S2 that MK yield of polished rice as raw material is higher than brown rice in the same variety. It is possible that the brown rice has a skin layer and an aleurone layer on the surface, making *Monascus* mycelium unable to fully utilize the rice nutrients, and consequently leading to lower MK yield. In summary, selection of polished rice of non-glutinous rice varieties as a raw material for solid fermentation of *Monascus* can help increase MK yield of red yeast rice.

The solid-state fermentation condition for lovastatin produced by *A. terreus* is similar to MK produced by *Monascus* (Kumar et al., 2000). The carbon nitrogen ratio of the matrix is also one main factor affecting the yield of lovastatin by *A. terreus*. The yeast extract and soybean meal are the preferred nitrogen sources. The carbon in the lovastatin synthesis pathway is much lower than the carbon converted to biomass. Therefore, nitrogen limitation (i.e., growth inhibition) helps divert more carbon to the synthesis of lovastatin (Lopez et al., 2003).

Miranda et al. (2014) showed that reactive oxygen species (ROS) would be produced during the fermentation of *A. terreus* and the ROS concentration was positively correlated with the lovastatin yield. ROS could increase the expression of *Lov E* and *Lov F* in the lovastatin biosynthetic gene cluster, thereby increasing lovastatin yield. When antioxidants were added, significant reduction was observed in expression levels of lovastatin biosynthetic gene clusters and lovastatin yield. This indicates that oxidative stress can promote the synthesis of lovastatin or MK, and reactive oxygen species can significantly increase the content of lovastatin or MK in the fermentation product.

Conclusions

In this study, the optimal combination of MK high-yield strains and rice raw materials was screened out: *Monascus purpureus* strain FG-2 and polished rice of rice variety "1a825" and MK yield was 3.2 times of the control strain and 2.7 times of the control rice raw material, respectively. The optimum fermentation conditions for the combination were: pre-fermentation temperature 30°C, seed solution pH 5.0, rice amount in fermentation bottle 40 g, fermentation time 17 d; the MK yield was 4.64 mg/g under the fermentation condition, which was 28.5% higher than before fermentation condition optimization, and about 8.3 times of commercially available Gutian red yeast rice. The high MK red yeast rice prepared herein has better cholesterol lowering effect than the commercially available Gutian red yeast rice.

Acknowledgments

This study was supported financially by the International Science and Technology Cooperation and Exchange Project of Fujian Agriculture and Forestry University (KXGH17001), the National Key Research and Development Program (2017YFD0100100), the Science and Technology Project Guidance Project of Fujian Province (2015N0001) and the Special Fund for Science and Technology Innovation of Fujian Agriculture and Forestry University (CXZX2017413) in China.

References

- Alberts, A.W., J. Chen, G. Kuron, V. Hunt, J. Huff, C. Hoffman, J. Rothrock, M. Lopez, H. Joshua, E. Harris, A. Patchett, R. Monaghan, S. Currie, E. Stapley, G. Albers-Schonberg, O. Hensens, J. Hirshfield, K. Hoogsteen, J. Liesch and A.J. Springer, 1980. Mevinolin: a highly-potent competitive inhibitor of hydroxyl methylglutaryl- coenzyme A reductase and a cholesterol-lowering agent. *Proc. Natl. Acad. Sci. USA*, 77: 3957–3961
- Awad, E., E.M. Othman and H. Stopper, 2017. Effects of Resveratrol, Lovastatin and the mTOR-Inhibitor RAD-001 on insulin-induced genomic damage *in vitro*. *Molecules*, 22: 2207
- Barrios-Gonzalez, J., J.G. Banos, A.A. Covarrubias and A. Garay-Arroyo, 2008. Lovastatin biosynthetic genes of *Aspergillus terreus* are expressed differentially in solid-state and in liquid submerged fermentation. *Appl. Microbiol. Biotechnol.*, 79: 179–186
- Calvo, A.M., R.A. Wilson, J.W. Bok and N.P. Keller, 2002. Relationship between secondary metabolism and fungal development. *Microbiol. Mol. Biol. Res.*, 66: 447–459
- Chen, Y.P., C.P. Tseng, L.L. Liaw, C.L. Wang, I.C. Chen, W.J. Wu, M.D. Wu and G.F. Yuan, 2008. Cloning and characterization of monacolin K biosynthetic gene cluster from *Monascus pilosus*. *J. Agric. Food Chem.*, 56: 5639–5646
- Chen, Y.P., G.F. Yuan, S.Y. Hsieh, Y.S. Lin, W.Y. Wang, L.L. Liaw and C.P. Tseng, 2010. Identification of the mok H gene encoding transcription factor for the upregulation of monacolin k biosynthesis in *Monascus pilosus*. *J. Agric. Food Chem.*, 58: 287–293
- Dhale, M.A., S. Divakar, S.U. Kumar and G. Vijayalakshmi, 2007. Isolation and characterization of dihydromonacolin-MV from *Monascus purpureus* for antioxidant properties. *Appl. Microbiol. Biot.*, 73: 1197–1202
- Endo, A., 1979. Monacolin K, a new hypocholesterolemic agent produced by a *Monascus* species. *J. Antibiototechnol. (Tokyo)*, 32: 852–854
- Erdoğan, Ö. and S. Azirak, 2004. Review of the studies on the red yeast rice (*Monascus purpureus*). *Turk. Electron. J. Biotechnol.*, 2: 37–49
- Ghayour, M.B., A. Abdolmaleki and M.B. Rassouli, 2017. Neuroprotective effect of Lovastatin on motor deficit induced by sciatic nerve crush in the rat. *Eur. J. Pharmacol.*, 812: 121–127
- Guo, C.J. and C.C.C. Wang, 2014. Recent advances in genome mining of secondary metabolites in *Aspergillus terreus*. *Front. Microbiol.*, 5: 717
- Hasan, H., M.H. Abd Rahim, L. Campbell, D. Carter, A. Abbas and A. Montoya, 2018. Overexpression of acetyl-CoA carboxylase in *Aspergillus terreus* to increase lovastatin production. *New Biotechnol.*, 44: 64–71
- Hong, M.Y., N.P. Seeram, Y. Zhang and D. Heber, 2008. Anticancer effects of Chinese red yeast rice versus monacolin K alone on colon cancer cells. *J. Nutr. Biochem.*, 19: 448–458
- Huang, Z.B., Y. Xu, Y.P. Li and Y.H. Wang, 2010. Conversion Investigation for Lovastatin and Its Derivatives By HPLC. *J. Chromatogr. Sci.*, 48: 631–636
- II Gum, S., P.A. Nguyen, J.R. Lee, Y.H. Han and M.K. Cho, 2017. The physico-chemical alteration of lovastatin and enhanced antioxidant effect of *Bacillus subtilis* fermented-red yeast rice product. *Food Chem.*, 232: 203–209
- Kennedy, J., K. Auclair, S.G. Kendrew, C. Park, J.C. Vederas and C.R. Hutchinson, 1999. Modulation of polyketide synthase activity by accessory proteins during lovastatin biosynthesis. *Science*, 284: 1368–1372
- Kumar, M.S., S.K. Jana, V. Senthil, V. Shashanka, S.V. Kumar and A.K. Sadhukhan, 2000. Repeated fed-batch process for improving lovastatin production. *Process Biochem.*, 36: 363–368
- Lee, C.I., C.D. Shih, C.L. Lee, S.L. Kuo, S.C. Jheng and J.J. Wang, 2013. Effect of red mold rice on blood coagulation and anticoagulation factors in a rat model of hyperlipidemia. *J. Funct. Foods*, 5: 1956–1965
- Lee, C.L., J.J. Wang, S.L. Kuo and T.M. Pan, 2006. *Monascus* fermentation of dioscorea for increasing the production of cholesterol-lowering agent - monacolin K and antiinflammation agent - monascin. *Appl. Microbiol. Biot.*, 72: 1254–1262
- Li, L., Y.C. Shao, Q. Li, S. Yang and F.S. Chen, 2010. Identification of Mga1, a G-protein α -subunit gene involved in regulating citrinin and pigment production in *Monascus ruber* M7. *Fems. Microbiol. Lett.*, 308: 108–114
- Lin, L., C.L. Wang, Z.J. Li, H.J. Wu and M.H. Chen, 2017. Effect of temperature-shift and temperature-constant cultivation on the monacolin k biosynthetic gene cluster expression in *Monascus* sp. *Food Technol. Biotech.*, 55: 40–47
- Lin, L., S.F. Wu, Z.J. Li, Z.Y. Ren, M.H. Chen and C.L. Wang, 2018. High expression level of mok E enhances the production of monacolin K in *Monascus*. *Food Biotechnol.*, 32: 35–46
- Liu, Y.Q., X.H. Tu, Q. Xu, C.X. Bai, C.X. Kong, Q. Liu, J.H. Yu, Q.Q. Peng, X.S. Zhou, Y.X. Zhang and M.H. Cai, 2018. Engineered monoculture and co-culture of methylotrophic yeast for de novo production of monacolin J and lovastatin from methanol. *Metab. Eng.*, 45: 189–199
- Lopez, J.L.C., J.A.S. Perez, J.M.F. Sevilla, F.G.A. Fernandez, E.M. Grima and Y. Chisti, 2003. Production of lovastatin by *Aspergillus terreus*: effects of the C: N ratio and the principal nutrients on growth and metabolite production. *Enzyme Microb. Tech.*, 33: 270–277
- Miranda, R.U., L.E. Gomez-Quiroz, M. Mendoza, A. Perez-Sanchez, F. Fierro and J. Barrios-Gonzalez, 2014. Reactive oxygen species regulate lovastatin biosynthesis in *Aspergillus terreus* during submerged and solid-state fermentations. *Fungal Biol-UK*, 118: 979–989
- Mulder, K.C.L., F. Mulinari, O.L. Franco, M.S.F. Soares, B.S. Magalhaes and N.S. Parachin, 2015. Lovastatin production: From molecular basis to industrial process optimization. *Biotechnol. Adv.*, 33: 648–665
- Panda, B.P., S. Javed and M. Ali, 2010. Optimization of fermentation parameters for higher lovastatin production in red mold rice through co-culture of *Monascus purpureus* and *Monascus ruber*. *Food Bioprocess Tech.*, 3: 373–378
- Pattanagul, P., R. Pinthong and A. Phianmongkhon, 2007. Review of Angkak production (*Monascus purpureus*). *Chiang Mai J. Sci.*, 34: 319–328
- Pengnoi, P., R. Mahawan, C. Khanongnuch and S. Lumyong, 2017. Antioxidant properties and production of monacolin K, citrinin, and red pigments during solid state fermentation of purple rice (*Oryza sativa*) varieties by *Monascus purpureus*. *Czech J. Food Sci.*, 35: 32–39
- Priatni, S., S. Damayanti, V. Saraswati, D. Ratnaningrum and M. Singgih, 2014. The utilization of solid substrates on *Monascus* fermentation for anticholesterol agent production. *Proc. Chem.*, 9: 34–39
- Ruscica, M., M. Gomaschi, G. Mombelli, C. Macchi, R. Bosisio, F. Pazzucconi, C. Pavane, L. Calabresi, A. Arnoldi, C.R. Sirtori and P. Magni, 2014. Nutraceutical approach to moderate cardiometabolic risk: Results of a randomized, double-blind and crossover study with Armolipid Plus. *J. Clin. Lipidol.*, 8: 61–68
- Sakai, K., H. Kinoshita and T. Nihira, 2009. Identification of mok B involved in monacolin K biosynthesis in *Monascus pilosus*. *Biotechnol. Lett.*, 31: 1911–1916
- Savitha, J., S.D. Bhargavi and V.K. Praveen, 2016. Complete genome sequence of soil fungus *aspergillus terreus* (km017963), a potent lovastatin producer. *Genom. Announ.*, 4: e00491–16
- Shannon, J., S. Tewoderos, M. Garzotto, T.M. Beer, R. Derenick, A. Palma and P.E. Farris, 2005. Statins and prostate cancer risk: A case-control study. *Am. J. Epidemiol.*, 162: 318–325

- Singgih, M., V. Saraswaty, D. Ratnaningrum, S. Priatni and S. Damayanti, 2014. The influence of temperature and ethanol concentration in monacolin k extraction from *Monascus* fermented rice. *Proc. Chem.*, 9: 242–247
- Subhagar, S., R. Aravindan and T. Viruthagiri, 2009. Response surface optimization of mixed substrate solid state fermentation for the production of lovastatin by *Monascus purpureus*. *Eng. Life Sci.*, 9: 303–310
- Venkateswaran, V. and G. Vijayalakshmi, 2010. Finger millet (*Eleusine coracana*) - an economically viable source for antihypercholesterolemic metabolites production by *Monascus purpureus*. *J. Food Sci. Tech.*, 47: 426–431
- Wei, Y. and D.G. Popovich, 2013. Red azaphilone pigments extracted from red yeast rice induces cellular senescence and reduces viability in HepG2 cells. *Biomed. Prevent. Nutr.*, 3: 331–337
- Yin, Y., M.H. Cai, X.S. Zhou, Z.Y. Li and Y.X. Zhang, 2016. Polyketides in *Aspergillus terreus*: biosynthesis pathway discovery and application. *Appl. Microbiol. Biotech.*, 100: 7787–7798
- Zhang, B.B., H.B. Xing, B.J. Jiang, L. Chen, G.R. Xu, Y. Jiang and D.Y. Zhang, 2018. Using millet as substrate for efficient production of monacolin K by solid-state fermentation of *Monascus ruber*. *J. Biosci. Bioeng.*, 125: 333–338
- Zhang, B.B., L.P. Lu and G.R. Xu, 2015. Why solid-state fermentation is more advantageous over submerged fermentation for converting high concentration of glycerol into Monacolin K by *Monascus purpureus* 9901: A mechanistic study. *J. Biotechnol.*, 206: 60–65
- Zhang, B.B., L.P. Lu, Y.J. Xia, Y.L. Wang and G.R. Xu, 2013. Use of agar as carrier in solid-state fermentation for Monacolin K production by *Monascus*: A novel method for direct determination of biomass and accurate comparison with submerged fermentation. *Biochem. Eng. J.*, 80: 10–13
- Zhang, C., J. Liang, L. Yang, B.G. Sun and C.T. Wang, 2017. De Novo RNA Sequencing and Transcriptome Analysis of *Monascus purpureus* and Analysis of Key Genes Involved in Monacolin K Biosynthesis. *PLoS One*, 12: e0170149
- Zhou, Y.J., H. Yang, X.D. Zhou, H.K. Luo, F. Tang, J. Yang, G. Alterovitz, L. Cheng and B. Ren, 2018. Lovastatin synergizes with itraconazole against planktonic cells and biofilms of *Candida albicans* through the regulation on ergosterol biosynthesis pathway. *Appl. Microbiol. Biotech.*, 102: 5255–5264
- Zhu, L., L.F. Yau, J.G. Lu, G.Y. Zhu, J.R. Wang, Q.B. Han, W.L. Hsiao and Z.H. Jiang, 2012. Cytotoxic dehydromonacolins from red yeast rice. *J. Agric. Food Chem.*, 60: 934–939
- Zlova, P.J., L. Martnkova and V.K. En, 1996. Secondary metabolites of the fungus *Monascus*: A review. *J. Ind. Microbiol.*, 16: 163–170

(Received 18 December 2018; Accepted 28 January 2019)

bioRxiv preprint doi: <https://doi.org/10.1101/2019.01.28.179111>; this version posted January 29, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.