



### Full Length Article

## Calorific Value of Cereals and Cereal by-products in Growing Male Broilers

Zafar Ullah<sup>1\*</sup>, Muhammad Sarwar<sup>1</sup>, Gulraiz Ahmed<sup>2</sup>, Javed Iqbal Sultan<sup>1</sup> and Ahsan ul Haq<sup>1</sup>

<sup>1</sup>Institute of Animal Sciences, University of Agriculture, Faisalabad, Pakistan

<sup>2</sup>Sadiq Feeds (Pvt.) Ltd. 48-C, Satellite Town, Chandni Chowk, Rawalpindi, Pakistan

\*For correspondence: zu2005@yahoo.com

### Abstract

An experiment was conducted to determine apparent metabolizable energy corrected for nitrogen (AMEn) of commonly used cereals and cereal by-products in poultry diet. The 3 samples of each of corn, rice broken (RB), rice polishings (RP) and wheat bran (WB) were assayed for AMEn by substitution method using 21-days-old broilers. The 12 experimental diets (4 ingredients x 3 samples) were formulated by substituting the corn-soybean meal basal diet with 30% test ingredients. The acid insoluble ash was added at 1% of all diets. The 468 day-old male broiler chicks (Hubbard x Hubbard) were reared on basal diet from day 1 to 13 and thereafter experimental diets were offered *ad libitum* to birds from 14 to 21 days. Each diet was offered to 36 birds, equally distributed to six replicate pens. The basal diet continued to feed the birds of 6 replicates from day 14 to 21. The excreta samples were collected for 3 days, day 19-21 and AMEn was calculated. The average daily feed intake (ADFI), average daily gain (ADG) and feed conversion ratio (FCR) were calculated from 14-21 days of age. Data regarding ADFI, ADG, FCR and AMEn were analyzed using GLM procedures of SAS while means were partitioned by t-test. The ADFI, ADG, FCR and AMEn remained unaltered ( $P>0.05$ ) by feeding diets based on samples of same feed ingredient except the RP and WB. The birds fed RP and WB based diets showed varying ( $P<0.05$ ) AMEn between the samples. The birds fed corn substituted basal diet had higher ( $P<0.05$ ) ADG (56.03 g) followed by RB (50.04 g), WB (45.87 g) and RP (43.61 g) diets. The ADFI of broilers was highest ( $P<0.05$ ) fed corn diets while it was lowermost with RB based diets. The birds showed improved ( $P<0.05$ ) FCR by feeding RB diet followed by corn, WB and RP diets. The average AMEn of RB and corn was 3372 and 3315 kcal/kg, respectively in cereals. In cereal by-products, the RP and WB had 1936 and 1760 kcal/kg AMEn, respectively. In conclusion, varying growth pattern of broilers was observed by feeding corn, RB, RP and WB substituted diets. The RB and RP's AMEn was higher in cereals and cereal by-products, respectively. © 2016 Friends Science Publishers

**Keywords:** Broiler; Apparent metabolizable energy corrected for nitrogen; Growth performance; Cereals; Cereal by-products

### Introduction

The feed cost represents more than 70% share of total broiler production cost, out of which the major portion is used to supply required dietary metabolizable energy (Jiang, 2004; Donohue and Cunningham, 2009). Cereals and cereal by-products contribute a major energy share in poultry diets (Sibbald, 1982). These ingredients' inclusion level in diet is based on their available nutrients particular the metabolizable energy (ME). The ME intake influences the broiler's feed intake and growth performance (Leeson and Summers, 2001; Morris, 2004) that's why poultry diets are formulated to fulfill bird's ME requirement (Nadeem *et al.*, 2005). Precise and accurate assessment of feed ingredient's ME in is mandatory to prepare a balanced diet with a definite calorific concentration (Scott *et al.*, 1998; Nadeem *et al.*, 2005; Macleod *et al.*, 2008). The feed ingredient's nutrient composition influences its

calorific value (Toghyani *et al.*, 2014) and this nutrient composition is affected by varying plant genetics, soil conditions, agronomic practices and climatic changes (Ravindran *et al.*, 2014). The ME value differs not only between ingredients but also amid samples of same feed ingredient (Jiang, 2004) produced at different locations of the same country.

The term apparent ME (AME) is classically used to represent ME of an ingredient and is the difference in gross energy (GE) intake and energy voided through feces and urine. For comparative purposes, these AME values are corrected for N retention (AMEn) to convert all data on N equilibrium basis (Fisher, 2000; Leeson and Summers, 2001; Macleod, 2002). The N correction is generally made to determine the nutrients potential to provide energy not their ability to promote N retention (McNab, 2000). These AMEn values are more authentic and additive approach compared to AME values (Jiang, 2004).

The advanced countries have developed the database regarding AMEn values of feed ingredients produced in their respective regions. But, unfortunately, the same information of our locally available ingredients is limited. The partial available information was determined using adult cockerels or roosters (Nadeem *et al.*, 2005; Pasha *et al.*, 2008). But the ME values are influenced by age (Zelenka, 1968; Lodhi *et al.*, 1969), species (Slinger *et al.*, 1964; Fisher and Shannon, 1973) and strains (Foster, 1968; March and Biely, 1971) of birds. Even broiler classes varying in growth performance resulted in different AME values (Jorgensen *et al.*, 1990; Ten Doeschate *et al.*, 1993). Broilers showed varying, generally, greater AME values than cockerels (Bourdillon *et al.*, 1990). Therefore, the feed ingredient's AME determined in older aged cockerels or roosters may not be equally useful in formulating precise and cost-effective growing broiler's diet. Therefore, the present study was planned to determine AMEn values of commonly used cereals and cereal by-products in growing male broilers.

## Materials and Methods

### Feed Ingredients

The 2 cereals and 2 cereal by-products determined for AMEn content included corn (*Zea mays* L.), rice broken (*Oryza sativa* L.), rice polishings (*Oryza sativa* L.) and wheat bran (*Triticum spp.*). The 3 samples of each feed ingredient were analyzed for AMEn value. These feed ingredients were assayed for dry matter (DM), crude protein (N x 6.25) by LECO® nitrogen analyzer (model FP-528, Leco Corporation, St. Joseph, MI), crude fat, crude fiber and ash content (AOAC, 2000) (Table 1).

### Test Diets

The 12 experimental diets (4 ingredients x 3 samples) were formulated by substituting the corn-soybean meal basal diet (Table 2) with 30% test ingredient (Table 3). The Celite® (acid insoluble ash) was added at 1% of each diet (Dourado *et al.*, 2010). The vitamins and minerals supplementation was similar across all diets. The feed ingredients' AMEn was determined by substitution method according to Macleod *et al.* (2008) using 3-weeks old male broilers as used by Woyengo *et al.* (2010).

### Bird's Management

The 468 day-old male broiler chicks (Hubbard x Hubbard) were arranged from hatchery (SB Hatchery, Rawalpindi, Pakistan) and reared in cages. All chicks were raised under same managemental conditions. The room temperature was kept at 32±1°C during 1<sup>st</sup> week and gradually reduced to 24°C by the end of 3<sup>rd</sup> week. The birds were provided continuous fluorescent light during the experimental period.

Chicks were vaccinated against Newcastle disease (ND), Infectious Bronchitis (IB) and Infectious Bursal disease (IBD). The fresh water's *ad-libitum* supply was available to chicks round the clock. All chicks were offered *ad-libitum* corn-soybean basal diet in crumble form to fulfill their nutritional requirements (NRC, 1994) for first 13 days of age. On day 14<sup>th</sup>, chicks were fasted overnight, individually weighed and randomly distributed to 78 replicate pens, each containing 6 birds. The difference in mean body weight of chicks between replicates was ± 10 g. The each of the basal diet as well as experimental diets was fed *ad-libitum* to six randomly selected replicates from day 14 to 21.

### Excreta Collection

The excreta samples from each replicate were collected 4 times a day, for 3 days (day 19 to 21). Feed and feather particles were removed from dropping samples as much as possible. The collected dropping samples were labeled according to diet and replicate and were refrigerated. After collection period, frozen samples were thawed, pooled replicate wise, homogenized and dried in hot air oven at 55°C for 72 h. Dried samples were ground in laboratory rotor mill (model ZM-200, Retsch GmbH, Haan, Germany) to pass through 0.5 mm sieve and stored for further analyses.

### Chemical Analysis

The experimental and basal diets and excreta samples were analyzed for DM (AOAC, 2000), nitrogen content by LECO® nitrogen analyzer (model FP-528, Leco Corporation, St. Joseph, MI) and gross energy using C2000 IKA adiabatic oxygen bomb calorimeter (GMBIT and CO KGD-79219, IKA Works Inc., Wilmington, NC). The acid insoluble ash of diets and dropping samples was determined by procedure reported in AOAC (2000).

### Calculations

The average daily feed intake (ADFI), average daily gain (ADG) of broiler chicks was recorded from day 14 to 21 and feed conversion ratio (FCR) was calculated. The AMEn value of all the test ingredient samples was calculated by data of diets and excreta analyses using following equations:

$$AME_n = GE_{Diet} - \frac{GE_{Excreta} \times AIA_{Diet}}{AIA_{Excreta}} - 8.22 \times N_{Retained}$$

Where, AMEn (kcal/kg) = N-corrected apparent metabolizable energy

GE<sub>Diet</sub> and GE<sub>Excreta</sub> (kcal/kg) = Gross energy content of diet and excreta, respectively

AIA<sub>Diet</sub> and AIA<sub>Excreta</sub> (%) = Acid insoluble ash in diet and excreta, respectively

8.22 (kcal/kg) = Energy value of uric acid

N<sub>Retained</sub> (g/kg) = N Retained by the birds per kg of feed intake

The retained N content was calculated by the following equation:

$$N_{\text{Retained}} = N_{\text{Diet}} - \frac{N_{\text{excreta}} \times AIA_{\text{Diet}}}{AIA_{\text{Excreta}}}$$

Where,  $N_{\text{Diet}}$  and  $N_{\text{Excreta}}$  (%) = N content of diet and excreta, respectively

AMEn of test ingredients was determined by the following formula:

$$AMEn_{\text{Ingredient}} = AMEn_{\text{Basal}} + \left[ \frac{AMEn_{\text{test}} - AMEn_{\text{Basal}}}{\text{Proportion of test ingredient in test diet}} \right]$$

### Statistical Analysis

Data regarding ADFI, ADG, FCR and AMEn were analyzed using GLM procedures of SAS (SAS, 1999). In case of significance, treatment means were compared by t-test.

### Results

The nutrient composition of feed ingredients is given in Table 1. The ingredient composition of basal diet as well as experimental diets is mentioned in Table 2 and 3, respectively. The ADFI, ADG and FCR in broilers remained same ( $P > 0.05$ ) fed diets based on different samples of same feed ingredients (Table 4). The AMEn of corn and RB was similar ( $P > 0.05$ ) between different samples; however, this AMEn varied ( $P < 0.05$ ) between samples of cereal by-products (RP and WB) (Table 4). The ADFI, ADG, FCR and AMEn was different ( $P < 0.05$ ) between different feed ingredients (Table 5). The greater ( $P < 0.05$ ) ADG in broilers was recorded by feeding corn diet (56.03 g) followed by RB (50.04 g), WB (45.87 g) and RP (43.61 g) diets. The more ( $P < 0.05$ ) ADFI was recorded when basal diet was substituted with corn while it was lowest with RB (Table 5). The improved ( $P < 0.05$ ) FCR in broilers was observed by feeding RB (1.94) diets followed by corn (2.02), WB (2.33) and RP (2.50).

The variation in different ingredient's AMEn was significant. The more ( $P < 0.05$ ) AMEn was recorded for RB (3372 kcal/kg) followed by corn (3315 kcal/kg), RP (1936 kcal/kg) and WB (1760 kcal/kg).

### Discussion

The nutrient composition of feed ingredients was concordant with published values (Sauvant *et al.*, 2004; Nadeem *et al.*, 2005; Rostagno *et al.*, 2005). The more ( $P < 0.05$ ) ADG in broilers fed corn substituted diets might be attributed to increased feed intake. The reduced ADFI of RP and WB diets might be because of more dietary fiber content. In this experiment, the basal diet was 30% substituted with test ingredients so the replacement of ingredients with higher fiber content resulted in increased dietary fiber. Kras *et al.* (2013)

**Table 1:** Chemical composition of corn, rice broken, rice polishings and wheat bran used in determination of AMEn (as-received basis)

Nutrient (%)	Ingredients			
	Corn	Rice broken	Rice polishings	Wheat bran
Moisture	8.67 ± 0.44	9.87 ± 0.40	6.84 ± 0.62	8.20 ± 1.28
Crude protein	8.81 ± 0.54	10.07 ± 0.21	12.38 ± 0.90	13.50 ± 0.71
Ether extract	3.61 ± 0.07	1.21 ± 0.05	10.74 ± 1.44	3.33 ± 0.47
Crude fiber	1.95 ± 0.12	0.68 ± 0.02	17.38 ± 4.06	10.21 ± 1.63
Ash	1.25 ± 0.17	1.01 ± 0.14	11.55 ± 0.63	5.09 ± 0.49
Acid insoluble ash	0.08 ± 0.06	0.17 ± 0.04	7.92 ± 0.72	1.20 ± 0.43

‡Data were average of three samples of each feed ingredient

**Table 2:** Ingredient and chemical composition of basal diet

Ingredient	Percentage composition
Corn	58.00
Soybean meal	35.52
Soybean oil	1.78
Di-calcium phosphate	2.17
Limestone	0.78
Sodium chloride	0.20
Sodium bicarbonate	0.23
Vitamin-mineral premix <sup>‡</sup>	0.32
Celite <sup>†</sup>	1.00
Chemical composition (%)	
Moisture	8.80
Crude protein	22.93
Ash	7.97
Crude fiber	2.40
Crude fat	4.02

<sup>†</sup>Celite: Acid insoluble ash

<sup>‡</sup> Provided per kg of diet: Retinyl acetate, 4400 IU; cholecalciferol, 118 µg; DL- $\alpha$ -tocopheryl acetate, 12 IU; menadione sodium bisulphite, 2.40 mg; thiamine, 2.5 mg; riboflavin, 4.8 mg; niacin, 30 mg; D-pantothenic acid, 10 mg; pyridoxine, 5 mg; biotin, 130 µg; folic acid, 2.5 mg; cyanocobalamin, 19 µg; manganese, 85 mg (MnSO<sub>4</sub>·H<sub>2</sub>O); Iron, 80 mg (FeSO<sub>4</sub>·H<sub>2</sub>O); Zinc, 75 mg (ZnO); Copper, 6 mg (CuSO<sub>4</sub>·5H<sub>2</sub>O); Iodine, 1 mg (ethylene diamine dihydroiodide); Selenium, 130 µg (Na<sub>2</sub>SeO<sub>3</sub>)

documented poor performance and lower energy retention of birds fed high fibrous diet. The lowest ( $P < 0.05$ ) ADFI of RB's diets might be due to its more AMEn content. Leeson and Summers (2001) and Morris (2004) reported that dietary ME value influenced broiler's feed intake and growth performance. The birds showed improved ( $P < 0.05$ ) FCR fed RB substituted diet might be because of lesser feed intake than that of birds fed other diets.

The samples of same feed ingredient resulted in varying AMEn values (Table 4). This AMEn variation was greater ( $P < 0.05$ ) among RP and WB's samples. The reason of such varying AMEn values between samples of same feed ingredient might be the differences in grains' nutrient composition cultivated in different areas. All those aspects influencing ingredient's digestibility affect its metabolizability. Like digestibility, the AMEn also depends upon ingredient's chemical composition because the plant genetics, soil conditions, agronomic practices and cultivar techniques affect both the ingredient's chemical composition and its nutritive value (Alvarenga *et al.*, 2013; Ravindran *et al.*, 2014). These differences were more

**Table 3:** Composition of experimental diets based on basal diet and test feed ingredients

Diets	Corn %			RB %			RP %			WB %			
	Basal Diet %	sample 1	sample 2	sample 3	sample 1	sample 2	sample 3	sample 1	sample 2	sample 3	sample 1	sample 2	sample 3
Corn diet 1	70	30	-	-	-	-	-	-	-	-	-	-	-
Corn diet 2	70	-	30	-	-	-	-	-	-	-	-	-	-
Corn diet 3	70	-	-	30	-	-	-	-	-	-	-	-	-
RB diet1	70	-	-	-	30	-	-	-	-	-	-	-	-
RB diet2	70	-	-	-	-	30	-	-	-	-	-	-	-
RB diet3	70	-	-	-	-	-	30	-	-	-	-	-	-
RP diet1	70	-	-	-	-	-	-	30	-	-	-	-	-
RP diet2	70	-	-	-	-	-	-	-	30	-	-	-	-
RP diet3	70	-	-	-	-	-	-	-	-	30	-	-	-
WB diet1	70	-	-	-	-	-	-	-	-	-	30	-	-
WB diet2	70	-	-	-	-	-	-	-	-	-	-	30	-
WB diet3	70	-	-	-	-	-	-	-	-	-	-	-	30

RB, RP and WB are the abbreviations of rice broken, rice polishings and wheat bran, respectively

prominent in by-products indicating variation in processing techniques and processing equipment's quality. These results elaborate the usefulness of determining available nutritive values of native feed resources rather than to use literature values in practical feed formulation.

In cereals, the RB had higher ( $P>0.05$ ) AMEn than that of corn's AMEn (Table 5). The AMEn of corn in this experiment was higher than that reported by Longo *et al.* (2004), Sauvant *et al.* (2004) and Nadeem *et al.* (2005) and less than energy value documented by Palmer-Jones and Halliday (1971), NRC (1994) and Rostagno *et al.* (2005). The reason of varying calorific values might be attributed to varying grain's nutrient composition. Palmer-Jones and Halliday (1971) reported that 3391 kcal/kg corn's energy with 3% fiber was reduced to 2532 kcal/kg when fiber content increased to 11%. Vieira *et al.* (2007) reported that corn hybrid's AMEn ranged from 3405 to 4013 kcal/kg and was related it to its varying lipid content. The ingredient's oil content influences its AMEn value (Zhou *et al.*, 2010). Wang and Parsons (1998b) reported that positive relationship existed in corn's energy level and its ether extract content. Meloche *et al.* (2013) documented decreased AMEn value of distillers dried grains with solubles exhibiting lower oil content than those of more oil percentage. The lipid content reduced the gastric emptying by increasing the cholecystokinin secretion in duodenum. This hormone enhanced the digestive enzymes secretions from pancreas and improved the protein and carbohydrate digestion thus increased the energy value (Alvarenga *et al.*, 2013). The contrasting results regarding corn's AMEn in this study and Longo *et al.* (2004) might be attributed to varying bird's age. Longo *et al.* (2004) used 1 week old broiler chicks in their study and digestibility and metabolizability is generally less in early aged birds than that of growing age.

The RB's AMEn in this study was contrasting to that reported by Sauvant *et al.* (2004), Nadeem *et al.* (2005) and Rostagno *et al.* (2005). Although, Nadeem *et al.* (2005) used native RB to determine energy value but higher energy value in this study might be because of varying rice variety

**Table 4:** Broilers' performance fed experimental diet based on different samples of feed ingredients from day 14 to 21 days of age

Parameters <sup>‡</sup>	Diet 1	Diet 2	Diet 3	SEM	P-value
<b>Corn</b>					
ADG (g)	56.81	56.45	54.83	0.64	0.424
ADFI (g)	113.31	112.83	113.10	0.14	0.420
FCR	1.99	2.0	2.07	0.02	0.423
AMEn (kcal/kg)	3303	3355	3286	26.89	0.574
<b>Rice broken</b>					
ADG (g)	49.51	50.66	49.97	0.40	0.521
ADFI (g)	96.37	97.65	98.06	0.49	0.367
FCR	1.95	1.94	1.96	0.01	0.779
AMEn (kcal/kg)	3343	3401	3376	23.18	0.680
<b>Rice polishings</b>					
ADG (g)	43.05	43.79	44.0	0.64	0.835
ADFI (g)	107.25	109.82	108.48	0.66	0.303
FCR	2.49	2.53	2.46	0.04	0.784
AMEn (kcal/kg)	2083 <sup>a</sup>	1752 <sup>b</sup>	1975 <sup>a</sup>	41.22	<0.001
<b>Wheat bran</b>					
ADG (g)	45.89	46.84	44.87	0.51	0.301
ADFI (g)	106.93	108.56	104.83	0.91	0.255
FCR	2.33	2.32	2.34	0.03	0.960
AMEn (kcal/kg)	1732 <sup>ab</sup>	1872 <sup>a</sup>	1678 <sup>b</sup>	29.35	0.010

Values sharing different superscripts (a-b) within rows differed significantly ( $P<0.05$ )

<sup>‡</sup>ADG = average daily gain/bird, ADFI = average daily feed intake/bird,

FCR = Feed conversion ratio

AMEn = Apparent metabolizable energy corrected for nitrogen

SEM = Standard error of mean

and experimental birds. The authors used 3-weeks old broilers in contrast to mature cockerels used by Nadeem *et al.* (2005). Bourdillon *et al.* (1990) documented greater energy values in broilers than cockerels and might be because of different metabolic rate of adult cockerels and growing broilers.

In this study, the RP possessed more ( $P<0.05$ ) AMEn than WB in cereal by-products. The RP's AMEn was less than that stated by Sauvant *et al.* (2004), Leeson and Summers (2001) and NRC (1994). The reason for such contrasting results might be due to varying processing conditions (Rao and Reddy, 1986). The RP is the by-product of rice processing industry thus processing affects

**Table 5:** Broilers' performance fed experimental diet based on different feed ingredients from day 14 to 21 days of age

Parameters‡	Diets based on feed ingredients				SEM	P- value
	Corn	Rice broken	Rice polishings	Wheat bran		
ADG (g)	56.03 <sup>a</sup>	50.04 <sup>b</sup>	43.61 <sup>d</sup>	45.87 <sup>c</sup>	0.62	< 0.001
ADFI (g)	113.08 <sup>a</sup>	97.35 <sup>c</sup>	108.52 <sup>b</sup>	106.77 <sup>b</sup>	0.74	<0.001
FCR	2.02 <sup>b</sup>	1.94 <sup>a</sup>	2.50 <sup>d</sup>	2.33 <sup>c</sup>	0.03	<0.001
AMEn (kcal/kg)	3315 <sup>a</sup>	3372 <sup>a</sup>	1936 <sup>b</sup>	1760 <sup>c</sup>	90.31	<0.001

Values sharing different superscripts (a-d) within rows differed significantly ( $P < 0.05$ )

‡ADG = average daily gain/bird, ADFI = average daily feed intake/bird, FCR = Feed conversion ratio

AMEn = Apparent metabolizable energy corrected for nitrogen

SEM = Standard error of mean

the chemical composition. The polishings and pressure during processing affects grain's quality (Ambreen *et al.*, 2006). Even varying temperature during processing can impact the energy level. The increasing drying temperature from 80 to 120°C resulted in decreased AMEn of corn without changing grain's gross energy and nutrient composition (Carvalho *et al.*, 2004). In addition, RP adulterated with rice husk (high in fiber) may be another aspect affecting its nutritional value (Ambreen *et al.*, 2006). It might also be due to inconsistent chemical composition of rice varieties (Rao and Reddy, 1986). That's why the AMEn of RP (1936 kcal/kg) was close to 1953 kcal/kg documented by Pasha *et al.* (2008) as they also used native RP samples in their experiment.

The determined WB's AMEn content was in-between the values stated by other workers (Kuzmicky *et al.*, 1978; Sauvant *et al.*, 2004; Nadeem *et al.*, 2005; Rostagno *et al.*, 2005). This difference in energy value might be attributed to variance in wheat type and soil or agronomic conditions. However, the AMEn of one WB sample (Table 4) was concordant with that reported by Nadeem *et al.* (2005). Cozannet *et al.* (2010a) reported that from proximate fractions, the only fiber content accounted for considerable AMEn variation in wheat by-products. Ravindran *et al.* (2014) reported that fiber was negatively correlated ( $r = -0.64$ ), while fat was positively correlated ( $r = 0.38$ ) to AME. Although an ample data regarding feed ingredient's energy value have been published. But several terminologies and methods had been used to determine these values. These methods included total collection or partial collection method with help of some appropriate marker. Different terms used included the apparent metabolizable energy (AME), apparent metabolizable energy corrected for nitrogen (AMEn), true metabolizable energy (TME) and true metabolizable energy corrected for nitrogen (TMEn). Furthermore, different animal species and strains of varying ages and sex were used, therefore, direct comparison of energy values determined in this trial and reported literature seems little valid.

In this trial, the higher AMEn value of RB from cereals and cereal by-products could be attributed to its more starch content. The cereals and their by-products being the rich sources of starches share equitable calories in poultry diet. The starch is the major energy source in grains (Moran, 1982; NRC, 1994) and its content may vary

between ingredients. There is less correlation between feed ingredient's energy value and starch content but there is very strong correlation between ingredient's calorific value and the digestible starch. The starch digestibility may vary not only between ingredients but also between samples of same ingredient. Rostagno *et al.* (2005) reported higher starch content in RB (74.45%) compared to corn (62.48), WB (54.93) and RP (27.40%) (Sauvant *et al.*, 2004).

## Conclusion

The varying nutrient composition of native feed ingredients and their by-products resulted in variations in broiler's growth performance and AMEn values. The RB had higher AMEn value amid cereals. Form cereal by-products, the RP exhibited more AMEn than WB. Formulating broiler diets using AMEn values determined from locally available feed ingredients may help in efficient broiler production.

## Acknowledgements

The study was supported by Higher Education Commission of Pakistan. The authors are grateful to Sadiq Feeds (Pvt.) Ltd. Rawalpindi, Pakistan, for providing chicks, feeds and laboratory facilities.

## Reference

- Alvarenga, R.R., M.G. Zangeronimo, P.B. Rodrigues, L.J. Pereira, R.C. Wolp and E.C. Almeida, 2013. Formulation of diets for poultry: The importance of prediction equations to estimate the energy value. *Arch. Zootec.*, 62: 1-11
- Ambreen, N., N.Q. Hanif and S. Khatoon, 2006. Chemical composition of rice polishing from different sources. *Pak. Vet. J.*, 26: 190-192
- AOAC, 2000. *Official Methods of Analysis*, 17<sup>th</sup> edition. Association of Official Analytical Chemists, Gaithersburg, MD, USA
- Bourdillon, A., B. Carre, L. Conan, M. Francesch, M. Fuentes, G. Huyghebaert, W.M.M.A. Jansen, B. Leclecq, M. Lessire, J. McNab, M. Rigoni and J. Wiseman, 1990. European reference method of *in vivo* determination of metabolizable energy in poultry: reproducibility, effect of age, comparison with predicted values. *Brit. Poult. Sci.*, 31: 567-576
- Carvalho, D.C., L.F.T. Albino, H.S. Rostagno, J.E. Oliveira, J.G. Vargas Jr., R.S. Toledo, C.H.R. Costa, S.R.F. Pinheiro and R.M. Souza, 2004. Chemical and energy composition of corn sample under different drying temperature and storage periods. *Braz. J. Vet. Anim. Sci.*, 33: 358-364
- Cozannet, P., M. Lessire, C. Gady, J.P. Metayer, Y. Primot, F. Skiba and J. Noblet, 2010a. Energy value of wheat dried distillers grains with solubles in roosters, broilers, layers, and turkeys. *Poult. Sci.* 89: 2230-2241

- Donohue, M. and D.L. Cunningham, 2009. Effects of grain and oilseed prices on the costs of US poultry production. *J. Appl. Poult. Res.*, 18: 325-337
- Dourado, L.R.B., J.C. Siqueira, N.K. Sakomura, S.R.F. Pinheiro, S.M. Marcato, J.B.K. Fernandes and J.H.V. Silva, 2010. Poultry feed metabolizable energy determination using total or partial excreta collection methods. *Braz. J. Poult. Sci.*, 12: 129-132
- Fisher, C. and D.W.F. Shannon, 1973. Metabolizable energy determinations using chicks and turkeys. *Brit. Poult. Sci.*, 14: 609-613
- Fisher, C., 2000. Advances in feed evaluation for poultry. In: *Feed Evaluation, Principles and Practice*, pp: 243-268. P.J. Moughan, M.W.A. Verstegen and M.I. Visser-Reyneveld (eds.). Wageningen Academic Press, Netherlands
- Foster, W.H., 1968. The response of brown leghorn and light Sussex laying flocks to dilution of the diet. *Record. Agric. Res.*, 17: 13-17
- Jiang, Z., 2004. *Putting Metabolizable Energy into Context: Animal Nutrition Group*, Vol. 12. Ajinomoto Co (Thailand) Ltd. Ratchathewi, Bangkok, Thailand
- Jorgensen, H., P. Sorensen and B.O. Eggum, 1990. Protein and energy metabolism in broiler chickens selected for either body weight gain or food efficiency. *Brit. Poult. Sci.*, 31: 517-524
- Kras, R.V., A.M. Kessler, A.M.L. Ribeiro, J.D.I. Henn, D.O.S. Santos II, D.P. Halfen and L. Bockor, 2013. Effect of dietary fiber and genetic strain on the performance and energy balance of broiler chickens. *Braz. J. Poult. Sci.*, 15: 15-20
- Kuzmicky, D.D., R.M. Saunders, R.H. Edwards and G.O. Kohler, 1978. Metabolizable energy values of treated wheat bran and wheat millrun protein concentrate. *Poult. Sci.*, 57: 763-767
- Leeson, S. and J.D. Summers, 2001. *Nutrition of the Chicken*, 4<sup>th</sup> edition. University Books, Guelph, Ontario, Canada
- Lodhi, G.N., R. Renner and D.R. Clandinin, 1969. Studies on the metabolizable energy of rapeseed meal for growing chickens and laying hens. *Poult. Sci.*, 48: 964-970
- Longo, F.A., J.F.M. Menten, A.A. Perdosio, A.N. Figueiredo, A.M.C. Racanicci, J.B. Gaiotto and J.O.B. Sorbara, 2004. Determination of the energetic value of corn, soybean meal and micronized full fat soybean for newly hatched chicks. *Braz. J. Poult. Sci.*, 6: 147-151
- Macleod, M.G., 2002. Energy utilization: Measurement and prediction. In: *Poultry Feedstuffs Supply, Composition and Nutritive Value*, pp: 191-217. J.M. McNab and K.N. Boorman (eds.). CABI Publications, Oxon, UK
- Macleod, M.G., J. Valentine, A. Cowan, A. Wade, L. McNeill and K. Bernard, 2008. Naked oats: Metabolizable energy yield from a range of varieties in broilers, cockerels and turkeys. *Brit. Poult. Sci.*, 49: 368-377
- March, B.E. and J. Biely, 1971. Factors affecting the response of chicks to diets of different protein value: breed and age. *Poult. Sci.*, 50: 1036-1040
- McNab, J.M., 2000. Rapid metabolizable energy assays. In: *Farm Animal Metabolism and Nutrition*, pp: 307-315. J.P.F. D'Mello (ed.). CABI Publications, Oxon, UK
- Meloche, K.J., B.J. Kerr, G.C. Shurson and W.A. Dozier III, 2013. Apparent metabolizable energy and prediction equations for reduced-oil corn distillers dried grains with solubles in broiler chicks from 10 to 18 days of age. *Poult. Sci.*, 92: 3176-3183
- Moran, Jr. E.T., 1982. Starch digestion in fowl. *Poult. Sci.*, 61: 1257-1267
- Morris, T., 2004. Nutrition of chicks and layers. *World's Poult. Sci. J.*, 60: 5-18
- Nadeem, M.A., A.H. Gillani, A.G. Khan and M. Nisa, 2005. True metabolizable energy values of poultry feedstuffs in Pakistan. *Int. J. Agric. Biol.*, 7: 990-994
- NRC, 1994. *Nutrition Requirements of Poultry*, 9<sup>th</sup> edition. National Academy Press, Washington, USA
- Palmer-Jones, R. and D. Halliday, 1971. *The Small-scale Manufacture of Compound Animal Feed*, Vol. 67. Tropical products Institute, London
- Pasha, T.N., F.M. Khattak, D.R. Khan and M.A. Jabbar, 2008. Effect of storage time on the rancidity and metabolizable energy of rice polishing in poultry. *Asian Aust. J. Anim. Sci.*, 21: 420-425
- Rao, E.P.V. and C.V. Reddy, 1986. Evaluation of chemical and nutrient composition in raw, de-oiled and parboiled rice polishing and maize. *Int. J. Poult. Sci.*, 21: 72-74
- Ravindran, V., M.R. Abdollahi and S.M. Bootwalla, 2014. Nutrient analysis, metabolizable energy and digestible amino acids of soybean meals of different regions for broilers. *Poult. Sci.*, 93: 2567-2577
- Rostagno, H.S., L.F.T. Albino, J.L. Donzele, P.C. Gomes, R.F. de Oliveira, D.C. Lopes, A.S. Ferreira and S.L.T. Barreto, 2005. *Brazilian Tables for Poultry and Swine: Feed Composition and Nutritional Requirements*, 2<sup>nd</sup> edition. UFV, Vicosa, Minas Gerais, Brazil
- SAS Institute Inc., 1999. *SAS Users Guide, Version 8.1*. SAS Institute Inc., Cary, NC, USA
- Sauvant, D., J.M. Perez and G. Tran, 2004. *Tables of Composition and Nutritional Value of Feed Materials: Pigs, Poultry, Cattle, Sheep, Goats, Rabbits, Horses, Fish*, 2<sup>nd</sup> edition. Wageningen Academic Publishers, the Netherlands and INRA, Paris, France
- Scott, T.A., F.G. Silversides, H.L. Classen, M.L. Swifts, M.R. Bedford and J.W. Hall, 1998. Broiler chick bioassay for measuring the feeding value of wheat and barley in complete diets. *Poult. Sci.*, 77: 449-455
- Sibbald, I.R., 1982. Measurement of bioavailable energy in poultry feeding stuffs: a review. *Can. J. Anim. Sci.*, 62: 983-1048
- Slinger, S.J., I.R. Sibbald and W.F. Pepper, 1964. The relative abilities of two breeds of chickens and two varieties of turkeys to metabolizable dietary energy and dietary nitrogen. *Poult. Sci.* 43: 329-333
- Ten Doeschate, R.A.H.M., C.W. Scheele, V.V.A.M. Schreurs and J.D. Van Der Klis, 1993. Digestibility studies in broiler chickens: influence of genotype, age, sex and method of determination. *Brit. Poult. Sci.*, 34: 131-146
- Toghyani, M., N. Rodgers, M.R. Barekatin, P.A. Iji and R.A. Swick, 2014. Apparent metabolizable energy value of expeller-extracted canola meal subjected to different processing conditions for growing broiler chickens. *Poult. Sci.*, 93: 2227-2236
- Vieira, R.O., P.B. Rodrigues, R.T.F. Freitas, G.A.J. Nascimento, E.L. Silva and R. Hespanhol, 2007. Chemical composition and metabolizable energy of corn hybrids for broilers. *Braz. J. Anim. Sci.*, 36: 832-838
- Wang, X. and C.M. Parsons, 1998b. Dietary formulation with meat and bone meal on a total versus a digestible or bioavailable amino acid basis. *Poult. Sci.*, 77: 1010-1015
- Woyengo, T.A., E. Kiarie and C.M. Nyachoti, 2010. Metabolizable energy and standardized ileal digestible amino acid contents of expeller-extracted canola meal fed to broiler chicks. *Poult. Sci.*, 89: 1182-1189
- Zelenka, J., 1968. Influence of age of the chicken on the metabolizable energy of poultry diets. *Brit. Poult. Sci.*, 9: 135-142
- Zhou, Z., H.F. Wan, Y. Lia, W. Chena, Z.L. Qia, P. Penga and J. Peng, 2010. The influence of the amylopectin amylase ratio in samples of corn on the true metabolizable energy value for ducks. *Anim. Feed Sci. Technol.*, 157: 99-103