



Full Length Article

Biochemical and Molecular Analysis of Laccase Enzyme in Saprobic Fungi; *Sordaria fimicola* (Roberge ex Desm.) Ces. & De Not

Muhammad Ishfaq^{1*}, Nasir Mahmood², Idrees Ahmad Nasir³ and Muhammad Saleem¹

¹Molecular Genetics Research Laboratory, Department of Botany, University of the Punjab (New campus), Lahore, Pakistan

²Department Biochemistry; Department of Human Genetics and Molecular Biology, University of Health Sciences, Lahore

³Centre of Excellence in Molecular Biology, University of the Punjab (New Campus), Lahore, Pakistan

*For correspondence: ishfaq.botany@pu.edu.pk; ishfaq80@hotmail.com

Abstract

Saprobic fungi play an important role in decomposition and thus contributing to the global carbon cycle. *Sordaria fimicola* strains collected from diverse environment were evaluated for their laccase enzyme activity, while *Aspergillus niger* was used as control fungus. In laccase assay, *S. fimicola* strain N6 collected from the station 6 located at the NFS of the EC 1, showed the maximum laccase enzyme activity (1.1 ΔA/min). In the next step, complete laccase gene (1.917 Kb) was amplified and sequenced from the biochemically efficient isolates of *S. fimicola* viz. S1, S2, SF13, N6 and N7 and on comparison with reported sequence of *Neurospora crassa* (Accession no. M18334.1) 245 point mutations, 65 amino acid and post-translational modifications (PTMs) changes were detected in the *in silico* translation proteins. In phylogenetic analysis, S1, S2, SF13, N6 and N7 strains of *S. fimicola* were found in clade-I alone, while reference sequence of *N. crassa* (Accession no. M18334.1) was present in clade-II. The laccase gene sequences were submitted to NCBI data base under accession numbers KM282173, KM282174, KM282172, KM282175 and KM282176 for *S. fimicola* strains S1, S2, SF13, N6 and N7, respectively. The biochemical as well as molecular data of the study depicted that environmental stresses affected the specific genes by bringing in mutations, which may result in genomic diversity among the organisms and their frontier molecules such as proteins. The observed laccase enzyme activity of *S. fimicola*-a non-pathogenic fungus, was found even better and comparable to *A. niger*-a pathogenic fungus. Therefore, being a saprophytic with short life cycle *S. fimicola* can become a fungus of choice to laccase enzyme at large scale. © 2017 Friends Science Publishers

Keywords: *Sordaria fimicola*; Laccase assay; Laccase gene; Phylogenetic analysis

Introduction

Laccases (EC 1.0.3.2) are used as a biocatalyst in many industrial processes including biopulping, bleaching and treatment of industrial waste water (Bourdais *et al.*, 2012). Laccase has ability to oxidize compounds like polyphenols, cyclic diamines, methoxy substituted phenols and other compounds (Baldrian, 2006). Fungi secrete enzymes in the growth media and purification of these secreted enzymes is considered relatively easier than from bacterial or plant sources, in which enzymes mostly are retained inside the cell as described by different workers (Archer *et al.*, 2008; Upadhyay *et al.*, 2016). Due to higher oxidation reduction potential (+800 mV) as compared to bacteria and plant, the fungal laccase have more application in biotechnology and food industry (Minussi *et al.*, 2002). Saprophytic fungi are unable to produce their own food and depend upon their enzymatic system, which breaks the complex biopolymers into simple nutritional

components and subsequently absorbed from their surroundings (Alexopolous *et al.*, 1996). Therefore, saprophytic fungus has potential to produce different extracellular enzymes, including laccases (Tellez-Jurado *et al.*, 2005). Laccase from *Neurospora crassa* is an inducible secretory enzyme and other fungi that have laccase activity include *Trichoderma* and *Botryosphaeria* (Pointing *et al.*, 2005; Ishfaq *et al.*, 2016a).

Genes encoding laccase enzymes have been studied in different filamentous fungi like *Aspergillus niger*, *Aspergillus oryzae* and *Trichoderma reesei* (Hoffmeister and Keller, 2007; Gomaa and Momtaz, 2015). Fungal species belong to Ascomycota have different laccase encoding genes which are involved in the oxidation of syringaldazine dye (Dedeyan *et al.*, 2000). In the current research *S. fimicola* fungus was evaluated for its laccase enzyme potential, further laccase gene analysis was carried out in order to exploit them to produce laccase enzyme at large scale.

Materials and Methods

Procurement of Fungal Strains and Culturing

Strains of *S. fimicola* (Roberge ex Desm.) Ces. & De Not., used in the study has been isolated from the 'Evolution Canyon 1 (EC 1), located at the Mount Carmel, Israel. Whereas, the EC I have two opposite slopes, the southern facing slope (SFS) with dry and harsh environment, which is quite different in flora and fauna from the northern facing slope (NFS) with moist and lush green environment (Nevo, 1995). On the SFS station 1 and 2 are located, while station 6 and 7 are present on the NFS and from each station 15 sample of *S. fimicola* were included, while one *S. fimicola* strain SF13 has been isolated from the premises of the University of Illinois at Urbana-Champaign (UIUC), USA. Potato Dextrose Agar (PDA) medium and PD broth were used for the culture reviving and sub-culturing of *S. fimicola*.

Laccase Assay

Fungal filtrate was separated from fungal mass by using Whatman filter paper No. 1. Bradford assay (Bradford, 1976) was used to estimate total secretory proteins from fungal filtrate. Laccase enzyme activity of *S. fimicola* was evaluated by using syringaldazine dye and absorbance was taken at 525 nm. The reaction mixture components were 0.1 mL syringaldazine (0.1 mM), 0.85 mL sodium citrate (0.1 M, pH-5.0) and 0.05 mL of the fungal growth medium filtrate containing 50 µg of total proteins. Absorbance value was read at 525 nm by using UV spectrophotometer (UV 1800 SHIMADZU), while 0.85 mL of 0.1 M sodium citrate (pH 5.0) and 0.1 mL of 0.1 mM syringaldazine was used as blank. Fifteen continuous readings were taken for each fungal strain filtrate sample with 1 min time interval. The growth medium filtrate if showed high absorbance at 525 nm in different time intervals then it indicated the high enzyme activity in that fungal strain (Kahraman and Gurdal, 2002).

DNA Extraction and Ribotyping

The manual DNA isolation was carried out by 1% CTAB method with some modifications (Saghai-Marooif *et al.*, 1984). The quality of extracted DNA was checked by 1% agarose gel electrophoresis and quantification was carried out by taking absorbance at 260 nm using UV spectrophotometer (UV 1800 SHIMADZU). *S. fimicola* strains found efficient in laccase enzyme production were subjected to ribotyping by amplification of 431 bases long hypervariable (V4) region of 18S rRNA gene as described earlier (Machouart-Dubach *et al.*, 2001).

Laccase Gene Amplification and Sequencing

S. fimicola strains found efficient in laccase assay were

further subjected to laccase gene targeting. Different primers were designed by using the *N. crassa* laccase gene (Accession no. M18334.1) as reference (Table 1). Amplification conditions were optimized by using different melting temperatures and MgCl₂ concentration (1.5-3.0 mM) during PCR, while all other parameters of reaction mixture were kept constant including Go *Taq* flexi[®] Buffer (1x), dNTPs mixture (0.2 mM), Go *Taq*[®] DNA Polymerase (2.5 U), 25 pmol of each upstream as well as downstream primers and template DNA (100 ng) was used in 50 µL reaction mixture. Temperature cycling condition used were; initial denaturation at 94°C for 4 min, followed by 40 cycles of denaturation temperature at 94°C for 1 min, annealing temperature at 50-60°C for 1 min, primer extension at 72°C for 1 min and final extension temperature was 72°C for 7 min, followed by hold at 4.0°C, while lid temperature was adjusted to 105°C to stop the evaporation. After laccase gene amplification, the PCR products were subjected to 1% agarose gel electrophoresis. The amplified DNA bands of laccase gene were purified from the gel and quantified by using *NanoDrop* ND-1000 (SPECTRAMax Plus) and were got sequenced in both directions from the Core Sequencing Facility available at UIUC, USA.

Bioinformatics Tools

The sequencing results of laccase gene from different strains of *S. fimicola* were submitted to NCBI database after peaks analysis by Chromas software. The laccase gene sequences were aligned by using ClastalW software in order to determine sequence variations. For finding protein sequence variations, gene sequences were translated by using protein translation and ExPASy Protparam tools. The post-translational modifications (PTMs) potential was predicted for laccase protein in *S. fimicola* by using different servers like *LysAcet* and *PredMod* for Acetylation, *BPS* for Methylation, *DISPHOS* and *YinOYang* for Phosphorylation and for Glycosylation *NetNGlyc 1.0* and *YinOYang* were used. Molecular Evolutionary Genetics Analysis (MEGA 6.0.5) software was used for phylogenetic analysis (Tamura *et al.*, 2013).

Results

Among the sixty one *S. fimicola* strains evaluated for their laccase enzyme activity, the isolate N6 collected from the station 6 of the EC 1 showed the maximum laccase enzyme activity (1.1 ΔA/min) (Fig. 1). The isolates N7 (1.0 ΔA/min), S1 (0.95 ΔA/min), IQ36.5 (0.9 ΔA/min), S2 (0.85 ΔA/min) and SF13 (0.82 ΔA/min) were found to be second, third, fourth, fifth and sixth most efficient isolates for laccase production (Fig. 1).

Five *S. fimicola* strains with maximum laccase enzyme activity from different environment viz. S1, S2, N6, N7 and SF13, were shortlisted for laccase gene amplification and sequencing. As regards DNA isolation, a genomic DNA

Table 1: Primers used to amplify laccase gene in *S. fimicola*

Sr. No.	Name	Sequence (5'-3')	Expected PCR Product (bp)
1	LACU-F	TCC AGA CTC GGA GGT GAA	798
2	LACU-R	GAA ATG CGA GTG GTA CCA C	
3	LACM-F	GGC ATG CAC CAG CGC AAC	799
4	LACM-R	TGA TGG GGA TCG TGT TGC	
5	LACL-F	GAA CAC CAA CAG CAT CGC	753
6	LACL-R	ACC AAG ACC AAC ACC AGC	

band of ~15 Kb was observed from *S. fimicola* strains by using 1% gel electrophoresis. After DNA extraction *S. fimicola* strains were confirmed by ribotyping for their purity of cultures. Ribotyping results showed that sequences of V4 domain of 18S rRNA gene were 100% similar with each other as well as with previously reported sequence of *S. fimicola* (Accession No AY545724.1). The sequences of V4 domain of *S. fimicola* strains S1, S2, N6, N7 and SF13 were submitted to NCBI data base under accession numbers KF487278, KF487279, KF487281, KF487282 and LM654514, respectively.

After amplification, complete laccase gene including exons and introns was got sequenced from all five shortlisted strain of *S. fimicola* and compared with the reference gene sequence of *N. crassa* (Accession no. M18334.1) by using ClastalW. The deduced laccase proteins from all five strains of *S. fimicola* as well as in reference gene of *N. crassa* (Accession no. M18334.1) were consisted of 619 amino acids (Table 2; Fig. 2). The PTMs of the laccase proteins in *S. fimicola* strains from the opposite slopes of the EC 1 were compared with each other and these were found to be common among all the five strains of *S. fimicola* except for the acetylation by server *predMod* (Table 3). The phylogenetic analysis of Laccase gene using 12 nucleotides sequences available at NCBI, placed the *S. fimicola* strains in clade-I and *N. crassa* in clad-II (Fig. 3).

Discussion

Out of total sixty one strains of *S. fimicola*, the strain N6 collected from the station 6 of the mild NFS showed the maximum laccase enzyme activity, while *S. fimicola* strains IQ36.6, S2, N7 and S1 were found second, third, fourth and fifth most efficient strains for laccase enzyme activity. The average laccase enzyme activity of *S. fimicola* was found comparable to the *A. niger* as described earlier (Ishfaq *et al.*, 2014). Laccase enzyme production has been reported from soil Ascomycota species (Levasseur *et al.*, 2010). Laccases are distributed in *Ascomycetes*, *Deuteromycetes* and *Basidiomycetes*, being particularly abundant in many white rot fungi which are involved in lignin metabolism (Gochev and Krastanov, 2007).

Ribotyping results of S1, S2, N6, N7 and SF13 revealed that sequences of hypervariable V4 domain of 18S rRNA gene were 100% similar with each other (Ishfaq *et al.*, 2016b) as well as with previously reported sequence of *S. fimicola* (Accession No AY545724.1),




Fig. 1: Comparison of *S. fimicola* and *A. niger* laccase enzyme activity

Key: S1: *S. fimicola* from the station 1. S2: *S. fimicola* from the station 2. N6: *S. fimicola* from the station 6. N7: *S. fimicola* from the station 7. AN744 and AN840: Wild isolates of *A. niger*. SF13: *S. fimicola* from UIUC. IQ36.5: *S. fimicola* strains from the station 2

which confirmed that *S. fimicola* strains were original and pure without any contamination. This was in line with the finding of other coworkers, who used 18S rRNA gene sequencing for species identification in different studies (Caetano-Anolles, 2002; Meyer *et al.*, 2010).

Genomic sequence of *Sordaria macrospora* was found most closely related to *N. crassa* (Nowrousian *et al.*, 2010) than other sequenced filamentous fungi; therefore, in the current research, gene sequences of *N. crassa* were used to design the primers for the amplification of *S. fimicola* genes. The complete laccase gene including exons and introns were amplified from all five strains of *S. fimicola* and sequencing results were aligned with the reported laccase gene of *N. crassa* (Accession # M18334.1). In all five strains of *S. fimicola*, a total of 245 base substitutions were observed in the exonic region of laccase gene. As a result of base substitutions, a total of 65 amino acids changes were observed in all five strains of *S. fimicola* (Fig. 2). These amino acids changes were common in all strains of *S. fimicola* with 100% prevalence except M(109)R amino acid shift which was present only in strain N6 and N7 with 40% prevalence (Table 2; Fig. 3) and this amino acid change may have impact on laccase enzyme activity of strains

Table 2: Analysis of laccase proteins derived from different strains of *S. fimicola* by ExPASy ProtParam tool

Strains	No. of Amino acid.	Mol. Wt. of Protein (Da)	PI. of Protein	Gene length (Exons) (bp)	Dominated Amino acid (%)
RefE	619	68120.7	7.67	1857	Gly (G) 9.5
LACS1	619	67956.9	8.31	1857	Gly (G) 9.7
LACS2	619	67956.9	8.31	1857	Gly (G) 9.7
LACN6	619	67981.9	8.50	1857	Gly (G) 9.7
LACN7	619	67981.9	8.50	1857	Gly (G) 9.7
LACSF13	619	67956.9	8.31	1857	Gly (G) 9.7

Key: Laccase from Strain S1: LACS1; Laccase from Strain S2: LACS2; Laccase from Strain N6: LACN6; Laccase from Strain N7: LACN7 and Laccase from Strain SF13: LACSF13. RefE: Laccase complete reference gene protein of *N. crassa* (Accession no. M18334.1); PI: Isoelectric point of protein

Table 3: Predictions of PTMs in laccase proteins by using different servers

Servers used to predict PTMs	Serial no.	Laccase			
		S1	N7	S2	N6
		Acetylation			
	1	K38	K38	K38	K38
<i>LysAcet</i>	2	K104	K104	K104	K104
	1	K2	K2	K2	
	2	Nil	K104	Nil	
	3	Nil	K114	Nil	
	4	K174	K174	K174	
<i>PredMod</i>	5	K256	K256	K256	
	6	K272	K272	K272	Nil
	7	K362	K362	K362	
	8	K461	K461	K461	
	9	K601	K601	K601	
	10	K616	K616	K616	
		Methylation			
	1	K274	K274	K274	K274
<i>BPS</i>	2	K611	K611	K611	K611
		Phosphorylation			
	1	T84	T84	T84	T84
<i>DISPHOS</i>	2	T88	T88	T88	T88
	1	S186	S186	S186	S186
<i>YinOYang</i>	2	S499	S499	S499	S499
		Glycosylation			
	1	N139	N139	N139	N139
<i>NetNGlyc 1.0</i>	2	N282	N282	N282	N282
	1	T81	T81	T81	T81
<i>YinOYang</i>	2	S186	S186	S186	S186

Note: Amino acids abbreviations, K-Lysine (Lys); T-Threonine (Thr); S-Serine (Ser); N-Asparagine (Asn); Key: The grey highlighted shows the PTMs changes in N7 and S2 strains of *S. fimicola*.

N6 and N7 as evident in biochemical assay performed for laccase enzyme. The sequence variations in laccase genes and their proteins were reported earlier in Ascomycetes fungi (Lyons *et al.*, 2003). The laccase genes amplified and sequenced from *Ganoderma lucidum*, *Phlebia brevispora* and *Trametes* exhibited 65–74% nucleotide sequence homology (Galhaup *et al.*, 2002). Laccase enzyme gene and protein isolation has been reported in different fungi, including *A. niger* and *A. oryzae* (Couto and Toca-Herrera, 2006, 2007).

The deduced laccase proteins consisted of 619 amino acids in all strains of *S. fimicola* as well in reference gene (Accession no. M18334.1) of *N. crassa* (Table 2); but *N. crassa* protein has different molecular weight (68120.7 Da) as compared to laccase proteins derived from strains of *S. fimicola* (Table 2). This difference in molecular weight is attributed to the amino acids substitutions at 65 different positions in all five samples of *S. fimicola* as compared to

reference control protein (Fig. 2). This is also evidenced from the laccase protein molecular weight, where molecular weight of the S1, S2 and SF13 were same (67956.9 Dalton); while the molecular weight of N6 and N7 were same (67981.9 Dalton) (Table 2). The laccase protein from *Cryphonectria parasitica* was 591 amino acids long and has 57% similarity with *N. crassa* as reported earlier by Choi *et al.* (1992). Several reports can be referred, on laccase genes in Ascomycota such as *Gaeumannomyces graminis* (Edens *et al.*, 1999), *Magnaporthe grisea* (Iyer and Chattoo, 2003) and *Mauginella* (Palonen *et al.*, 2003). Laccase is monomers having a molecular mass in the range of 40000-130000 Da with a covalently linked carbohydrate content of 10-25% in fungi and 20-45% in plants (Claus, 2003). As regard isoelectric point (pI) value of laccase proteins, N6 and N7 strains showed 8.50, while strains S1, S2 and SF13 showed 8.31; while the (pI) value of reference control laccase protein was 7.67 (Table 2).

CLUSTAL 2.1 multiple sequence alignment

```

LACN6  MFFFGTAAVAGLAFSLVIGAFALGNDRWLLTPVVRQDFVAFGGGGGACNCPFN 60
LACN7  MFFFGTAAVAGLAFSLVIGAFALGNDRWLLTPVVRQDFVAFGGGGGACNCPFN 60
LACSF13 MFFFGTAAVAGLAFSLVIGAFALGNDRWLLTPVVRQDFVAFGGGGGACNCPFN 60
LACS2  MFFFGTAAVAGLAFSLVIGAFALGNDRWLLTPVVRQDFVAFGGGGGACNCPFN 60
LCAS1  MFFFGTAAVAGLAFSLVIGAFALGNDRWLLTPVVRQDFVAFGGGGGACNCPFN 60
RefE   MFFFGTAAVAGLAFSLVIGAFALGNDRWLLTPVVRQDFVAFGGGGGACNCPFN 60
*****

LACN6  RQCMSLGNLDFEGLPTPTGTRRYTLTLSTWNIIGFQVGRKRVWYNNKLIIGPTL 120
LACN7  RQCMSLGNLDFEGLPTPTGTRRYTLTLSTWNIIGFQVGRKRVWYNNKLIIGPTL 120
LACSF13 RQCMSLGNLDFEGLPTPTGTRRYTLTLSTWNIIGFQVGRKRVWYNNKLIIGPTL 120
LACS2  RQCMSLGNLDFEGLPTPTGTRRYTLTLSTWNIIGFQVGRKRVWYNNKLIIGPTL 120
LCAS1  RQCMSLGNLDFEGLPTPTGTRRYTLTLSTWNIIGFQVGRKRVWYNNKLIIGPTL 120
RefE   RQCMSLGNLDFEGLPTPTGTRRYTLTLSTWNIIGFQVGRKRVWYNNKLIIGPTL 120
*****

LACN6  QANMGDYLETVINLAKANGTSLHBMGMBQNTNLDGVNGVTECFIPFNGSSVYFRPA 180
LACN7  QANMGDYLETVINLAKANGTSLHBMGMBQNTNLDGVNGVTECFIPFNGSSVYFRPA 180
LACSF13 QANMGDYLETVINLAKANGTSLHBMGMBQNTNLDGVNGVTECFIPFNGSSVYFRPA 180
LACS2  QANMGDYLETVINLAKANGTSLHBMGMBQNTNLDGVNGVTECFIPFNGSSVYFRPA 180
LCAS1  QANMGDYLETVINLAKANGTSLHBMGMBQNTNLDGVNGVTECFIPFNGSSVYFRPA 180
RefE   QANMGDYLETVINLAKANGTSLHBMGMBQNTNLDGVNGVTECFIPFNGSSVYFRPA 180
*****

LACN6  TVIGTSMWHSFSAVCGNGIVSIVNGFASANYDVLGFPELMDYVTEADKLVLLTQR 240
LACN7  TVIGTSMWHSFSAVCGNGIVSIVNGFASANYDVLGFPELMDYVTEADKLVLLTQR 240
LACSF13 TVIGTSMWHSFSAVCGNGIVSIVNGFASANYDVLGFPELMDYVTEADKLVLLTQR 240
LACS2  TVIGTSMWHSFSAVCGNGIVSIVNGFASANYDVLGFPELMDYVTEADKLVLLTQR 240
LCAS1  TVIGTSMWHSFSAVCGNGIVSIVNGFASANYDVLGFPELMDYVTEADKLVLLTQR 240
RefE   TVIGTSMWHSFSAVCGNGIVSIVNGFASANYDVLGFPELMDYVTEADKLVLLTQR 240
*****

LACN6  AGFPPSNVLPNGFARHPTTGAQVATVSLTKGKRRLRLINTSVEHFVLSVYNSMT 300
LACN7  AGFPPSNVLPNGFARHPTTGAQVATVSLTKGKRRLRLINTSVEHFVLSVYNSMT 300
LACSF13 AGFPPSNVLPNGFARHPTTGAQVATVSLTKGKRRLRLINTSVEHFVLSVYNSMT 300
LACS2  AGFPPSNVLPNGFARHPTTGAQVATVSLTKGKRRLRLINTSVEHFVLSVYNSMT 300
LCAS1  AGFPPSNVLPNGFARHPTTGAQVATVSLTKGKRRLRLINTSVEHFVLSVYNSMT 300
RefE   AGFPPSNVLPNGFARHPTTGAQVATVSLTKGKRRLRLINTSVEHFVLSVYNSMT 300
*****

LACN6  IAAVLVPPQFVYDLSLGGVGRYDVIIDAAQIGNRYWYVFFSGGSLGCSNMPAFAL 360
LACN7  IAAVLVPPQFVYDLSLGGVGRYDVIIDAAQIGNRYWYVFFSGGSLGCSNMPAFAL 360
LACSF13 IAAVLVPPQFVYDLSLGGVGRYDVIIDAAQIGNRYWYVFFSGGSLGCSNMPAFAL 360
LACS2  IAAVLVPPQFVYDLSLGGVGRYDVIIDAAQIGNRYWYVFFSGGSLGCSNMPAFAL 360
LCAS1  IAAVLVPPQFVYDLSLGGVGRYDVIIDAAQIGNRYWYVFFSGGSLGCSNMPAFAL 360
RefE   IAAVLVPPQFVYDLSLGGVGRYDVIIDAAQIGNRYWYVFFSGGSLGCSNMPAFAL 360
*****

LACN6  PRYQAFNGLPTRNGVAFPPQCLDVLDFLQSLATNSIALATNTYTFITLGGPWR 420
LACN7  PRYQAFNGLPTRNGVAFPPQCLDVLDFLQSLATNSIALATNTYTFITLGGPWR 420
LACSF13 PRYQAFNGLPTRNGVAFPPQCLDVLDFLQSLATNSIALATNTYTFITLGGPWR 420
LACS2  PRYQAFNGLPTRNGVAFPPQCLDVLDFLQSLATNSIALATNTYTFITLGGPWR 420
LCAS1  PRYQAFNGLPTRNGVAFPPQCLDVLDFLQSLATNSIALATNTYTFITLGGPWR 420
RefE   PRYQAFNGLPTRNGVAFPPQCLDVLDFLQSLATNSIALATNTYTFITLGGPWR 420
*****

LACN6  VNGTALDVMKFLVLYMGLNYSYSDKLVKVDGQWYKWLIEDQGFSPFPIH 480
LACN7  VNGTALDVMKFLVLYMGLNYSYSDKLVKVDGQWYKWLIEDQGFSPFPIH 480
LACSF13 VNGTALDVMKFLVLYMGLNYSYSDKLVKVDGQWYKWLIEDQGFSPFPIH 480
LACS2  VNGTALDVMKFLVLYMGLNYSYSDKLVKVDGQWYKWLIEDQGFSPFPIH 480
LCAS1  VNGTALDVMKFLVLYMGLNYSYSDKLVKVDGQWYKWLIEDQGFSPFPIH 480
RefE   VNGTALDVMKFLVLYMGLNYSYSDKLVKVDGQWYKWLIEDQGFSPFPIH 480
*****

LACN6  LRGHDFLLGRSFDVAISQRYVNFVAVMRKGNPFRDVMAPKMGILAFRFD 540
LACN7  LRGHDFLLGRSFDVAISQRYVNFVAVMRKGNPFRDVMAPKMGILAFRFD 540
LACSF13 LRGHDFLLGRSFDVAISQRYVNFVAVMRKGNPFRDVMAPKMGILAFRFD 540
LACS2  LRGHDFLLGRSFDVAISQRYVNFVAVMRKGNPFRDVMAPKMGILAFRFD 540
LCAS1  LRGHDFLLGRSFDVAISQRYVNFVAVMRKGNPFRDVMAPKMGILAFRFD 540
RefE   LRGHDFLLGRSFDVAISQRYVNFVAVMRKGNPFRDVMAPKMGILAFRFD 540
*****

LACN6  NPGMAMCHIAMHVSGLSNQFLERAGQNGKISNAKQAFNRCMAYFFONAPFF 600
LACN7  NPGMAMCHIAMHVSGLSNQFLERAGQNGKISNAKQAFNRCMAYFFONAPFF 600
LACSF13 NPGMAMCHIAMHVSGLSNQFLERAGQNGKISNAKQAFNRCMAYFFONAPFF 600
LACS2  NPGMAMCHIAMHVSGLSNQFLERAGQNGKISNAKQAFNRCMAYFFONAPFF 600
LCAS1  NPGMAMCHIAMHVSGLSNQFLERAGQNGKISNAKQAFNRCMAYFFONAPFF 600
RefE   NPGMAMCHIAMHVSGLSNQFLERAGQNGKISNAKQAFNRCMAYFFONAPFF 600
*****

LACN6  KDDSGLSGVRARVYNDM- 619
LACN7  KDDSGLSGVRARVYNDM- 619
LACSF13 KDDSGLSGVRARVYNDM- 619
LACS2  KDDSGLSGVRARVYNDM- 619
LCAS1  KDDSGLSGVRARVYNDM- 619
RefE   KDDSGLSGVRARVYNDM- 619
*****
    
```

Fig. 2: Multiple sequences alignment of laccase proteins isolated from different strains of *S. fimicola* along with reference protein derived from nucleotide sequence of *N. crassa* laccase complete gene (Accession no. M18334.1) with exons only
 Key: RefE: Laccase protein derived from gene of *N. crassa* (Accession no. M18334.1); Laccase from strain S1: LACS1; Laccase from strain S2: LACS2; Laccase from strain N6: LACN6; Laccase from strain N7: LACN7 and Laccase from strain SF13: LACSF13

However, this finding was found in contrary to an earlier report of Terraza *et al.* (2007), according to which (pI) of fungal laccase was in range of 4-5.

As regards PTMs, these were common among all strains of *S. fimicola* except for the acetylation, which was predicted in N7 strain at K104 and K114 (Table 3). *Ceriporiopsis subvermisporea* has a single laccase gene and its multiple isoforms are formed by the process of PTMs




Fig. 3: Phylogenetic relationship of *S. fimicola* based on the laccase gene by using Maximum likelihood method

possibly by glycosylation and phosphorylation as reported by different coworkers (Larrondo *et al.* 2003; Feng *et al.*, 2015).

The Phylogenetic analysis placed the S1, S2, SF13, N6, and N7 in the clade-I along with *S. macrospora* (XM003349355.1) depicting no or negligible intraspecific variation between *Sordaria* spp., while reference sequence of *N. crassa* (Accession no. M18334.1) was found in clade-II (Fig. 3). The fifty specific laccase gene sequences from different organisms including fungi were compared by multiple sequence alignment and results of the phylogenetic analysis reported diversity of the derived proteins (Satpathy *et al.*, 2013). The protein coding genes have been used to study the phylogenetic analysis among Ascomycetes and Zygomycetes fungi as reported earlier by several coworkers (Diezmann *et al.*, 2004; Tanabe *et al.*, 2004). The biochemical and molecular analysis revealed the laccase activity of *S. fimicola* strains better and comparable to *A. niger*. Therefore, being a non-pathogenic with a short life cycle *S. fimicola* can be used as an alternative fungus for *A. niger* to produce laccase on a larger scale.

Acknowledgements

The work was financially supported by Higher Education Commission (HCE), Islamabad, under International Research Support Initiative Program and by the Special research grant given by the Vice Chancellor, University of the Punjab (New campus) Lahore. We gratefully acknowledge Dr. Youfu “Frank” Zhao, Crop Sciences Department, University of the Illinois at Urbana-Champaign, USA, for his help.

References

Alexopolous, C.J., C.W. Mims and I.M. Blackwel, 1996. *Introductory Mycology*, 4th edition. John Wiley and Sons, Inc. New York, USA
 Archer, D.B., I.F. Connerton and D.A. MacKenzie, 2008. Filamentous fungi for production of food additives and processing aids. *Adv. Biochem. Eng. Biotechnol.*, 111: 99–147

- Baldrian, P., 2006. Fungal laccases-occurrence and properties. *FEMS Microbiol. Rev.*, 30: 215–242
- Bourdais, A., F. Bidard, D. Zickler, V. Berteaux-Lecellier, P. Silar and E. Espagne, 2012. Wood utilization is dependent on catalase activities in the filamentous fungus, *Podospora anserine*. *PLoS ONE*, 7: e29820
- Bradford, M.M., 1976. Rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.*, 72: 248–254
- Caetano-Anolles, G., 2002. Tracing the evolution of RNA structure and the rooting of the universal tree of the life. *Nucleic Acids Res.*, 30: 2575–2587
- Choi, G.H., G.L. Thomas and L.N. Donald, 1992. Molecular analysis of the laccase gene from the chestnut blight fungus and selective suppression of its expression in an isoelectric hypovirulent strain. *Mol. Plant-Microbe Int.*, 5: 119–128
- Claus, H., 2003. Laccases and their occurrence in prokaryotes. *Arch Microbiol.*, 179: 145–150
- Couto, S.R. and J.L. Toca-Herrera, 2006. Industrial and biotechnological applications of laccases: a review. *Biotech. Adv.*, 24: 500–513
- Couto, S.R. and J.L. Toca-Herrera, 2007. Laccase production at reactor scale by filamentous fungi. *Biotech. Adv.*, 25: 558–569
- Dedeyan, B., A. Klonowska, S. Tagger, T. Tron, G. Iacazio, G. Gil and J.L. Petit, 2000. Biochemical and molecular characterization of a laccase from *Marasmius quercophilus*. *Appl. Environ. Microbiol.*, 66: 925–929
- Diezmann, S., C.J. Cox, G. Schonian, R.J. Vilgalys and T.G. Mitchell, 2004. Phylogeny and evolution of medical species of *Candida* and related taxa: a multigenic analysis. *J. Clin. Microbiol.*, 42: 5624–5635
- Edens, W.A., T.Q. Goins, D. Dooley and J.M. Henson, 1999. Purification and characterization of a secreted laccase of *Gaeumannomyces graminis* var. *tritici*. *Appl. Environ. Microbiol.*, 65: 3071–3074
- Galhaup, C., H. Wagner, B. Hinterstoisser and D. Haltrich, 2002. Increased production of laccase by the wood-degrading Basidiomycete *Trametes pubescens*. *Enzyme Microbial. Technol.*, 30: 529–536
- Feng, B.Z., P.Q. Li, L. Fu and X.M. Yu, 2015. Exploring laccase genes from plant pathogen genomes: a bioinformatics approach. *Genet. Mol. Res.*, 14: 14019–14036
- Gochev, V.K. and A.I. Krastanov, 2007. Fungal laccases. *Bulg. J. Agric. Sci.*, 13: 75–83
- Gomaa, O.M. and O.A. Momtaz, 2015. Copper induction and differential expression of laccase in *Aspergillus flavus*. *Braz. J. Microbiol.*, 46: 285–292
- Hoffmeister, D. and N.P. Keller, 2007. Natural products of filamentous fungi: enzymes, genes and their regulation. *Nat. Prod. Rep.*, 24: 393–416
- Ishfaq, M., N. Mahmood, I.A. Nasir and M. Saleem, 2014. Molecular and biochemical screening of local *Aspergillus niger* strains efficient in catalase and laccase enzyme production. *Int. J. Agric. Biol.*, 16: 177–182
- Ishfaq, M., N. Mahmood, Q. Ali, I.A. Nasir and M. Saleem, 2016a. Biochemical and molecular studies of various enzymes activity in fungi. *Mol. Plant Breed.*, 7: 1–16
- Ishfaq, M., N. Mahmood, I.A. Nasir and M. Saleem, 2016b. Biochemical and molecular analysis of superoxide dismutase in *Sordaria fimicola* and *Aspergillus niger* collected from different environments. *Pol. J. Environ. Stud.*, PJOES-00785-2016-01
- Iyer, G. and B.B. Chattop, 2003. Purification and characterization of laccase from the rice blast fungus, *Magnaporthe grisea*. *FEMS Microbiol. Lett.*, 227: 121–126
- Kahraman, S.S. and I.H. Gurdal, 2002. Effect of synthetic and natural culture media on laccase production by white rot fungi. *Bioresour. Technol.*, 22: 2–7
- Larrondo, L.F., M. Avila, L. Sala, D. Cullen and T. Vicun, 2003. Heterologous expression of laccase cDNA from *Ceriporiopsis subvermispota* yields copper-activated apoprotein and complex isoform patterns. *Microbiology*, 149: 1177–1182
- Levasseur, A., M. Saloheimo, D. Navarero, M. Andberg, P. Pontarotti, K. Kruus and E. Record, 2010. Exploring laccase-like multicopper oxidase genes from the ascomycete *Trichoderma reesei*: a functional, phylogenetic and evolutionary study. *BMC Biochem.*, 24: 11–32
- Lyons, J.I., S.Y. Newell, A. Buchan and M.A. Moran, 2003. Diversity of ascomycete laccase gene sequences in a southeastern US salt marsh. *Microb. Ecol.*, 45: 270–281
- Machouart-Dubach, C.L., F.D. Martine, I. Chauvin, L.E. Gall, G. Catherine, L. Frederick and D. Francis, 2001. Rapid discrimination among dermatophytes, *Scytalidium* spp. and other fungi with a PCR-Restriction Fragment Length Polymorphism Ribotyping method. *J. Clin. Microbiol.*, 29: 685–690
- Meyer, A., C. Todt, N.T. Mikkelsen and B. Lieb, 2010. Fast evolving 18S rRNA sequences from Solenogastres (Mollusca) resist standard PCR amplification and give new insights into mollusk substitution rate heterogeneity. *BMC Evol. Biol.*, 10: 70
- Minussi, R.C., G.M. Pastore and N. Duran, 2002. Potential applications of laccase in the food industry. *Trends Food Sci. Technol.*, 13: 205–216
- Nevo, E., 1995. Asian, African and European biota meets at “Evolution Canyon” Israel. Local tests of global biodiversity and genetic diversity patterns. *Proc. R. Soc. Lond.*, 262: 149–155
- Nowrousian, M., J.E. Stajich, M. Chu, I. Engh, E. Espagne, K. Halliday, J. Kamerewerd, F. Kempken, B. Knab, H.C. Kuo, H.D. Osiewacz, S. Poggeler, N.D. Read, S. Seiler, K.M. Smith, D. Zickler, U. Kuck and M. Freitag, 2010. *De novo* assembly of a 40 Mb eukaryotic genome from short sequence reads: *Sordaria macrospora*, a model organism for fungal morphogenesis. *PLoS Genetics*, 6: e1000891
- Palonen, H., M. Saloheimo, L. Viikari and K. Kruus, 2003. Purification, characterization and sequence analysis of a laccase from the ascomycete *Mauginiella* sp. *Enzyme Microb. Technol.*, 33: 854–862
- Pointing, S.B., A.L. Pelling, G.J.D. Smith, K.D. Hyde and C.A. Reddy, 2005. Screening of Basidiomycetes and Xylariaceae fungi for lignin peroxidase and laccase gene-specific sequences. *Mycol. Res.*, 109: 115–124
- Saghai-Marouf, M.A., K.M. Soliman, R.A. Jorgensen and R.W. Allard, 1984. Fungal DNA isolation. *Proc. Natl. Acad. Sci.*, 81: 8014–8018
- Satpathy, R., R. Behera, S.K. Padhi and R.K. Guru, 2013. Computational phylogenetic study and data mining approach to laccase enzyme sequences. *J. Phylogenetic Evol. Biol.*, 1: 108
- Tamura, K., G. Stecher, D. Peterson, A. Filipski and S. Kumar, 2013. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol. Biol. Evol.*, 30: 2725–2729
- Tanabe, Y., M. Saikawa, M.M. Watanabe and J. Sugiyama, 2004. Molecular phylogeny of Zygomycota based on EF-1 α and RPB1 sequences: limitations and utility of alternative markers to rDNA. *Mol. Phylogenet. Evol.*, 30: 438–449
- Tellez-Jurado, A., A. Arana-Cuenca, A.E. Gonzalez-Becerra and O. Viniegra-Gonzalez Loera, 2005. Expression of heterologous laccase by *Aspergillus niger* cultured by solid state and submerged fermentations. *Enzyme Microbial. Technol.*, 38: 665–669
- Terraza, C.A., L.H. Tagle, F. Concha and L. Poblete, 2007. Synthesis and characterization of new bi-functional monomers based on germerylene or silarylene units: 4,4'-(R1R2-silylene) bis (phenyl chloroformates) and 4,4'-(R1R2-germylene) bis (phenyl chloroformates). *Desig. Monomers Polym.*, 10: 253–261
- Upadhyay, P., R. Shrivastava and P.K. Agrawal, 2016. Bioprospecting and biotechnological applications of fungal laccase. *Biotechnology*, 6: 15

(Received 08 October 2015; Accepted 01 December 2016)