



Full Length Article

Improvement of Salt Tolerance on *Escherichia coli* by Expression of Agglutinin from *Amygdalus mira* (Koehne) Yü et Lu.

Liping Xu¹, Ximei Ji¹, Guangze Jin², Fachun Guan^{3,4} and Qiuxiang Luo^{1,5*} and Fanjuan Meng^{1*}

¹College of Life Science, Northeast Forestry University, Harbin 150040, China

²Key Laboratory of Sustainable Forest Ecosystem Management-Ministry of Education, School of Forestry, Northeast Forestry University, Harbin 150040, China

³Jilin Academy of Agricultural Science, Changchun 130033, China

⁴Tibet Agriculture and Animal Husbandry College, Nyingchi 860000, China

⁵Key Laboratory of Saline-Alkaline Vegetation Ecology Restoration (SAVER), Ministry of Education, Alkali Soil Natural Environmental Science Center, Northeast Forestry University, Harbin 150040, People's Republic of China

*For correspondence: 18846831735@163.com; 327273176@qq.com

Abstract

Amygdalus mira (Koehne) Yü et Lu also called Tibetan peach, is a wild species belonging to the *Amygdalus* genus, *Roseaceae*. It is distributed in Tibet and possesses tolerance to extreme environment, such as salt, cold and drought and also possesses resistance to disease. In present research, we cloned the *AmAGG* gene in the root of *A. mira* and analyzed its expression pattern in drought roots of *Amygdalus mira* (Koehne) Yü et Lu by qRT-PCR. The result showed the expression of *AmAGG* increased under drought and decreased after re-watered, indicating that the *AmAGG* was induced by drought. Accordingly, the recombinant *AmAGG* protein (pET-*AmAGG*) was induced and purified from *E. coli* strain (*Rossetta*). The optimal expression temperature was 28°C, the optimal induction time was 6 h and the optimum concentration of IPTG was 0.1 mM, through which the protein was induced and expressed with soluble form in *Rossetta* strain. The salt tolerance of *E. coli* expressing the agglutinin was estimated by measuring at OD₆₀₀ (optical density at 600 nm) value of *Rossetta* by supplementing different concentrations of NaCl into the culture media. The results indicated that *Rossetta* strain harboring the *AmAGG* protein was more tolerant to salt stress than that of harbored pET-15b empty vector, revealed the *AmAGG* might improve the salt tolerance of *Rossetta* cells. Our data suggest that *AmAGG* plays a crucial role in drought and salinity. © 2020 Friends Science Publishers

Keywords: *Amygdalus mira* (Koehne) Yü et Lu; Agglutinin; Expression; Purification; Salt stress; Tolerance

Introduction

Biotic and abiotic stress such as drought and salinity greatly impact agricultural development (Naz *et al.*, 2018). Salinity limit water absorb and increase accumulation of sodium in plants. Therefore, it is a top priority to solving the problem of salinity (Nawaz *et al.*, 2019).

To successfully survive in their environment, plants have carefully evolved and able to confront complex and adverse environmental conditions including salt, drought, cold, heat, insect and pathogen infections. Now, salt stress is one of the main adverse environmental factors which limit plant growth, development, distribution and production worldwide. Currently, much researches are focused on the molecular mechanisms that control plant response to salt stress. Therefore, many related genes and/or signaling pathways have been discovered and identified (Liu *et al.*, 2019; Wu *et al.*, 2019).

To date, the role of agglutinin in protecting cells against damage related to biotic and abiotic stresses has been well proved in plants (Jin *et al.*, 2012; Suzuki *et al.*, 2014; He *et al.*, 2017; Eggermont *et al.*, 2018; Mostofa *et al.*, 2018; Mostafa *et al.*, 2018). Agglutinin is a non-immune protein or glycoprotein that is found in plants, animals, microorganisms, and viruses and it specifically combines with different carbohydrates (Nakata *et al.*, 2017). Among them, plant agglutinin is easily purified and their many features have been extensively studied (Umadevi *et al.*, 2017). These specific features may help us to follow up with a more in-depth understanding. However, there are large researches on the role of agglutinin in response to environmental stresses in different plants, studies on agglutinin of *Amygdalus mira* (Koehne) have been rarely reported (Kil'dibekova *et al.*, 2004; Bezrukova *et al.*, 2008).

Amygdalus mira (Koehne) Yü et Lu is an important wild peach species that possesses strong adaptability to cold, drought, barrenness and high ultraviolet, which has

additional long life (over 1000 years), abundant fruit and other excellent features, means that it is a valuable resource with high ecological and economic merits and is reputed as an ideal wild peach germplasm for improving cultivated peach plants (Liu *et al.*, 2009; Peng *et al.*, 2015; Cao *et al.*, 2017). As a wild species, it can be used to regain vegetation and control soil run off while showing a tolerance to various environmental stresses (Rzepka-Plevnes *et al.*, 2007; Xing *et al.*, 2015).

Recombinant DNA technologies have made it possible to obtain a broad range of diverse proteins in various types living organisms, even mutant proteins that do not exist in nature (Gessler and Patocchi, 2007; Fujita *et al.*, 2016; Misbah *et al.*, 2017). Accordingly, recombinant proteins expressions and purification have become fundamental techniques in many aspects of life science research (Noi and Chung, 2017). However, the results of these techniques often remain frustratingly elusive (Jensen, 1996). The efficient expression of soluble functional proteins isolated from highly complex crude protein lysates in the host, is necessary for successful purification of recombinant proteins. Currently, *E. coli* was one of the most frequently used host organisms for recombinant protein expression. However, over-expression of eukaryotes proteins in *E. coli* often yields inadequate soluble protein (Chae *et al.*, 2017).

Materials and Methods

Plant Culture and Drought Treatment

The experiments were carried out at Northeast Forestry University. All seeds were obtained from the Tibetan plateau. After being rinsed and immersed in water for 24 h, the seeds were embedded with wet sand and stored at 4°C for 3 months. After the seeds germinated, the seedlings were planted into the polyvinyl chloride (PVC) pots of (12-cm diameter and 12-cm height) filled with wet sand. Young and fresh roots of water deficit treated for 4, 8, 12, 16 and 20 days were gathered and frozen in liquid nitrogen, then stored at -80°C for total RNA extraction. Three independent repetitions were performed for each treatment.

Gene Cloning and Bioinformatics Analysis

The total RNA of the *A. mira* roots was extracted using a plant RNA extraction kit (TSINGKE, China) according to the instructions. cDNA was composited using reverse transcriptase (promega, USA). Specific primers for *AmAGG* cloning were designed by using Premier 5.0 and BioEdit. The *AmAGG*-F (sequence: 5' - ATGGCAGACTTACCAAAGTTTTATGGCAGACTTACCAAAGTTTTT - 3') and *AmAGG*-R (sequence: 5' - TCATCCTGGAAGTTTTTCGAC - 3') were synthesized as the forward and reverse primer, respectively. cDNA of *A. mira* roots for 16 days of drought were used as the templates for *AmAGG* cloning. The obtained PCR fragment was

purified and ligated with the pMD18-T vector (Takara, Japan), and then transformed into the *E. coli* (JM109 strain), and the colonies containing the *AmAGG* were sequenced by TSINGKE biological company.

The *AmAGG* protein sequence was analyzed using the ExPASy website (<https://www.expasy.org/>). Protein hydrophobicity, signal prediction were performed. The phylogenetic tree of *AmAGG* was constructed using MEGA version 5 (Tamura, Peterson, Stecher).

Expression of the *AmAGG* in Root of *Amygdalus mira* (Koehne) Yü et Lu

The qRT-PCR was performed using SYBR Premix Ex Taq II Real-time PCR Master Mix (TAKARA, USA) with a LightCycler480 (Roche, USA) in a total volume of 20 µL. Master Mix are composed of *TaKaRa Ex Taq*® HS, dNTP Mixture, Mg²⁺, Tli RNaseH, SYBR® Green I. The relative expression of the target genes was analysed by the $\Delta\Delta CT$ method. Specific primer pairs were determined using Primer Premier 5.0 software. The forward primer was *AmAGG*1-F: 5'- GAAGACAGCCCCGTGGATT-3' and reverse primer was *AmAGG*1-R: 5'-GGCATTCTCGTITGAAGACC-3'.

Expression and Purification of the *AmAGG* Protein in *E. coli* Cells

For protein prokaryotic expression vector construction, the *AmAGG* was amplified with forward primer *AmAGG*p-F: 5' -CTCGAGATGGCAGACTTACCAA-3' and reverse primer *AmAGG*p-R: 5' -GGATCCTCATCCAAGTTTT-3' (the underlined are XhoI and BamHI sites, respectively) and the purified fragment was constructed into the prokaryotic expression vector pET-15b to generate the recombinant expression plasmid pET-*AmAGG*. Then the recombinant plasmid was transformed into the *E. coli* strain *Rossetta*, which was cultured at 37°C with shaking at 180 rpm. The *E. coli* cells grew until the OD₆₀₀ value reached about 0.5, using the IPTG in the bacterium solution to induce the protein expression. To obtain the appropriate induction conditions, a time gradient and IPTG concentration gradient were set. To obtain the optimal IPTG concentration, 0.1, 0.5 and 1.0 mM were added for protein induction, the time gradient were set for 0, 15, 30, 60, 120, 240, 360 min respectively.

The *Rossetta* cells induced by IPTG at 28°C were collected at 13000 rpm for 2 min and then adding a protein loading buffer on an ice bath for 1 h. The loading buffer contained 125 mM Tris-HCl (pH 6.8), 4% SDS, 20% Glycerol and 200 mM DTT and a trace of bromophenol blue, then the cell suspension was centrifuged and the supernatant was soluble protein. The precipitation was dissolved in a protein lysis buffer and was classified as insoluble protein. The lysis buffer contained 7 mol/L urea, 2 mol/L thiourea, 2% Chaps, 50 mM Tris (pH 8.0), trace bromophenol blue and DTT. The recombinant fusion

protein was dissociated by 14.5% SDS-PAGE and visualized Coomassie brilliant blue (CBB) staining.

To obtain the purified *AmAGG* recombinant protein, the *Rossetta* bacteria solution were cultured at 180 rpm for 2.5 h at 37°C until the OD_{600nm} of the bacterial solution reached 0.4-0.6. 0.1 mM IPTG was added to bacterial solution and incubated at 28°C, 180 rpm for 6 h. The bacteria culture was harvested and resuspended in the binding buffer (50 mM NaH₂PO₄, 300 mM NaCl, 10 mM imidazole, pH 8.0) and the crude protein mixture was centrifuged at 10000 rpm for 10 min. Then, the supernatant was the sample for the recombinant protein to be purified using Novagen's Nitia following the manufacture's guidelines. In brief, 6 mL of 50% Nitia was added to the column and equilibrated with five resin-bed volume of binding buffer, the crude protein was loaded onto the resin-bed. After the resin-bed was washed with washing buffer containing 30, 50 and 80 mM imidazole successively, the purified *AmAGG* protein was eluted by five resin-bed volume of elution buffer containing 150 mM imidazole. The samples of all purification steps were collected for SDS-PAGE analysis.

The purity of the purified *AmAGG* recombinant protein was assessed with Quantity One v4.62 and then the concentration was measured by the Bradford method with bovine serum albumin (BSA) as a standard.

Salt Stress Tolerance Assay of *E. coli* Expressing the Agglutinin Protein

Tolerance to salt stress of OD₆₀₀ was tested in an LB (Luria-Bertani) medium supplemented with different concentration of NaCl. The OD₆₀₀ reached a value of 0.5 and was induced with 0.1 mM IPTG for 6 h as described above. Then, the samples were diluted 100-fold by LB medium supplemented with NaCl to the final concentration of 0, 200, 400, 600, 800 mM, respectively. Empty vector transformed *Rossetta* cells was used as control and the bacteria growth rate was determined by measuring the OD₆₀₀ value at 0, 2, 4, 6, 8 and 10 h, respectively.

Results

Gene Cloning and Bioinformatical Analysis of *AmAGG*

The *AmAGG* gene was amplified using cDNA libraries from drought roots of *A. mira* by ordinary PCR. The obtained *AmAGG* fragment was then ligated into the PET-15b prokaryotic expression vector and was sequenced.

A multiple sequence alignment of *AmAGG* protein and its bioinformatical characters are shown in Fig. 1A-D. The alignment of the *AmAGG* amino acid sequence and other agglutinins from *Rosaceae* show that the *A. mira* was closest to the *prunus persica* (*Amygdalus persica* Linn. var. *persica* f. *duplex* Rehd), with only one amino acid difference between them, i.e, the 74^{CYS} in *A. mira* and 74^{SER} in *prunus persica* (Fig. 1A). Phylogenetic tree analysis of the amino




Fig. 1: Multiple sequence alignment and bioinformatical analysis of *AmAGG*. (A) Alignment of *AmAGG* protein and other agglutinins from *Rosaceae*. The χ^2 test statistic was 0.00 ($P = 1.00000$ with 1 degree[s] of freedom). P -value less than 0.05 is often used to reject the null hypothesis of equal rates between lineages. The analysis involved 3 amino acid sequences. All positions containing gaps and missing data were eliminated. There are a total of 158 positions in the final dataset. Evolutionary analyses were conducted in MEGA5. (B) *AmAGG* protein hydrophobicity and hydrophilicity prediction and (E) signal peptide prediction of *AmAGG* protein (C) Phylogenetic tree analysis of the deduced *AmAGG* protein. *prunus persica* (*Amygdalus persica* Linn. var. *persica* f. *duplex* Rehd), *prunus mume* (*Armeniaca mume* Sieb), *malus domestica* (*Malus pumila* Mill), *pyrus x bretschneideri* (*Pyrus bretschneideri* Rehd), *cucumis sativus* (*Cucumis sativus* L.), *cucumis melo* (*Cucumis melo* L.), *vitis vinifera* (*Vitis* L.)

acid sequence of *AmAGG* indicates that, compared with *malus domestica* (*Malus pumila* Mill), *pyrus x bretschneideri* (*Pyrus bretschneideri* Rehd), *cucumis sativus* (*Cucumis sativus* L.), *cucumis melo* (*Cucumis melo* L.) and *vitis vinifera* (*Vitis* L.), *AmAGG* of *A. mira* was closest to that of *prunus persica* and *prunus mume* (*Armeniaca mume* Sieb) (Fig. 1C).

Protein hydrophobic amino acid composition is the main driving force of protein folding and hydrophilic prediction can reflect the folding of the protein. Hydrophobic/hydrophobic analysis of the *AmAGG* protein was performed using the Kyte and Doolittle algorithm of ProtScale (> 0.5). The hydrophobic region was found in the region of < 0.5 , and the hydrophilic region was in the range of < -0.5 and between $+0.5$ and -0.5 Gender area. Hydrophobicity and Hydrophilicity prediction of the *AmAGG* protein was shows that the *AmAGG* protein had 8 hydrophobic regions and 8 hydrophilic regions (Fig. 1B). Signal peptides prediction results indicated that there was no signal peptide was found in N-terminal of the *AmAGG* protein (Fig. 1D).

Expression of the *AmAGG* in Root of *Amygdalus mira* (Koehne) Yü et Lu

A qRT-PCR analysis of the *AmAGG* was performed to determine changes at the *AmAGG* transcription levels were associated with drought (Fig. 2). To verify how drought




Fig. 2: Quantitative real-time PCR analysis of *AmAGG* in root tissues in response to drought at different time intervals. 0,8,12,16 means day 0,8,12,16 under drought treatment, 20 means day 4 under re-watering




Fig. 3: Expression of *AmAGG* fusion protein. (A) Solubility analysis of expressed *AmAGG*. Coomassie-stained gel was used in SDS-PAGE analysis. Lane M, protein ladder; lane 1, 2, 3, 4 are insoluble protein induced with 0, 0.1, 0.5 and 1.0 mM IPTG respectively; lane 5, 6, 7, 8 are soluble protein induced with 0, 0.1, 0.5 and 1.0 mM IPTG respectively. (B) Time-course of *AmAGG* expression. Lane M, protein ladder; lane 1, 2, 3, 4, 5, 6 and 7 are induced by 0.1 mM IPTG for 0, 15, 30, 60, 120, 240, 360 min respectively

influenced the expression of the *AmAGG* in the of *A. mira* root, we observed the change of expression of *AmAGG*. In Fig. 2, The expression of *AmAGG* was up-regulated in response to drought and exhibited the highest expression at day 16, recovering at day 20. These results suggested that the *AmAGG* was induced by drought.

Expression and Purification of the *AmAGG* Recombinant Protein in *E. coli*

In our study, the prokaryotic expression vector pET-

AmAGG was successfully constructed by genetic engineering method, and the expression condition of pET-*AmAGG* recombinant protein was optimized by designing a series of induction time gradient and inducer concentration gradients. The optimal expression temperature of pET-*AmAGG* was 28°C, the optimal induction time was 6 h, and the optimum concentration of IPTG was 0.1 mM, through which the protein was induced and expressed in a soluble form in the *Rossetta* strain.

To optimize the expression conditions of recombinant the *AmAGG* protein, solubility, IPTG concentration gradient and time course were performed. The results of SDS-PAGE are shown in Fig. 3. In Fig 3A, at 0.1, 0.5 and 1.0 mM IPTG induction, there was a markedly enhanced protein band (below 25 kD), whereas in the control (lane 1 and lane 5) there was no band, indicating that this band may be target protein. Furthermore, most of the recombinant *AmAGG* protein was soluble and a small part was the insoluble inclusion body. Soluble expression can greatly simplify the following protein purification and activity analysis. To obtain the optimal IPTG concentration, 0.1, 0.5 and 1.0 mM were added for protein induction, yet there was no difference in soluble *AmAGG* protein expression with the increase of IPTG concentration (lane 6, 7 and 8). To reduce the damage of IPTG to the cells of *Rossetta* strain, 0.1 mM was confirmed as the optimum concentration for IPTG induction. In Fig. 3B, a time gradient was set to optimize the induction time of the recombinant protein and provide a guarantee for the subsequent induction and purification. The expression of the target protein increased with the increase of induction time (0-360 min), indicating that 360 min was the optimal induction time. The optimal over-expression was obtained with the pET-*AmAGG* after 6 h of induction with IPTG at 28°C.

A Large-scale purification of the induced *AmAGG* recombinant protein with a concentration gradient of imidazole in washing buffer was performed with column chromatography and samples corresponding to each stage subjected to SDS-PAGE are shown in Fig. 4. In Fig. 4, the *AmAGG* recombinant protein was purified after washing with 30 mM imidazole in a washing buffer, and the purified protein was obtained via elution with 50 and/or 80 mM imidazole in a washing buffer. At this point, the elution was not complete and the remaining purified target protein could be further eluted with 150 mM imidazole in an elution buffer. These results indicated that 30 mM imidazole in a washing buffer and 150 mM imidazole in an elution buffer might be the optimal conditions for *AmAGG* recombinant protein purification. The purity of the purified *AmAGG* recombinant protein was assessed with Quantity One v4.6.2 (lane 6-12) and the purity was about 95%. The purified *AmAGG* recombinant protein was then collected and quantified and the results showed that the concentration of the purified protein was 0.28 mg/mL.




Fig. 4: Purification of *AmAGG* recombinant protein expressed in *E. coli*. Lane M, protein ladder; lane 1, crude protein sample extract from *E. coli*; lane 2, sample flow through the column (FT); lane 3-5, lane 6-7 and lane 8-9, washing with 30, 50 and 80 mM imidazole in the washing buffer respectively; lane 10-12, eluting with 150 mM imidazole in the elution buffer




Fig. 5: The effects of *AmAGG* expression on the growth of *E. coli* under salt stress conditions. Compared to the control of *E. coli* in normal condition (A), *E. coli* cell growth after 200 mM NaCl treatment (B), 400 mM NaCl treatment (C), 600 mM NaCl treatment (D) and 800 mM NaCl treatment (E) was determined. pET-15b denoted *E. coli* with empty vector and represented control in (A-E). pET-*AmAGG* denoted *E. coli* with pET-*AmAGG*. The mean value of each transgenic line represented a statistically significant difference with respect to the control, as determined by the LSD-t test (P0.05). Vertical bars indicate the mean \pm SE of three biological independent experiments

Stress Tolerance Assay of *E. coli* Expressing the Agglutinin Protein

To identify the *AmAGG* expression to salt tolerance in *E. coli*, the effect of *AmAGG* expression on salt tolerance of *E.*

coli was determined (Fig. 5). In the control experiment, as shown in Fig. 5A, there was no obvious difference in cell survival between the *AmAGG*-expressing strain and the empty vector control strain in 0-2 h, yet in 2-10 h, the control strain grew better than the *AmAGG*-expressing strain. After treatment with, the *AmAGG*-expressing strain showed higher under 200-800 mM NaCl ($P < 0.05$) survival compared to the control strain, indicating that the expression of *AmAGG* might be conducive for the survival and growth of *E. coli* NaCl stress (Fig. 5 B-E). Specifically, under 800 mM NaCl, both the empty vector and *AmAGG*-expressing strain showed sharply lower OD₆₀₀ value than other conditions throughout the growing period, which means they both suffered from serious salt stress. Compared with the empty vector control strain, however, the *AmAGG*-expressing strain showed a significantly higher OD₆₀₀ value during 2-10 h treatment, especially in 0-4 h, which almost no growth for the control strain, while the *AmAGG*-expressing strain could grow slowly (Fig. 5E). These results indicated that the empty vector was inhibited more severely than the *AmAGG*-expressing strain under salt stress. In other words, the expression of *AmAGG* might increase the salt tolerance of *E. coli* to salt stress.

Discussion

Plants are often influenced by abiotic stresses, such as salinity, drought and low temperature. These stresses can cause adversely affected in plant growth and crop productivity. Therefore, to research the mechanisms of plant adjust to these stresses, many genes have been examined. (Deng *et al.*, 2013).

Even though the research on the role of agglutinin in response to environmental stresses in diverse plants, few studies on *A. mira* have been reported. As a tolerant drought and salt stress plant, there is no genome and transcriptional information of *A. mira* in the database (Bai *et al.*, 2009; Guan *et al.*, 2014). Therefore, currently we cannot explain the mechanism of peptide expression in *E. coli*, the agglutinin gene has been reported to award tolerance to drought and salinity (Alqahtany *et al.*, 2019). In previous studies, some drought response proteins have been identified by proteomic analysis (Cao *et al.*, 2017). To obtain the gene sequences of these drought response proteins, a series of primers were designed according to the results of previous proteomic sequencing and the transcriptional information of *Prunus persica* in NCBI database, which is close to the *A. mira* in phylogenies and might share high protein sequences identity with *A. mira*. Thus, we cloned the *AmAGG* gene from *A. mira* and deduced amino acid of the full-length sequence of agglutinin gene from *A. mira*, and its expression pattern under drought was also examined. It showed only one amino acid different between *A. mira* and *P. persica*, the 74^{CYS} in *A. mira* and 74^{SER} in *P. persica*, which maybe the reason for different resistance in *A. mira* and *P. persica*.

As salt and drought share the similar response pathways (Song et al., 2019), we presumed the agglutinin from *A. mira* might also respond to salt stress and play some roles in salt tolerance of *A. mira*. Recombined *AmAGG* protein was induced and purified from *E. coli*. In the exploration of the conditions of expression, although 37°C is a typically used bacteria breeding temperature, a lower temperature of 28°C was selected as the choice according to experience. Generally, under relatively high temperature (37°C), bacteria are prone to form inclusion bodies showing low or no biological activity. The purified *AmAGG* recombinant protein might provide an essential foundation for exploring the characters and activities of the *AmAGG* protein.

Functional identification of genes in *E. coli* and yeast has been widely used for studying salt-sensitive (Latef et al., 2019). The salt tolerance of *E. coli* expressing the agglutinin was estimated via measurement of the OD₆₀₀ value of the *E. coli* strain which was supplied with different concentrations of NaCl into the cultural media (Udawat et al., 2014; Pathak and Ikeno, 2017; Biradar et al., 2018; Daldoul et al., 2018). Compared with empty vector controls, *AmAGG*-expressing *E. coli* showed increased survival rates following salt stress. Under the same concentration of salt treatment, the growth state of *AmAGG*-expressing *E. coli* was better than that of empty vector, which may be due to increased tolerance to salt stress after recombinant protein expression. It was shown that exogenous supplement of WGA, a wheat germ agglutinin, can reduce the oxidative stress level of wheat seedlings and have a protecting effect on the cell growth in wheat seedling roots under salinity (Kil'dibekova et al., 2004; Bezrukova et al., 2008). Therefore, in our study, the *AmAGG* might have a similar effect on the salt tolerance of plants, and was presumed to be involved in the salt response and play a pivotal role in plant stress tolerance. However, the molecular mechanism has not been completely explained and further experiments are necessary to explore how *AmAGG* is induced in these abiotic stress responses.

Conclusion

In conclusion, expression of the *AmAGG* from drought leaves of *Amygdalus mira* (Koehne) Yü et Lu may improve the salt tolerance of *E. coli*. This study has provided a foundation for further study of the activity and characteristics of agglutinin.

Acknowledgements

This study was supported by the Fundamental Research Funds for Central Universities (No. 2572018CG05; No.2572019AA20 ; No.2572017CA10), the National Natural Science Foundation of China (No. 31500317, No. 31660552) and Open Foundation of Open Grant for Key

Laboratory of Sustainable Forest Ecosystem Management from Northeast Forestry University, Ministry of Education (KFJJ2019YB03).

References

- Alqahtany, M., P. Khadka, I. Niyonshuti, V.R. Krishnamurthi, A.A. Sadoon, Challapalli S.D. ChenJ and Y. Wang, 2019. Nanoscale reorganizations of histone-like nucleoid structuring proteins in *Escherichia coli* are caused by silver nanoparticles. *Nanotechnology*, 30: 14
- Bai, T., Y. Wang, X.Z. Zhang, X.F. Xu and Z.H. Han, 2009. The effect of ultraviolet radiation on water-logging resistance in Tibetan peach. *Afr. J. Biotechnol.*, 8: 6596–6602
- Bezrukova, M., A. Kildibekova and F. Shakirova, 2008. WGA reduces the level of oxidative stress in wheat seedlings under salinity. *J. Plant. Growth. Regul.*, 54: 195–201
- Biradar, H., K. Ratna and P.K. Subudhi, 2018. Transgene pyramiding of salt responsive protein 3-1 (SaSRP3-1) and SaVHAc1 from *Spartina alterniflora* L. enhances salt tolerance in rice. *Front. Plant. Sci.*, 9: 1304
- Cao, Y., Q.X. Luo, Y. Tian and F.J. Meng, 2017. Physiological and proteomic analyses of the drought stress response in *Amygdalus Mira* (Koehne) Yu et Lu roots. *BMC. Plant. Biol.*, 17: 53
- Chae, Y.K., S.H. Kim and J.L. Markley, 2017. Relationship between recombinant protein expression and host metabolome as determined by two-dimensional NMR spectroscopy. *Plos. One.*, 12: e0177233
- Daldoul, S., A.B. Amar, M. Gargouri, H. Limam, A. Mliki and T. Wetze, 2018. A grapevine-inducible gene Vv- α -gal/SIP confers salt and desiccation tolerance in *Escherichia coli* and tobacco at germinative stage. *Biochem. Genet.*, 56: 78–92
- Deng, Y., R. Srivastava and S.H. Howell, 2013. Protein kinase and ribonuclease domains of IRE1 confer stress tolerance, vegetative growth, and reproductive development in Arabidopsis. *P Natl. Acad. Sci. USA*, 110: 19633–19638
- Eggermont, L., K. Stefanowicz and E.J.M. Van Damme, 2018. Nictaba homologs from arabidopsis thaliana are involved in plant stress responses. *Front. Plant. Sci.*, 8: 2218
- Fujita, T., M. Yuno and H. Fujii, 2016. Efficient sequence-specific isolation of DNA fragments and chromatin by in vitro enChIP technology using recombinant CRISPR ribonucleoproteins. *Genes. Cells.*, 21: 370–377
- Gessler, C. and A. Patocchi, 2007. Recombinant DNA technology in apple. *Adv. Biochem. Eng. Biot.*, 107: 113–132
- Guan, F.C., S.P. Wang, R.Q. Li, M. Peng and F.J. Meng, 2014. Genetic diversity of wild peach (*Prunus mira* Koehne kov et. kpst) from different altitudes in the tibetan plateau by pollen morphous and RAPD markers. *Hortscience*, 49: 1017–1022
- He, X., L. Li, H. Xu, J. Xi, X. Cao, H. Xu, S. Rong, Y. Dong, C. Wang, R. Chen, J. Xu, X. Gao and Z. Xu, 2017. A rice jacalin-related mannose-binding lectin gene, OsJRL, enhances *Escherichia coli* liability under high salinity stress and improves salinity tolerance of rice. *Plant Biol.*, 19: 257–267
- Jensen, E.S., 1996. Grain yield, symbiotic N₂ fixation and interspecific competition for inorganic N in pea-barley intercrops. *Plant Soil*, 182: 25–38
- Jin, S., X. Zhang and H. Daniell, 2012. Pinellia ternata agglutinin expression in chloroplasts confers broad spectrum resistance against aphid, whitefly, lepidopteran insects, bacterial and viral pathogens. *Plant Biotechnol. J.*, 10: 313–327
- Kil'dibekova, A.R., M.V. Bezrukova, A.M. Aval'baev, R.A. Fatkhutdinova and F.M. Shakirova, 2004. Mechanisms of protective action of wheat germ agglutinin on cell growth in wheat seedling roots under salinity. *Tsitologiya*, 46: 312–316
- Liu, D., C. Han, X. Deng, Y. Liu, N. Liu and Y. Yan, 2019. Integrated physiological and proteomic analysis of embryo and endosperm reveals central salt stress response proteins during seed germination of winter wheat cultivar zhengmai 366. *BMC. Plant. Bio.*, 19: 1

- Liu, W.S., W. Wei and M. Dong, 2009. Clonal and genetic diversity of *Carex moorcroftii* on the Qinghai-Tibet plateau. *Biochem. Syst. Ecol.*, 70e377
- Latef, A.A.H.A., M.G. Mostofa, M.M. Rahman, I.B. Abdel-Farid and L.S.P. Tran, 2019. Extracts from yeast and carrot roots enhance maize performance under seawater-induced salt stress by altering physio-biochemical characteristics of stressed plants. *J. Plant Growth Regul.*, 1–14
- Misbah, M.H., M. Santos, L. Quintanilla, C. Gunter, M. Alonso, A. Taubert and J.C. Rodriguez-Cabello, 2017. Recombinant DNA technology and click chemistry: a powerful combination for generating a hybrid elastin-like-statherin hydrogel to control calcium phosphate mineralization. *Beilstein. J. Nanotech.*, 8: 772–783
- Mostafa, A., J. Sudisha, J.B. David and L.S.P. Tran, 2018. Legume genetic resources and transcriptome dynamics under abiotic stress conditions. *Plant. Cell. Environ.*, 1: 1972–1983
- Mostofa, M.G., W. Li, K.H. Nguyen, M. Fujita and L.S.P. Tran, 2018b. Strigolactones in plant adaptation to abiotic stresses: An emerging avenue of plant research. *Plant. Cell. Environ.*, 41: 2227–2243
- Naz, N., S. Fatima, M. Hameed, M. Ashraf, M. Naseer, F. Ahmad and A. Zahoor, 2018. Structural and Functional Aspects of Salt Tolerance in Differently Adapted Ecotypes of *Aeluropus lagopoides* from Saline Desert Habitats. *Intl. J. Agric. Biol.*, 20: 41–51
- Nawaz, M., A. Khan and M.Y. Ashraf, 2019. Screening Mungbean Germplasm for Salt Tolerance using Growth Indices and Physiological Parameters. *Intl. J. Agric. Biol.*, 22: 401–406
- Nakata, H., T. Wakayama, T. Asano, T. Nishiuchi and S. Iseki, 2017. Identification of sperm equatorial segment protein 1 in the acrosome as the primary binding target of peanut agglutinin (PNA) in the mouse testis. *Histochem. Cell. Biol.*, 147: 27–38
- Noi, N.V. and Y.C. Chung, 2017. Optimization of expression and purification of recombinant S1 domain of the porcine epidemic diarrhea virus spike (PEDV-S1) protein in *Escherichia coli*. *Biotechnol. Biotech. Eq.*, 31: 619–629
- Pathak, N. and S. Ikeno, 2017. In vivo expression of a short peptide designed from late embryogenesis abundant protein for enhancing abiotic stress tolerance in *Escherichia coli*. *Biochem. Bioph. Res. Co.*, 492: 386–390
- Peng, M., F.C. Guan, L. Tao, R.Q. Li, C. Wang and F.J. Meng, 2015. Analysis of genetic relationship on *Amygdalus mira* (koehne) Ricker with other peach species using simple sequence repeat (SSR). *Biochem. Syst. Ecol.*, 62: 98–105
- Rzepka-Plevnes, D., D. Kulpa, M. Smolik and M. Glowka, 2007. Somaclonal variation in tomato *L-pennelli* and *L-peruvianum* f. *gladulosum* characterized in respect to salt tolerance. *J. Food. Agric. Environ.*, 5: 194–201
- Suzuki, N., R.M. Rivero, V. Shulaev, E. Blumwald and R. Mittler, 2014. Abiotic and biotic stress combinations. *New. Phytol.*, 203: 32–43
- Song, Y., X. Yang, S. Yang and J. Wang, 2019. Transcriptome sequencing and functional analysis of sedum lineare thunb. upon salt stress. *Mol. Genet. Genom.*, 5: 1–13
- Udawat, P., A. Mishra and B. Jha, 2014. Heterologous expression of an uncharacterized universal stress protein gene (SbUSP) from the extreme halophyte, *Salicornia brachiata*, which confers salt and osmotic tolerance to *E. coli*. *Gene.*, 536: 163–70
- Umadevi, C., P. Kalaivani, H. Puschmann, S. Murigan, P.S. Mohan and R. Prabhakaran, 2017. Substitutional impact on biological activity of new water soluble Ni(II) complexes: Preparation, spectral characterization, X-ray crystallography, DNA/protein binding, antibacterial activity and in vitro cytotoxicity. *J. Photoch. Photobio. B.*, 167: 45–57
- Wu, J., Y. Jiang, Y. Liang, L. Chen, W. Chen and B. Cheng, 2019. Expression of the maize myb transcription factor zmyb3r enhances drought and salt stress tolerance in transgenic plants. *Plant. Physiol. Bioch.*, 137: 179
- Xing, C., Y. Tian, F.C. Guan and F.J. Meng, 2015. Evaluation of genetic diversity in *Amygdalus mira* (Koehne) Ricker using SSR and ISSR markers. *Plant. Syst. Evol.*, 301: 1055–1064

(Received 15 May 2019; Accepted 15 October 2019)