



Full Length Article

Effects of Fertilizers on Development of Root-knot Nematode, *Meloidogyne javanica*

Ayatollah Saeedizadeh^{*1}, Fahimeh Niasti², Mohammad Ali Baghaei³, Sara Hasanpour³ and Kayvan Agahi²

¹Assistant Professor, Faculty of Agriculture, Shahed University, Tehran, Iran

²Senior Expert, Faculty of Agriculture, Shahed University, Tehran, Iran

³Khorram Bahar Atis Company, Shahed University, Tehran, Iran

*For correspondence: ayatsaeed314@gmail.com

Abstract

Root-knot nematodes (*Meloidogyne* spp.) are known as one of the most important pests of summer crops. Although the use of synthetic nematicides is as a traditional effective method for their control, but heavy costs and adverse environmental effects are the concerns of this strategy. Hence, as an eco-friendly approach, an experiment was conducted to control *Meloidogyne javanica* infestation on tomato seedlings using four fertilizers including a chemical fertilizer (NPK), an organic fertilizer (Humic acid), a semi-organic fertilizer (Agro-health) and a bio-fertilizer (Azotobarvar-1). Nematicidal capacity of the fertilizers were evaluated on *M. javanica* populations *in vitro* and *in vivo* assays, as well as the effect of the fertilizers on plant growth properties were measured in a free-nematode assay. The fertilizers increased second instar juveniles (J₂s) mortality and decreased egg hatching *in vitro*. The highest J₂s mortality rate was found in Agro-health (64.4%), NPK (56.9%), Azotobarvar-1 (51.8%) and Humic acid (48.6%), respectively. Also, the highest reduction on nematode activity (galls, egg masses, and the reproduction factor) was observed in Agro-health (16.4%), NPK (23.8%), Azotobarvar-1 (26.7%) and Humic acid (30.0%) treatments, respectively, *in vivo*. According to the results, using Agro-health at the dose of 8 g kg⁻¹ soil led to an acceptable level of nematode control. Although the fertilizers were not as effective as synthetic nematicides in controlling *M. javanica*, some of them could be a good alternative to nematicides, because, along with the effect of repressive on the nematode, they also improved host plant growth indices. © 2020 Friends Science Publishers

Keywords: Alternative to pesticide; Eco-friendly strategy; Nematicidal potential; Soil amendment; Suppressive effect

Introduction

Root-knot nematodes, *Meloidogyne* spp. (RKNs) are the most economically damaging plant parasitic nematodes (PPNs) and cause considerable horticultural and field crop losses worldwide (Hajji-Hedfi *et al.*, 2018; Kiriga *et al.*, 2018). Summer crops producers often use soil chemotherapy methods to control RKNs in Iran. Although synthetic nematicides provide a quick and effective control, but the heavy costs and the environmental concerns are their disadvantages. Hence, finding an effective and eco-friendly material or method has been considered to suppression of RKNs populations (Julio *et al.*, 2017).

Research has shown that soil amendments, including micro- and macro-nutrients (Mansourabad *et al.*, 2016), bacterial bio-fertilizers (El-Hadad *et al.*, 2011), chemical fertilizers (Habash and Al-Banna, 2011), composts (Hu and Cao, 2008; Hu and Qi, 2010; Ketabchi *et al.*, 2016), plant extracts (Ntalli and Caboni, 2012) and chopped plant tissues (Soheili and Saeedizadeh, 2017), have repressive effects on populations of RKNs. Habash and Al-Banna (2011) found that phosphonate fertilizers, including calcium

phosphonate, magnesium phosphonate and potassium phosphonate were able to stop egg hatching and increased mortality of *M. incognita* and *M. javanica* infectious juveniles *in vitro*. These fertilizers also reduced root galling of *M. javanica* on host plants in a greenhouse. According to Soheili and Saeedizadeh (2017), treating the soil with Brassicaceous tissues [rapeseed (*Brassica napus* L. cv. Okapi); garden cress, (*Lepidium sativum* L. cv. local of Tehran); and cabbage (*B. oleracea* L. cv. Snow crown)], as green manures caused population suppression and decreased activity (galling and reproduction) of *M. javanica* in a greenhouse.

Despite these promising findings, the effective control of RKNs is still under investigation, and synthetic nematicides are widely used to tackle the menace. Since, the world's attention and approach to organic farming is on the rise, therefore, it has been considered the replacement of eco-friendly materials rather than synthetic pesticides. The current study focused on determining the suppressive effect of the chemical, organic, semi-organic and biological fertilizers on populations of RKN, *M. javanica* in Petri dish and pot experiments.

Materials and Methods

Preparation of *M. javanica* Inoculum

The inoculum of *M. javanica* was prepared according to the single egg mass method. For this purpose, first, tomato roots (cv. Super Chief) with symptoms of RKN were collected from farms in Karaj, Iran. The roots were then washed gently in running tap water. Afterward, *M. javanica* was identified morphologically based on the perineal pattern of females using a microscope with 40x magnification (Cetintas and Cakmak, 2016). Subsequently, in order to proliferate the nematode, the egg masses isolated from the galled roots were added to pots filled with steam-sterilized sandy soil and a tomato seedling, cv. Super Chief. The pots were then stored in a greenhouse (25±2°C temperature, 12 h lighting). After two months, infected roots were harvested, washed, crushed, and disinfected with 1% NaOCl solution. The eggs suspension was then incubated in distilled water at 25±2°C for a week. Afterward, the obtained second-stage juvenile (J₂s) were collected, counted, and considered as *M. javanica* inoculum (Saeedizadeh, 2016).

In vitro Assays

In vitro assays were arranged to determine the effect of the fertilizers; bio-fertilizer (Azotobaryar-1[®], Granular 10%, containing Azotobacter sp., Green Biotech Co., Iran), semi-organic fertilizer (Agro-health[®], Emulsifiable Concentrate 10%, Khorram Bahar Atis Co., Iran), chemical fertilizer (NPK, Powder 20-20-20, Zagross Co., Iran), and organic fertilizer (Humic acid, Emulsifiable Concentrate 10%, Zagross Co., Iran); on J₂s mortality and egg hatching rate (%) of *M. javanica* in Petri dish experiments. The egg hatching assay was conducted using eggs suspension as a population under test. The density of the eggs suspension was estimated by an inverted microscope (40x). The suspension was transferred to plastic Petri dishes (5 cm diameters) with 200 eggs as a replicate. The eggs were exposed to fertilizers at doses of 1, 2, 4 and 8 g L⁻¹ distilled water, individually. A treatment of a reference nematicide, Fenamiphos (Nemacur[®] Granular 10%, Bayer Crop Science AG, Germany) was arranged as a positive control at dosage of 0.5, 1, 1.5 and 2 g L⁻¹ distilled water. Also eggs in distilled water (no fertilizer and no nematicide) were considered as a negative control. The eggs suspensions were incubated at 25±2°C, relative humidity 80% and without light for eight days. Egg hatching rate (%) was determined based on the number of hatched eggs by using an inverted microscope (40x) on the second, fourth, sixth, and eighth days from the start of the assay. Each treatment had three replicates. To check recovery, the eggs were transferred in fresh water and incubated after eighth day of exposure in order to ensure the nematocidal or nematostatic effect of the fertilizers (Habash and Al-Banna, 2011).

To evaluate the effect of the fertilizers on J₂s mortality rate (%) of *M. javanica*, an assay was arranged using J₂s

suspension. A suspension of eggs was incubated at 25±2°C for a week. Then, the emerged J₂s were collected and counted as a population under test. J₂s population was set to 200 J₂s per plastic Petri dish (with 5 cm diameter). The J₂s were exposed to the fertilizers at four doses of 1, 2, 4 and 8 g L⁻¹ distilled water, and the nematicide, Fenamiphos with four doses of 0.5, 1, 1.5 and 2 g L⁻¹ distilled water at 25±2°C, relative humidity 80% and without light. A negative control was also considered (no fertilizer and no nematicide). To estimate J₂s mortality rate (%) the number of dead J₂s was determined every day up to four days using stereomicroscope (40x). Each treatment had three replicates (Petri dishes) (Habash and Al-Banna, 2011).

In vivo Assays

A pot assay was conducted to evaluate suppressive effect of the fertilizers (Azotobarvar-1, Agro-health, NPK and humic acid) on *M. javanica* infestation at the greenhouse. The soil required for the assay was collected from the same field as the inoculum, because it was a previously suitable environment for the growth and proliferation of nematode. The soil texture was a loamy sand with 85% sand, 11% silt, 4% clay, 0.3% organic matter, and a pH of 7.2. The collected soil, approximately 366 kg, was mixed, sieved through a screen (5 mesh) and autoclaved. Then, the suspension of the nematode inoculum was thoroughly mixed in the soil (10 J₂s g⁻¹ soil). The nematode-inoculated soil was divided to 61 masses (5 amendments × 4 doses × 3 host plants) + 1 control (dose 0) in quantities of 6 kg (2 kg for a pot × 3 replicates). The soil masses were thoroughly mixed with the fertilizers (at doses 1, 2, 4, and 8 g kg⁻¹ soil), according to the treatments, individually. It was arranged a treatment of Fenamiphos (at doses 0.5, 1, 1.5 and 2 g kg⁻¹ soil) as a positive control; and a negative control (dose 0 or no fertilizer no nematicide). Then, the treated soils was transferred to plastic pots in quantities of 2 kg. After that, a four-leaf seedling of the host plant, previously grown in a sterile sandy substrate, were planted in the pots, according to treatments, individually. Host plants included tomato, *Lycopersicon esculentum* Mill. cv. Super Chief. It was considered three replicates for each dose. A pot containing one plant was represented as a replicate. The pots were kept in the greenhouse for two months; and were sufficiently irrigated.

In the pot assay, the studied traits of the nematode activity on the host plants included the number of galls and egg masses per root, and reproduction factor (RF). Two months after planting, the roots were collected and washed gently in running tap water. Then, the number of galls per root was counted. To determine egg mass numbers per root, the roots were crushed into pieces of 2 to 3 cm and exposed to a solution of Floxin B, 0.15 g L⁻¹. Then, highlighted egg masses were counted using a stereomicroscope (20x). Also, RF was calculated as an indicator of reproduction and pathogenicity of *M. javanica* as follows: $RF = Pf.Pi^{-1}$, where Pf is the final population (eggs, females, J₃s and J₄s into the

plant's root system plus J₂s and males in the soil) and Pi is the inoculum's initial population (Maleita *et al.*, 2012; Soheili and Saeedizadeh, 2017). To determine of the nematode population in the soil, a 100 g subsample of well mixed soil from each replicate (pot) was processed according to centrifuge or sugar flotation method. The extracted nematode population was counted using a stereomicroscope (40x) and then it was applied to estimate the population of nematodes per 1 kg soil (pot). To estimate the number of females and juveniles (J₃s and J₄s) into the roots, 2 g subsample of well mixed chopped roots was macerated in a Waring blender, and then, the nematode population was counted using a stereomicroscope (40x) in the obtained suspension. The numbers of nematodes present in a root were calculated by multiplying the numbers of nematodes present in the subsample of the root, considering the total weight of the root (Soheili and Saeedizadeh, 2017).

In the second pot assay, the fertilizers effect on plants seedling growth properties were evaluated under the similar conditions to the first pot assay and free of the nematode inoculum. In this assay, after soil treatment with the amendments (according to the doses) and transferring to the pots, three seeds of the host plant (tomato, Super Chief cv.) were sowed in each pot. After germination, the seedlings were thinned so that in each pot remained a seedling. The pots were kept in the greenhouse; and the plants were irrigated as needed. The pot containing one plant was represented a replicate. Two months after sowing seeds, plants were taken, rewetted, weighed fresh root and shoot, and measured length of stem, as host plants growth properties (Soheili and Saeedizadeh, 2017).

Data Analysis

The data was analyzed using SAS software version 9 (2002), SAS Institute Inc., Cary, NC, USA. Charts were drawn using Microsoft Excel 2013 software. Three replicated pots per treatment were arranged in a completely randomized design.

Results

In vitro Assays

The most suppressive effect of the fertilizers was observed on egg hatching rate (%) of *M. javanica* by Agro-health (16.4%), NPK (23.8%), bio-fertilizer (26.7%) and Humic acid (30.0%) treatments, respectively. The egg hatching decreased by increasing the dosage of the fertilizers at the same exposure times (Fig. 1 and 2). Also, the use of the fertilizers increased the mortality rate of J₂s population. So that, the highest level of J₂s mortality was resulted in Agro-health (64.4%), NPK (56.9%), bio-fertilizer (51.8%) and Humic acid (48.6%) treatments, respectively. The dead J₂s population increased as a result of increasing the dose and exposure time (Fig. 3 and 4). The J₂s mortality rate reached the highest level (100%) in treatments of Agro-health and NPK at the final dose (8 g L⁻¹

distilled water), however, the treatments in this dose could not stop the egg hatching completely on the last exposure time (eighth day) (Fig. 1 and 3).

In vivo Assays

All treatments significantly ($P \leq 0.05$) reduced the pathogenicity and reproduction indices (the number of gall and egg mass per root and RF) of *M. javanica*. The suppressive effect of Agro-health on *M. javanica* was ranked second after the nematicide, Fenamiphos. Moreover, its nematode controlling effect was more than that of the bio-fertilizer, Azotobarvar-1, NPK and Humic acid, respectively. The concentration of 8 g kg⁻¹ soil of Agro-health and Azotobarvar-1 was found to have the highest suppression effect, whereas the application of Fenamiphos at 2 g kg⁻¹ of soil highly significantly ($P \leq 0.05$) reduced the galling and reproduction of *M. javanica* (Fig. 5).

Notably, the highest values of plant growth parameters were obtained when the seedlings were treated with both Azotobarvar-1 and Agro-health, while Fenamiphos did not improve plant growth compared to the control plants. Moreover, no significant ($P \leq 0.05$) difference was found for shoot and root fresh weight between those pots treated with Azotobarvar-1 and Agro-health (Fig. 6).

Discussion

The acceptance of organic and semi-organic fertilizers for controlling PPNs is considerably increasing, since most old synthetic nematicides have been put aside due to application heavy costs and their environmental concerns. The results of the current study are supported by findings of others in the field of non-chemical control of phytonematodes, through a soil amendment by plant-based and biological fertilizers, which show the use of non-chemical strategies is the remarkable solution to manage PPNs (Hu and Qi, 2010; Oka, 2010; Mcorley, 2011; Tranier *et al.*, 2014; Mansourabad *et al.*, 2016; Kepenekci *et al.*, 2017; Soheili and Saeedizadeh, 2017). Furthermore, application of plant extracts has been effective in controlling these nematodes *in vitro* and *in vivo* (Ntalli and Caboni, 2012; Aissani *et al.*, 2013; Asadi Sardari *et al.*, 2015). Likewise, the results of the current work showed that Agro-health, as a fertilizer based on herbal extracts, has a considerable potential to contribute as a successful environmental-friendly fertilizer in the management of *M. javanica*.

Active ingredient of Agro-health contains a mixture of herbal essential oils; castor-oil (*Ricinus communis* L.), cinnamon (*Cinnamomum verum* J. Presl); acetic acid and nitrate fertilizer. It is expected the suppressive effect of Agro-health on the RKN is related to the presence of plant and non-herbal compounds in the product. The anti-nematode ability of castor-oil, cinnamon, nitrate and acetic acid have been previously reported (Kong *et al.*, 2007; Dong *et al.*, 2012; Pan *et al.*, 2015; Ntalli *et al.*, 2016). Ozdemir and Gozel (2017)




Fig. 1: Effect of amendments (NPK, Humic acid, Agro-health, Azotobarvar-1 and Fenamiphos) and their dose levels on egg hatching of *Meloidogyne javanica* at exposure times

* Defined doses at the X axis (0, I, II, III and IIII) respectively correspond to 0, 1, 2, 4 and 8 g L⁻¹ distilled water for the fertilizers; and 0.5, 1, 1.5, 2 and 2.5 g L⁻¹ distilled water for the nematicide (Fenamiphos)




Fig. 2: Effect of amendments (NPK, Humic acid, Agro-health, Azotobarvar-1 and Fenamiphos) on egg hatching of *Meloidogyne javanica* at exposure times




Fig. 3: Effect of amendments (NPK, Humic acid, Agro-health, Azotobarvar-1 and Fenamiphos) and their dose levels on J₂s mortality of *Meloidogyne javanica* at exposure times

* Defined doses at the X axis (0, I, II, III and IIII) respectively correspond to 0, 1, 2, 4 and 8 g L⁻¹ distilled water for the fertilizers; and 0.5, 1, 1.5, 2 and 2.5 g L⁻¹ distilled water for the nematicide (Fenamiphos)

found that herbal essential oils have had a high potential to become an alternative control strategy against PPNs. In a pot experiment by using urea coated with different doses, Wani and Bhat (2012) reported soil amendment with castor-oil

reduced *M. incognita* populations, and developed plant growth properties and increased chlorophyll content of host plant leaves. Also, the intercropping of castor-oil with cucumber (as a main crop) reduced the pathogenicity activity




Fig. 4: Effect of amendments (NPK, Humic acid, Agro-health, Azotobarvar-1 and Fenamiphos) on J2s mortality of *Meloidogyne javanica* at exposure times




Fig. 5: Effects of treatments of soil amendments (NPK, Humic acid, Agro-health, Azotobarvar-1 and Fenamiphos) on the number of galls, egg masses, and reproduction factor of *Meloidogyne javanica* on the root of tomato plants




Fig. 6: Effect of soil amendments (NPK, Humic acid, Agro-health, Azotobarvar-1 and Fenamiphos) on growth properties (fresh weight of root and shoot and length of stem) of tomato seedlings in the free-nematode treatments

of *M. incognita* by 71.54% (Dong *et al.*, 2012). Kong *et al.* (2007) reported that cassia [*Cinnamomum cassia* (L.) J. Presl] and cinnamon oil compounds have suppressive activity on pinewood nematode, *Bursaphelenchus xylophilus* (Sterner and Buhner) Nickle. Cassia and cinnamon oils and test compounds described merit further study as

potential nematicides or leads for the control of pine wilt disease caused by *B. xylophilus*. The results of current study showed that, in addition to controlling *M. javanica* galling and reproduction, Agro-health also caused a significant promotion in plant growth indices which is due to the existence of a significant amount of nitrate in it.

It has been reported about the potential nematicides nitrogenous compounds (Farahat *et al.*, 2012; Pan *et al.*, 2015). The use of organic compounds such as acetic acid, leading to structural deformities J₂s of *M. incognita*. Obviously, an increase in the number of damaged and dead J₂s will reduce their penetration and pathogenicity (Ntalli *et al.*, 2016).

It seems the application of the semi-organic fertilizers; a mixture of essential oils of plants and chemical fertilizers; could be sound alternative strategy of RKNs management as it may lessen the problems of application of synthetic nematicides therewith improve the net income product. Although the active ingredients in Agro-health are not known as a reference nematicide, but they reduced the pathogenicity and reproduction indices of the RKN to an acceptable level and hence this product could be a suitable alternative to chemical nematicides because of the significant improvement in plant growth indices as well as its eco-friendly properties. It means, they have not been able to control the severe infestations of PPNs on field soils, but they are able to inhibit the development of PPNs populations by repressive properties.

Since short-term non-chemical strategies do not help to control of RKN, long-term use of Agro-health will certainly have more effective results. However, the effect of organic and semi-organic fertilizers has not been as much as synthetic nematicides in RKNs control, but the fertilizers will not lead to ecological concerns. In addition to developing crop performance, the fertilizers have positive results on nutrition, physical conditions and microbial populations of the soil. It seems these fertilizers need time to demonstrate their nematicidal ability and to be introduced as a real nematicide. More research is needed for better understanding of semi-organic fertilizers, and their nematicidal capacity. However, soil amendment with the fertilizers is recommended to control RKNs due to the positive effect of the fertilizers on growth of plants, and also the suppressive capacity of these materials on soil-inhabitant PPNs populations.

Conclusion

The semi-organic fertilizer, Agro-health could be a suitable alternative to synthetic nematicides, with regard to the significant improvement of plant growth indices as well as its eco-friendly properties. However, further studies are needed for a better understanding of the effects of non-chemical nematicides on the activity of RKNs. Because of the contribution to plant growth and due to the lack of destructive effects on the population of soil saprophytic microbes, the use of the fertilizer would have a positive feedback in the integrated management of RKNs in the future.

Acknowledgment

The authors are grateful to Shahed University for financial support of this research.

References

- Aissani, N., P. Tedeschi, A. Maietti, V. Brandolini, V.L. Garau and P. Caboni, 2013. Nematicidal activity of allylthiocyanate from horseradish (*Armoracia rusticana*) roots against *Meloidogyne incognita*. *J. Agric. Food Chem.*, 61: 4723–4727
- Asadi Sardari, A., A.A. Hojat Jalali, S. Bahraminejad and D. Safaei, 2015. Effect of plant extracts on the mortality of root-knot nematodes' J₂, *Meloidogyne javanica*. *Arch. Phytopathol. Plant Prot.*, 48: 365–375
- Cetintas, R. and B. Cakmak, 2016. *Meloidogyne* species infesting tomatoes, cucumbers and eggplants grown in Kahramanmaraş Province, Turkey. *Turk. J. Entomol.*, 40
- Dong, L., C. Huang, L. Huang, X. Li and Y. Zuo, 2012. Screening plants resistant against *Meloidogyne incognita* and integrated management of plant resources for nematode control. *Crop Prot.*, 33: 34–39
- El-Hadad, M.E., M.I. Mustafa, S.M. Selim, T.S. El-Tayeb, A.E.A. Mahgoub and N.H.A. Aziz, 2011. The nematicidal effect of some bacterial biofertilizers on *Meloidogyne incognita* in sandy soil. *Braz. J. Microbiol.*, 42: 105–113
- Farahat, A.A., A.A. Alsayed, H.S. El-Bbeltagi and N.M. Mahfoud, 2012. Impact of organic and inorganic fertilizers on nematode reproduction and biochemical alterations on tomato. *Not. Sci. Biol.*, 4: 48–55
- Habash, S. and L. Al-Banna, 2011. Phosphonate fertilizers suppressed root-knot nematodes *Meloidogyne javanica* and *M. incognita*. *J. Nematol.*, 43: 95–100
- Hajji-Hedfi, L., E. Rebai, A. Larayedh, H. Regaieg and N. Horrigne-Raouani, 2018. Biological control of *Meloidogyne javanica* on tomato with Dazitol® and soil solarization. *Environ. Sci. Pollut. Res. Intl.*, 25: 17278–17282
- Hu, C. and Z.-P. Cao, 2008. Nematode community structure under compost and chemical fertilizer management practice, in the north china plain. *Exp. Agric.*, 44: 485–496
- Hu, C. and Y. Qi, 2010. Effect of compost and chemical fertilizer on soil nematode community in a Chinese maize field. *Eur. J. Soil Biol.*, 46: 230–236
- Julio, L.F., A. González-Coloma, J. Burillo, C.E. Diaz and M.F. Andrés, 2017. Nematicidal activity of the hydrolate byproduct from the semi industrial vapor pressure extraction of domesticated *Artemisia absinthium* against *Meloidogyne javanica*. *Crop Prot.*, 94: 33–37
- Kepekci, I., O. Dura and S. Dura, 2017. Determination of nematicidal effects of some biopesticides against root-knot nematode (*Meloidogyne incognita*) on kiwifruit. *J. Agric. Sci. Technol.*, 11: 546–551
- Ketabchi, S., H. Charehgani and S. Majzoub, 2016. Impact of rhizosphere antagonistic bacteria and urea fertilizer on root knot nematode (*Meloidogyne incognita*) under greenhouse condition. *J. Anim. Plant Sci.*, 26: 1780–1786
- Kiriga, A.W., S. Haukeland, G.M. Kariuki, D.L. Coyne and N.V. Beek, 2018. Effect of *Trichoderma* spp. and *Purpureocillium lilacinum* on *Meloidogyne javanica* in commercial pineapple production in Kenya. *Biol. Contr.*, 119: 27–32
- Kong, J.O., S.M. Lee, Y.S. Moon, S.G. Lee and Y.J. Ahn, 2007. Nematicidal activity of cassia and cinnamon oil compounds and related compounds toward *Bursaphelenchus xylophilus* (Nematoda: Parasitaphelenchidae). *J. Nematol.*, 39: 31–36
- Maleita, C.M.N., R.H.C. Curtis, S.J. Powers and I. Abrantes, 2012. Host status of cultivated plants to *Meloidogyne hispanica*. *Eur. J. Plant Pathol.*, 133: 449–460
- Mansourabad, M.A., A. Kargar Bideh and M. Abdollahi, 2016. Effects of some micronutrients and macronutrients on the root-knot nematode, *Meloidogyne incognita*, in greenhouse cucumber (*Cucumis sativus* cv. Negin). *J. Crop Prot.*, 5: 507–517
- Mcsorley, R., 2011. Overview of organic amendments for management of plant-parasitic nematodes, with case studies from Florida. *J. Nematol.*, 43: 69–81
- Ntalli, N.G. and P. Caboni, 2012. Botanical nematicides in the mediterranean basin. *Phytochem. Rev.*, 11: 351–359

- Ntalli, N.G., M. Ratajczak, C. Oplos, U. Menkissoglu-Spiroudi and Z. Adamski, 2016. Acetic acid, 2-undecanone, and (E)-2-decenal ultrastructural malformations on *Meloidogyne incognita*. *J. Nematol.*, 48: 248–260
- Oka, Y., 2010. Mechanisms of nematode suppression by organic soil amendments-A review. *Appl. Soil Ecol.*, 44: 101–115
- Ozdemir, E. and U. Gozel, 2017. Efficiency of some plant essential oils on root-knot nematode *Meloidogyne incognita*. *J. Agric. Sci. Technol.*, 7: 178–183
- Pan, K., P. Gong, J. Wang, Y. Wang, C. Liu, W. Li and L. Zhang, 2015. Applications of nitrate and ammonium fertilizers alter soil nematode food webs in a continuous cucumber cropping system in Southwestern Sichuan, China. *Eurasian J. Soil. Sci.*, 4: 287–300
- Saeedizadeh, A., 2016. *Trichoderma viride* and *Pseudomonas fluorescens* CHA0 against *Meloidogyne javanica* in the rhizosphere of tomato plants. *Hellenic Plant Prot. J.*, 9: 28–34
- Soheili, A. and A. Saeedizadeh, 2017. Suppression of brassicaceous tissue on *Meloidogyne javanica* in a rhizosphere. *Intl. J. Agric. Biol.*, 19: 1012–1018
- Tranier, M.S., J. Pognant-Gros, R.D. Quiroz, C.N.A. González, T. Mateille and S. Roussos, 2014. Commercial biological control agents targeted against plant-parasitic root-knot nematodes. *Braz. Arch. Biol. Technol.*, 57: 831–841
- Wani, A. and M.Y. Bhat, 2012. Control of root-knot nematode, *Meloidogyne incognita* by urea coated with Nimin or other natural oils on mung, *Vigna radiata* (L.) R. Wilczek. *J. Biopestic.*, 5: 255–258

(Received 06 May 2019; Accepted 12 July 2019)

IMPRESS