



Full Length Article

Differential Response of Micronutrients and Novel Insecticides to Reduce Cotton Leaf Curl Virus Disease and its Vector in *Gossypium hirsutum* L. Varieties

Humaira Kalsoom¹, Safdar Ali¹, Ghulam Mustafa Sahi¹, Amer Habib¹, Muhammad Ahmad Zeshan², Romana Anjum¹, Maryam Yousaf² and Ahsan Abdullah¹

¹Department of Plant Pathology, University of Agriculture, Faisalabad, 38000, Pakistan

²Department of Plant Pathology, University of Sargodha, Sargodha, Pakistan

*For correspondence: safdar147m@uaf.edu.pk

Abstract

The present research evaluated insecticides flonicamid (ulala, 1 g/L) and pyriproxyfen (flyban, 5 mL/L) applied to control the adults and nymphs of *Bemisia tabaci* under field conditions. Micronutrients (33% ZnSO₄ (zinc sulphate)) 3 g/L of water and 13% H₃BO₃ (boric acid) 2 g/L of water were applied to young cotton plants. Minimum disease severity (26.77%) was recorded in flonicamid treated plants and maximum (27.76%) was in pyriproxyfen as compared to the control (77.25%). Whitefly mortality was higher due to flonicamid application than other treatments. The boric acid application showed good control of whitefly and subsequent reduction in disease severity followed by zinc sulphate. The results revealed that treated plants showed significant response to decrease the disease severity (65.34%) and whitefly infestation (58.62%) in all the varieties. The application of micronutrients proved useful for the management of cotton leaf curl virus disease. © 2019 Friends Science Publishers

Keywords: CLCuV; Viruliferous; Infestation; Flonicamid; Pyriproxyfen; Insecticides

Introduction

Cotton leaf curl disease caused by cotton leaf curl virus (CLCuV) is the most damaging to cotton. CLCuV has single stranded DNA genome that belongs to genus Begomovirus of family Geminiviridae (Idris and Brown, 2004). In Pakistan, the disease gained importance in last few decades when it became a severe problem after 1991-1992 and continued till the development of cotton leaf curl resistant variety CIM-1100 in 1996 (Sharma and Bombawale, 2008). This problem appeared again during 2001 in Burewala, breaking the resistance of all the available cotton cultivars/lines (Mansoor *et al.*, 2003). CLCuV is transmitted by whitefly (*Bemisia tabaci*) that belongs to order Hemiptera and family Aleyrodidae. Virus is transmitted in a persistent and circulative manner (Oliveira *et al.*, 2001).

The symptoms of leaf curl disease comprises of upward curling of leaves with thickened veins which are most distinct on the underside (Mansoor *et al.*, 2003). Infected plants show development of leaf enations (oval or cuplike foliar growth) on the underside of leaf. Leaves become brittle and thicker than normal and produce a rattling sound when plant is shaken (Harrison *et al.*, 1997). Susceptibility of cotton plants to viral attack is based upon the stage of plant growth. Plants in early

stage are more vulnerable to disease attack as compared to later stages (Alegbejo, 2001).

As the virus is not transmitted through soil and seed, the whitefly being the most efficient vector is focused upon in order to reduce the CLCuV transmission (Ali *et al.*, 1995, 2014). Whitefly is mainly controlled by using insecticides with different mode of actions under various agricultural systems (Jadhav *et al.*, 2018; Schlaeger *et al.*, 2018). Conventional insecticides became less effective in controlling whitefly due to development of resistance by repeated and excessive exposure. The problem of insecticidal resistance has put a pressure on the novel insecticides (Nauen, 2015). Flonicamid is a novel insecticide used against sucking insects of order Hemiptera. It produces antifeeding behavior and convulsions in hemipterous insects exposed to it (Morita *et al.*, 2007). Flonicamid through its selective activities increase mortality of whitefly and cause delay in nymphal development (Roditakis *et al.*, 2014). Pyriproxyfen controls all growth stages of whitefly with considerable efficiency. It is one of the novel insecticides to which *B. tabaci* has not developed resistance (Caballero *et al.*, 2013). The other way to reduce whitefly infestation and virus transmission is to use nutrient solutions. Soil conditions have promising effect on improving plant health that is helpful in controlling the insect infestation (Altieri and Nicholls, 1999).

Nutrient application affects the relationship between plant and insect (Abro *et al.*, 2004). Insect pests and pathogenic attack is more prevalent on weak plants that have nutritional deficiency (Huber and Thompson, 2007).

Novel insecticides may give the effective control of whitefly and reduce the CLCuV transmission while nutrients may assist the plant to defend against viral attacks. The present research was conducted to assess the efficiency of insecticides and micronutrients application for the management of CLCuV and its vector.

Materials and Methods

The experiment was conducted at research area Department of Plant Pathology, University of Agriculture, Faisalabad. Land was prepared by properly pulverizing the soil and ploughing. Delinted cotton seed of four varieties (BS-70, CIM-616, MM-58 and S-32) were sown by using randomized complete block design (RCBD) with three replications (10 plants/treatment/variety in each replication). There was 75 cm row to row and 30 cm plant to plant distance. Management of CLCuV and its insect vector was done by using two insecticides (pyriproxyfen and flonicamid (ulala) and two micronutrients solutions (H₃BO₃ and ZnSO₄).

Insecticides were sprayed on cotton plants when whitefly population reached economic threshold level (ETL) *i.e.*, 4-5/leaf. Nutrient solutions *i.e.*, 33% ZnSO₄ and 13% H₃BO₃ was sprayed 15 days after sowing, thereafter, insecticides and micronutrients were sprayed to plants at weekly intervals. Applications were done early in the morning to minimize the chances of drift. The treatments of insecticides pyriproxyfen (flyban) and flonicamid (ulala) and the micronutrients solutions of H₃BO₃ and ZnSO₄ were applied at the dose of 5 ml/L, 1 g/L, 2 g/L and 3 g/L, respectively (Table 1). The control plants were treated only with distilled water.

Disease severity was recorded by counting number of infected leaves and total number of leaves in one plant. Three plants were randomly selected in each row and average was taken and disease severity was determined by following the formula:

$$\text{Disease severity} = \frac{\text{Number of infected leaves}}{\text{Total number of healthy leaves}} \times 100$$

Whiteflies were counted from upper, middle and lower leaves of randomly selected three plants in each row. After counting, their mean was taken and recorded against each variety (Razaq *et al.*, 2003).

Efficacy of treatments was determined by using following formula:

$$\text{Efficacy} = \frac{\text{Disease severity in control} - \text{Disease severity in treated plants}}{\text{Disease severity in control}} \times 100$$

$$\text{Efficacy} = \frac{\text{Whitefly infestation in control} - \text{Whitefly infestation in treated plants}}{\text{Whitefly infestation in control}} \times 100$$

Analysis of data was performed statistically and comparison of means was performed using Least Significance Test

(LSD-Test) with 5% significance level. Statistix 8.1 software was used for analysis of the data (Steel *et al.*, 1997).

Results

Response of Cotton Germplasm against Whitefly Infestation after Treatment with Novel Insecticides

The insecticides (Pyriproxyfen and Flonicamid) were applied to check the whitefly mortality and reduction in CLCuV transmission. Flonicamid at its recommended dose showed maximum mortality of whiteflies and thus found to be the most effective in reducing the whitefly infestation. Pyriproxyfen also gave significant decrease in whitefly infestation as compared to untreated plants (Table 4).

The efficacy of insecticides was also recorded in different cotton cultivars. In case of Flonicamid, S-32 gave maximum mortality and minimum whitefly infestation (1.67/leaf) followed by CIM-616 (1.79/leaf). Minimum mortality and maximum infestation was recorded in MM-58 (2.64/leaf) followed by MM-58 (2.33/leaf). Pyriproxyfen proved most effective in BS-70 with minimum whitefly infestation (2.05/leaf) while BS-70 (2.38/leaf), S-32 (2.91/leaf) and CIM-616 (3.03/leaf) as compared to control (5.10/leaf).

Response of Cotton Germplasm against Whitefly Infestation after Treatment with Nutrients

Application of micronutrients affected the whitefly population. Boric acid at 13% lowered the whitefly population to 37.01%. Zinc sulphate effectively reduced the infestation (29.65) at 33% when compared to non-treated plants (Table 5).

Such effect of micronutrients also varied in cotton varieties individually. Minimum infestation of whitefly was shown by S-32 as a result of boric acid application followed by CIM-616, S-32 and MM-58. Whitefly population decreased to a minimum level in BS-70 when treated with zinc sulphate whereas BS-70, CIM-616 and MM-58 demonstrated promising control of whitefly respectively than untreated check (Table 5).

Efficacy of Insecticides against CLCuV Disease Severity in Different Varieties

The control of whitefly through insecticides significantly reduced CLCuV transmission and disease severity. Flonicamid was the most effective in reducing disease severity (65.34%) while pyriproxyfen also gave considerable (64.06%) disease control as compared to control (Table 2).

Flonicamid and pyriproxyfen affected the disease severity in different varieties. The lowest disease severity was recorded in S-32 treated with flonicamid. Maximum disease severity was shown by MM-58 in case of flonicamid application followed by CIM-616 and BS-70.

Table 1: Treatments applied for the management of CLCuV and its vector

Active ingredient	Trade name	Category	Dose/ha
Pyriproxyfen	FlyBan	Insecticide	1235ml
Fonicamid	Ulala	Insecticide	247g
Zinc sulphate	Kiran	Micronutrient	741g
Boric acid	Boric acid	Micronutrient	494g

Table 2: Effect of insecticides on CLCuV disease severity in different cotton varieties

Insecticides	Varieties				Mean
	BS-70	CIM-616	MM-58	S-32	
Disease severity (%)					
Fonicamid	26.17	27.67	29.23	25.00	26.77
Pyriproxyfen	28.14	27.53	29.35	26.02	27.76
Control	74.26	72.03	79.06	83.62	77.25

 LSD = 2.075 at $P < 0.05$
Table 3: Effect of nutrients on CLCuV disease severity in different cotton varieties

Nutrients	Varieties				Mean
	BS-70	CIM-616	MM-58	S-32	
Disease severity (%)					
Boric acid	37.17	36.35	39.11	35.07	36.93
Zinc sulphate	37.23	36.85	38.62	34.98	36.92
Control	79.68	73.25	82.14	80.05	78.78

 LSD = 2.027 at $P < 0.05$
Table 4: Effect of insecticides on whitefly infestation in different cotton varieties

Insecticides	Varieties				Mean
	BS-70	CIM-616	MM-58	S-32	
Whitefly population					
Fonicamid	2.33	1.79	2.64	1.67	2.11
Pyriproxyfen	2.38	3.03	2.05	2.91	2.59
Control	5.43	5.01	4.99	4.97	5.10

 LSD = 0.012 at $P < 0.05$

Pyriproxyfen performed best on S-32 and exhibited minimum disease severity. CIM-616 responded effectively to pyriproxyfen treatment and showed significantly low disease severity as compared to control. Slightly higher disease severity was recorded in BS-70 and MM-58 respectively).

Efficacy of Nutrients against CLCuV Disease Severity in Different Varieties

Proper plant nutrition strengthens its vigour and stimulates the defence mechanisms. Zinc sulphate gave maximum control (53.13%) of the disease as compared to control. The efficacy of boric acid was also best in reducing the disease severity (53.12%). Zinc sulphate and boric acid have very slight difference in their efficacies.

Zinc sulphate provided minimum disease severity on variety S-32 as compared to non-treated plants of the same variety. Maximum disease severity was found in MM-58 followed by BS-70 and CIM-616. The efficacy of boric acid against CLCuV disease severity was more in S-32 than BS-70, CIM-616 and MM-58 in that order (Table 3).

Table 5: Effect of nutrients on whitefly infestation in different cotton varieties

Nutrients	Varieties				Mean
	BS-70	CIM-616	MM-58	S-32	
Whitefly population					
Boric acid	3.47	3.31	4.09	2.12	3.25
Zinc sulphate	3.29	3.64	4.44	3.13	3.63
Control	4.95	5.23	5.48	4.98	5.16

 LSD = 0.033 at $P < 0.05$
Table 6: Evaluation of different treatments against whitefly population

Treatments	Whitefly population	Efficacy (%)
Fonicamid	1.53	81.18a*
Pyriproxyfen	2.06	74.66b
Boric acid	2.79	65.68c
Zinc sulphate	3.62	55.47d
Control	8.13	

 *Different letters indicating insignificant means at $P < 0.05$
Table 7: Evaluation of different treatments against CLCuV

Treatments	Disease severity (%)	Efficacy (%)
Fonicamid	9.82	74.01b
Pyriproxyfen	11.14	70.61c
Boric acid	8.05	78.75a
Zinc sulphate	7.81	79.38a*
Control	37.79	

 *Different letters indicating insignificant means at $P < 0.05$

Evaluation of Insecticides and Nutrients against Whitefly Infestation and Severity

Whitefly population was significantly reduced by the application of fonicamid (81.18%) followed by pyriproxyfen (74.66%) was the second most efficient insecticide in controlling whitefly followed by boric acid and zinc sulphate (Table 6). Zinc sulphate treatment reduced CLCuV disease severity with 79.38% efficiency. The efficacy of boric acid was high (78.75%) as compared to pyriproxyfen (70.61%) and fonicamid (74.01%, Table 7).

Discussion

Cotton leaf curl virus causes huge losses worldwide and 30% yield reduction is recorded in Pakistan (Hassan *et al.*, 2015). Different insecticides are used against whitefly to minimize the virus transmission and disease severity (Aktar *et al.*, 2008). Commonly used insecticides from organophosphate (OP), organochlorine (OC) and carbamates groups become less effective against whitefly after repeated use (Nauen, 2015). Resistance to insecticides is pretended as the failure of the chemical. That's why OPs and OCs have been replaced with neonicotinoids and novel insecticides (Jeschke *et al.*, 2010). One of the unique novel insecticide is fonicamid that blocks the feeding of sucking insects like whitefly (Roditakis *et al.*, 2014).

In present study, the mean whitefly population was 2.11 from all the varieties treated with fonicamid. Fonicamid was 81.18% efficient in controlling the whitefly

infestation. Previously 95% whitefly mortality have been recorded after exposure to flonicamid (Roditakis *et al.*, 2014). Devine and Denholm (1998) studied the effect of flonicamid application on nymphs of whitefly and found delay in its development. The results of the following experiment revealed that whitefly management by flonicamid affected the disease severity and 74% efficacy was calculated. Flonicamid inhibits the feeding of sucking insects and it has no side effect on natural enemies (Morita *et al.*, 2007).

The whitefly population in pyriproxyfen treated plants was 2.59 and its efficiency was 74.66%. Wen *et al.* (2014) recorded 77% mortality of whitefly eggs through the use of pyriproxyfen. It was further concluded that pyriproxyfen is more toxic against whitefly eggs than neonicotinoids. It binds to the ryanodine receptors and kills the eggs and larvae of the *B. tabaci* (Jeanguenat, 2013). Li *et al.* (2012) reported pyriproxyfen is an anthranilic diamide to which whitefly has no resistance. Reduction in egg hatchability and mortality of larvae was described by Lee *et al.* (2002). Kumar *et al.* (2014) used pyriproxyfen to manage whitefly infestation in tomato plantations 200 g/acre. Pooled analysis of treatments applied showed that maximum whitefly mortality was calculated from pyriproxyfen with 76% efficiency. Pyriproxyfen act as insect growth regulator that suppresses the process of embryo formation and development of adult by badly interrupting with hormones of insects (Ishaaya and Horowitz, 1998). Bi *et al.* (2002) sprayed estereem (Pyriproxyfen) on the leaves of strawberry and found it effective against whitefly egg hatching. These findings are further strengthened by (Mckee and Zalom, 2007) who described that foliar spray of pyriproxyfen at early growth stages minimize the whitefly population. In this study, CLCuV disease severity was reduced upto 70% in the plants treated with flonicamid due to maximum mortality of the insect vector. Detailed study was made on novel insecticides pyriproxyfen, buprofezin, imidacloprid, pymetrozine and diafenthiuron in 1990. It was observed that the insecticides do not directly kill or affect the *Bemisia tabaci* as conventional pesticides (immediately knocking down the pests). Buprofezin adversely affect the female's viability, eggs fertility and larvae molting of insects is highly affected. Pyriproxyfen directly or indirectly affects all stages of life cycle (Dittrich *et al.*, 1990). Three successful applications of novel pesticides pyriproxyfen and flonicamid control the whitefly infestation and subsequently CLCuV transmission successfully. Pyriproxyfen (juvenile hormone mimic) suppresses embryogenesis and adult formation of whitefly (Imtiaz *et al.*, 2010).

Pyriproxyfen mode of action revealed that it acts like analog hormone of juvenile stage and mimics the hormone action (playing major role in control of whiteflies). Pyriproxyfen played significant role in control of *B. tabaci* especially in Israel and North America (Palumbo *et al.*, 2001).

Micronutrients enriched plants are less vulnerable to insect attack as compared to under nourished cotton plants

(Rashdi, 1998). Khan *et al.* (2003) used salicylic acid at 3% concentration that effectively controlled adult whitefly, emergence and egg hatchability. In present study, whitefly infestation was controlled with 65% efficiency in different cotton cultivars by using boric acid. The insecticidal efficacy of boric acid was studied against the sucking insect pests and 81% mortality was recorded (Javaid *et al.*, 2018). Gore and Schal (2004) observed significant mortality of sucking insects in laboratory when treated with aqueous solution of boric acid. Boric acid has been used as a successful chemical to control a wide range of sucking insects in different plantations (Bicho *et al.*, 2015). Deterrence of insects was also observed during the contact with boric acid treated sites (Tsunoda, 2001). Boric acid destroys the inner portion of foregut and produces toxin in the neuron of insects (Habes *et al.*, 2006). Murray (1998) described that boric acid is less toxic to environment and highly water soluble. In present experiment, CLCuV disease severity minimized up to 78% due to boric acid application. Boric acid increases the plant integrity and thus helps to manage the physiological disturbance caused by viral infection. Boric acid caused significant increase in total chlorophyll content and proline of the plants (Köksal *et al.*, 2018). Evaluation of zinc sulphate against whitefly population indicated 55% mortality. This finding is supported by the results obtained from experiments of (Sharaby *et al.*, 2013) where plant yield is increased due to zinc sulphate application. Actually, zinc sulphate acts as protective chemical that disrupt the endocrine system of insect. Padhee and Mishara (1993) recorded that zinc application induce resistance in plants against insect pests. Sarwar (2011) found that zinc sulphate affects the growth, development and reproduction of insects.

Plant nutrition should be improved to provide an active defence system against attacking virus (Pervez *et al.*, 2007). Nutrients can decrease disease occurrence to an acceptable level by improving plant growth and development. Application of adequate nutrients improves plant resistance by enhancing defence system. In current study, boric acid and zinc sulphate effectively reduced CLCuV disease severity and their efficacy was 79 and 75, respectively. Provision of zinc sulphate and boric acid controlled disease incidence to an appropriate level (Abro *et al.*, 2004). Zinc is involved in many physiological functions and limited availability of this vital nutrient limits plant yield by stunting the growth, increasing crop maturity time, chlorosis, shorter leaves and inferior production quality (Butler *et al.*, 1991). Virus infections alters the synthesis and functions of plant hormones especially indole acetic acid. Zinc takes part in the synthesis of tryptophan that is used for the indole acetic acid formation. Indole acetic acid triggers the plant defense mechanism (Spiegel-Roy and Goldschmidt, 2008). Puzina (2004) described that zinc sulphate and boric acid application enhances the hormonal status of plants which contributes towards increased growth and development. Geminiviruses interrupt with plant cell

cycle by disturbing proteins that are integral part of cell cycle (Kong *et al.*, 2000). Boron increases the vegetative growth by boosting carbohydrate and protein metabolism (Asad *et al.*, 2003) and provides integrity to the cell wall in plants (Altieri and Nicholls, 1999).

Conclusion

It can be concluded that insecticides (Fonicamid and Pyriproxyfen) effectively controlled the whitefly infestation that resulted in significant reduction in disease severity. Nutrients application not only enhanced the plant defence against CLCuV but also created harsh conditions for the whitefly vector.

References

- Abro, G., T. Syed and M. Unar, 2004. Effect of application of a plant growth regulator and micronutrient on insect pest infestation and yield components of cotton. *J. Entomol.*, 3: 243–248
- Aktar, M.M., M.S. Akhter and A.M. Akanda, 2008. Impact of insecticides and organic oil spray on the growth and yield of tomato under TYLCV infected condition. *Bangl. Res. Publ. J.*, 1: 199–205
- Alegbejo, M., 2001. Whitefly transmitted plant viruses in Nigeria. *J. Sustain. Agric.*, 17: 99–109
- Ali, M., Z. Ahmad, M. Tanveer and T. Mahmood, 1995. *Cotton Leaf Curl Virus in the Punjab: Current Situation and Review of Work*. Central Cotton Research Institute, Multan/CLCV Project, Ministry of Food, Agriculture and Livestock, Government of Pakistan/Asian Development Bank
- Ali, S., M.A. Khan, S.T. Sahi, M. Atiq and A. Hannan, 2014. *Cotton Leaf Curl Virus Disease Predictive Model based on Environmental Variables*. In: *Improvement of Crops in the Era of Climatic Changes*, pp: 323–335. Springer
- Altieri, M.A. and C.I. Nicholls, 1999. Biodiversity, ecosystem function, and insect pest management in agricultural systems. *Agric. Ecosys. Environ.*, 74: 19–31
- Asad, A., F.P.C. Blamey and D.G. Edwards, 2003. Effects of boron foliar applications on vegetative and reproductive growth of sunflower. *Ann. Bot.*, 92: 565–570
- Bi, J.L., N.C. Toscano and G.R. Ballmer, 2002. Greenhouse and field evaluation of six novel insecticides against the greenhouse whitefly *Trialeurodes vaporariorum* on strawberries. *Crop Prot.*, 21: 49–55
- Bicho, R.C., S.I. Gomes, A.M. Soares and M.J. Amorim, 2015. Non-avoidance behaviour in enchytraeids to boric acid is related to the GABAergic mechanism. *Environ. Sci. Pollut. Res.*, 22: 6898–6903
- Butler Jr, G., S. Puri and T. Henneberry, 1991. Plant-derived oil and detergent solutions as control agents for *Bemisia tabaci* and *Aphis gossypii* on cotton. *Southwestern Entomol.*, 16: 331–337
- Caballero, R., S. Cyman, D.J. Schuster, H.E. Portillo and R. Salter, 2013. Baseline susceptibility of *Bemisia tabaci* (Genn.) biotype B in southern Florida to cyantraniliprole. *Crop Prot.*, 44: 104–108
- Devine, G.J. and I. Denholm, 1998. An unconventional use of piperonyl butoxide for managing the cotton whitefly, *Bemisia tabaci* (Hemiptera: Aleyrodidae). *Bull. Entomol. Res.*, 88: 601–610
- Dittrich, V., S. Uk and G. Ernst, 1990. Chemical control and insecticide resistance of whiteflies. In: *Whiteflies: their Bionomics, Pest Status and Management*, pp: 263–285. Gerling, D. (ed.). Intercept, Andover, Hants
- Gore, J.C. and C. Schal, 2004. Laboratory evaluation of boric acid sugar solutions as baits for management of German cockroaches infestations. *J. Econ. Entomol.*, 97: 581–587
- Habes, D., S. Morakchi, N. Aribi, J.P. Farine and N. Soltani, 2006. Boric acid toxicity to the German cockroach, *Blattella germanica*: alterations in midgut structure, and acetylcholinesterase and glutathione S-transferase activity. *Pest. Biochem. Physiol.*, 84: 17–24
- Harrison, B., Y. Liu, S. Khalid, S. Hameed, G. Otim-Nape and D. Robinson, 1997. Detection and relationships of cotton leaf curl virus and allied whitefly-transmitted geminiviruses occurring in Pakistan. *Ann. Appl. Biol.*, 130: 61–75
- Hassan, F., A. Qayyum, W. Malik, A. Maqbool, M. Hassan, M.A. Rehman, M. Shoaib, M. Shehzad, S. Ahmad, L. Ahmad and M. Arshad, 2015. Cotton Leaf Curl Virus (CLCuV) Disease in Pakistan: A Critical Review. *Appl. Sci. Business Econ.*, 3: 8–14
- Huber, D.M. and I.A. Thompson, 2007. Nitrogen and plant disease. In: *Mineral Nutrition and Plant Disease*, pp: 278–279. Datnoff, L.E., W.H. Elmer and D.M. Huber (eds.). APS Press, USA
- Idris, A. and J. Brown, 2004. Cotton leaf crumple virus is a distinct western hemisphere begomovirus species with complex evolutionary relationships indicative of recombination and reassortment. *Phytopathology*, 94: 1068–1074
- Imtiaz, M., A. Rashid, P. Khan, M. Memon and M. Aslam, 2010. The role of micronutrients in crop production and human health. *Pak. J. Bot.*, 42: 2565–2578
- Ishaaya, I. and A.R. Horowitz, 1998. Insecticides with novel modes of action: An overview. In: *Mechanism and Application*, pp: 1–24. Ishaaya, I. and D. Degheele (eds.). Springer-Verlag, Berlin Heidelberg
- Jadhav, R.D., D.R. Kadam, P.B. Jadhao and G.K. Kangale, 2018. Bioefficacy of newer insecticides against brinjal whitefly and their natural enemies. *J. Entomol. Zool. Stud.*, 6: 1187–1191
- Javaid, M.A., A. Rasool and A. Yaseen, 2018. Bio-efficacy of enhanced diatomaceous earth with boric acid against Citrus mealy bug (*Planococcus citri*). *J. Agric. Sci. Bot.*, 2: 29–33
- Jeanguenat, A., 2013. The story of a new insecticidal chemistry class: the diamides. *Pest Manag. Sci.*, 69: 7–14
- Jeschke, P., R. Nauen, M. Schindler and A. Elbert, 2010. Overview of the status and global strategy for neonicotinoids. *J. Agric. Food Chem.*, 59: 2897–2908
- Khan, M., Q. Nadeem, S. Khan and M. Nasir, 2003. Effect of salicylic acid, KH_2PO_4 and K_2HPO_4 on the egg hatchability, adult emergence and population of *Bemisia tabaci* and cotton leaf curl virus severity. *Pak. J. Bot.*, 35: 983–986
- Köksal, K., K. Kakyüz, M. Yıldıztekin, H.S. Civelek, S. Nadeem and A.L. Tuna, 2018. Antioxidative Defence Mechanisms in Tomato (*Lycopersicon esculentum* L.) Plants Sprayed with Different Pesticides and Boron compounds. *Intl. J. Sec. Met.*, 5: 200–209
- Kong, L.J., B.M. Orozco, J.L. Roe, S. Nagar, S. Ou, H.S. Feiler, T. Durfee, A.B. Miller and W. Gruissem, 2000. A geminivirus replication protein interacts with the retinoblastoma protein through a novel domain to determine symptoms and tissue specificity of infection in plants. *EMBO J.*, 19: 3485–3495
- Kumar, V., R.S. Chandi, H.S. Bhullar and A. Dhawan, 2014. Pyriproxyfen against whitefly, *Bemisia tabaci* (Gennadius) on tomato. *Pest. Res. J.*, 26: 144–149
- Li, X.C., B.A. Degain, V.S. Harpold, P.G. Marçon, R.L. Nichols, A.J. Fournier, S.E. Naranjo, J.C. Palumbo and P.C. Ellsworth, 2012. Baseline susceptibilities of B- and Q-biotype *Bemisia tabaci* to anthranilic diamides in Arizona. *Pest Manage. Sci.*, 68: 83–81
- Lee, Y.S., E.C. Park, J.H. Kim and G.H. Kim, 2002. Comparative toxicities of pyriproxyfen and thiamethoxam against the sweetpotato whitefly *Bemisia tabaci* (Homoptera: Aleyrodidae). *J. Asia Pac. Entomol.*, 5: 117–122
- Mansoor, S., I. Amin, S. Iram, M. Hussain, Y. Zafar, K. Malik and R. Briddon, 2003. Breakdown of resistance in cotton to cotton leaf curl disease in Pakistan. *Plant Pathol.*, 52: 784–784
- McKee, G.J. and F.J. Zalom, 2007. Management and yield impact of the greenhouse whitefly on California strawberries. *Hortic. Sci.*, 42: 285–288
- Morita, M., T. Ueda, T. Yoneda, T. Koyanagi and T. Haga, 2007. Fonicamid, a novel insecticide with a rapid inhibitory effect on aphid feeding. *Pest Manage. Sci.*, 10: 969–973
- Murray, F.J., 1998. A comparative review of the pharmacokinetics of boric acid in rodents and humans. *Biol. Trace Elem. Res.*, 66: 331–341

- Nauen, R., 2015. Development of a lateral flow test to detect metabolic resistance in *Bemisia tabaci* mediated by CYP6CM1, a cytochrome P450 with broad spectrum catalytic efficiency. *Pest. Biochem. Physiol.*, 121: 3–11
- Oliveira, M., T. Henneberry and P. Anderson, 2001. History, current status, and collaborative research projects for *Bemisia tabaci*. *Crop Prot.*, 20: 709–723
- Padhee, A.K. and D.S. Mishara, 1993. Influence of Zinc application on the influence of rice leafhopper, *Cnaphalocrocis medinalis* Guenee. *Ind. J. Entomol.*, 55: 335–336
- Palumbo, J., A. Horowitz and N. Prabhaker, 2001. Insecticidal control and resistance management for *bemisia tabaci*. *Crop Prot.*, 20: 739–765
- Pervez, H., M. Ashraf, M.I. Makhdom and T. Mahmood, 2007. Potassium nutrition of cotton (*Gossypium hirsutum* L.) in relation to cotton leaf curls virus disease in arid soils. *Pak. J. Bot.*, 39: 529–539
- Puzina, T.I., 2004. Effect of zinc sulfate and boric acid on the hormonal status of potato plants in relation to tuberization. *Russ. J. Plant Pathol.*, 51: 209–215
- Rashdi, S.M., 1998. *Host Plant Resistance of Bioregulator Treated Cotton to Bollworms and Sucking Complex and its Impact on Yield and Yield Components*, p: 37. Final Research Report. Nuclear Institute of Agriculture, Tandojam, Pakistan
- Razaq, M., M. Aslam, M. Iqbal, M. Hussain and A. Shahzad, 2003. Efficacy of six Insecticides with different mode of actions against sucking insect pests of cotton. *Sarhad J. Agric.*, 19: 97–100
- Roditakis, E., N. Fytrou, M. Staurakaki, J. Vontas and A. Tsagkarakou, 2014. Activity of flonicamid on the sweet potato whitefly *Bemisia tabaci* (Homoptera: Aleyrodidae) and its natural enemies. *Pest Manage. Sci.*, 70: 1460–1467
- Sarwar, M., 2011. Effect of Zinc fertilizer application on the incidence of rice stem borer (*Scirpophaga* species) (Lepidoptera: Pyralidae) in rice (*Oryza sativa* L.). *Crop J. Cereal Oilseeds*, 2: 61–65
- Schlaeger, S., J.A. Pickett and M.A. Birkett, 2018. Prospects for management of whitefly using plant semiochemicals, compared with related pests. *Pest Manage. Sci.*, 74: 2405–2411 DOI: 10.1002/ps.5058.
- Sharaby, A.M., F.M. El-Hawary and S.S. Moawad, 2013. Zinc Sulfate as a growth disruptor for *Spodoptera littoralis* with reference to histological changes in larval endocrine glands. *IOSR J. Agric. Vet. Sci.*, 5: 67–74
- Sharma, O. and O. Bambawale, 2008. Integrated management of key diseases of cotton and rice. In: *Integrated Management of Diseases Caused by Fungi, Phytoplasma and Bacteria*, pp: 271–302. Springer
- Spiegel-Roy, P. and E. Goldschmidt, 2008. *Biology of Citrus*, pp: 140–184. Cambridge University Press
- Steel, R.G.D., J.H. Torrie and D.A. Dickey, 1997. *Principles and Procedures of Statistics: A Biometrical Approach*, pp: 400–428, 3rd edition. McGraw Hill Book Co. Inc. New York
- Tsunoda, K., 2001. Preservative properties of vapor-boron treated wood and woodbased composites. *J. Wood Sci.*, 47: 149–153
- Wen, X., Y. Liu, S. Wang, Q. Wu, H. Pan, X. Yang, L. Guo and Y. Zhang, 2014. Sensitivity of *Bemisia Tabaci* (Hemiptera: Aleyrodidae) to several new insecticides in China: effects of insecticide type and whitefly species, strain and stage. *J. Insect Sci.*, 14: 261–263

(Received 06 August 2019; Accepted 19 August 2019)