



Full Length Article

Agronomic Efficiency and Economics of Crop Establishing Techniques and Nitrogen Application in Fine Aromatic Rice (*Oryza sativa* L.)

Shawaiz Iqbal^{1*}, Usama Bin Khalid¹, Muhammad Usman Saleem¹, Adila Iram¹, Naeem Ahmad¹, Nadeem Iqbal¹, Muhammad Sabar² and Tahir Hussain Awan¹

¹Department of Agronomy, Rice Research Institute, Kala Shah Kaku, Lahore-39018, Pakistan

²Department of Plant Breeding and Genetics, Rice Research Institute, Kala Shah Kaku, Lahore-39018, Pakistan

*For correspondence: shawaiziqbal@gmail.com

Abstract

Different crop establishing techniques and nitrogen application levels have crucial role in determining the production of rice. A field experiment was conducted at experimental farm of Rice Research Institute, Kala Shah Kaku, Pakistan during 2015 and 2016 to compare four planting methods *i.e.*, broadcast sowing (BS), drill sowing (DS), ridge sowing (RS) and puddled transplanted rice (PTR) along with four nitrogen rates *viz.*, N₀: control, N₁₃₃: 133 kg ha⁻¹, N₁₆₆: 166 kg ha⁻¹ and N₁₉₉: 199 kg ha⁻¹ using fine aromatic rice cultivar Super Basmati. Highest tiller density m⁻², panicle length, filled grain density per panicle, 1000 grain weight, grain yield, straw yield and harvest index were achieved for RS followed by DS while the minimum grain yield along with other yield contributing factors was recorded in PTR. Likely, highest filled grain density per panicle, grain yield, harvest index and minimum sterility percentage was recorded with 166 kg N ha⁻¹ compared to control. For the interaction highest grain yield and harvest index was obtained in RS for 166 kg N ha⁻¹ whereas minimum in PTR. Highest net profit and cost-benefit ratio was recorded with RS and 166 kg N ha⁻¹. Hence, RS with application of 166 kg N ha⁻¹ can be recommended to harvest the maximum grain yield and net profit in rice. © 2019 Friends Science Publishers

Keywords: Dry seeded rice; Transplanted rice; Nitrogen rates; Grain yield; Profit; Cost-benefit ratio

Introduction

More than half of the world's population consumes rice as a staple food and it is also the second largest cereal crop of Pakistan as well an important exportable national commodity. It accounts for 3.1 percent of value added in agriculture and 0.6 percent of GDP. During 2017-2018, rice crop was sown on an area of 2.72 million hectares whereas the production was 6.85 million tons. (Anonymous, 2018).

In Asian rice growing countries, transplanting rice seedling manually into puddled soil is most widely adopted method of rice planting but it consumes abundant quantity of water (Bouman and Tuong, 2001). As the water resources are declining, thus judicious use of irrigation water has gained a pivotal importance to meet the water and food needs of rapidly growing human population (Hanjar and Qureshi, 2010; Mahajan and Timsina, 2011; Mahajan *et al.*, 2012). The shrinking water resources are threatening the rice growers of Asia for adequate availability of irrigation water in the coming years (Mahajan *et al.*, 2012) that may adversely affect the sustainability of rice production in irrigated areas (Chauhan, 2012; Chauhan *et al.*, 2014). Thus, there is **dire** need to explore water saving approaches in farming and water saving of 11-18% and 40% in direct seeded rice as compared to transplanting in puddled soil has

been reported by Cabangon *et al.* (2002) and Tabbal *et al.* (2002), respectively. Furthermore, dry seeding on raised beds instead of puddled soil saves even more irrigation water (Bouman and Tuong, 2001). Moreover, applying irrigation precisely on the basis of soil surface tension (-20 KPa at 20 cm depth) in DSR, showed even higher saving of about 50% of water (Bhushan *et al.*, 2007; Sudhir *et al.*, 2007; Jat *et al.*, 2009). DSR as water saving technology, owing to its popularity is getting momentum over puddled transplanted rice (Farooq *et al.*, 2011).

Transplanting rice seedlings in puddled soil not only consumes more water but also involves intensive labour. It requires 25-50 **man-days** ha⁻¹ as compared to direct seeded rice which needs about 5 **man-days** ha⁻¹ (Dawe, 2005). Scarcity of skilled labour results in low plant population, eventually low crop productivity in PTR (Mann *et al.*, 2007; Gill *et al.*, 2014). Moreover, increasing wages result in an increase in cost of production, another factor contributing towards gradually replacing this conventional planting method with DSR (Bhushan *et al.*, 2007; Saharawat *et al.*, 2010). However, where labour is easily available and affordable, seedling transplanting could be chosen but on other hand, then it is more economical to opt for direct seeding (Laary *et al.*, 2012). Moreover, direct seeded rice plays pivotal role to achieve optimum plant population and

high-water use efficiency in water scarce areas that translates into improvement in the yield components and eventually in the final grain yield (0.16 t ha^{-1}) over PTR (Sarkar *et al.*, 2003; Farooq *et al.*, 2011). Furthermore, the DSR crop does not face the transplanting shock, that favors early crop establishment, accelerates the growth and hastens physiological maturity (Tuong, 2008) leading to shortening of crop duration (Ali *et al.*, 2006) by seven days (Rana *et al.*, 2014).

Therefore, sowing of rice crop by direct seeding is being practiced successfully in Europe, USA, Brazil, India, Philippines, Thailand, Malaysia and Sri Lanka (Farooq *et al.*, 2011; Kumar and Ladha, 2011; Weerakoon *et al.*, 2011).

Planting methods not only significantly affect yield and grain quality of rice crop but also influence health and productivity of soil. The yield of DSR can be increased by adopting suitable planting pattern which is considered an important agronomic factor (Iqbal, 2014; Sanjitha Rani and Jayakiran, 2010). Moreover, direct seeding had better plant establishment producing higher yield than transplanting (Laary *et al.*, 2012). However, various site-specific changes in method of land preparation and crop establishment may be applied in directly sown crop to enhance the crop yield (Farooq *et al.*, 2008). In addition, highest cost benefit ratio (1:1.91) had been recorded in DSR (Aslam *et al.*, 2008) along with about 20% reduction in production cost (Rana *et al.*, 2014).

Maximum paddy yield in drill sowing, broadcasting and bed sowing was achieved while lowest was found in puddled transplanted rice (Ali *et al.*, 2012) whereas Pasha *et al.* (2011) found that among different methods of direct rice seeding, highest paddy yield was achieved by drilling the dry seed. Moreover, in survey-based study of different rice growers in Pakistan, it was revealed that 0.70 to 0.89 t ha^{-1} higher yield obtained in DSR over conventional method of transplanting. The succeeding wheat crop also gave higher yield in field where previous rice crop was sown by DSR (Ali *et al.*, 2014). In a previous study found that mechanical transplanting, DSR by broadcast, bed sowing and drill sowing have superior results equally over farmers conventional transplanting (Ali *et al.*, 2012). In another study, DSR cultivation on ridges resulted in increased grain yield as compared to DSR by drill and broadcast sowing (Jamil *et al.*, 2017). Moreover, in China DSR ridge sowing also produced better results due to improved tillers, higher flag leaf chlorophyll content, slowed leaf aging, higher net photosynthetic rate, increased leaf area index, promotion of root development in upper soil at grain filling stage and finally resulting higher grain yield (Zhang *et al.*, 2003; Zhang *et al.*, 2009; Ting *et al.*, 2013).

The deficiency of nitrogen generally causes chlorosis in leaves leading to stunted growth. This is because of the poor formation of assimilates which ultimately results in early flowering and overall growth cycle of the plant is shortened, conversely, when nitrogen applied in excessive amount it accelerates vegetative growth of above ground parts of the plant due to enhanced chlorophyll content thus,

giving dark greenish colour and succulence to the crop, making crop more vulnerable to disease incidence, insect attack and severe lodging (Alam *et al.*, 2005). Generally, nitrogen uptake by the plant is merely 40% of the total nitrogen applied and this lower efficiency of the nitrogen fertilizer is due to its losses through denitrification, runoff, volatilization and leaching (Ladha *et al.*, 2005). In rice cultivation systems, nitrogen nutrition is the key factor which limits both the growth and quality of the crops (Khan *et al.*, 2012) as almost all agronomic parameters are affected with the changes of nitrogen input (Abou-Khalifa, 2010; Anisuzzaman *et al.*, 2010; Pramanik and Bera, 2013; Yoseftabar, 2013; Singh *et al.*, 2015) and grain yield is enhanced by altering nitrogen rate in different ecological zones (Sharma *et al.*, 2007; Ahmed *et al.*, 2015; Singh *et al.*, 2015). Moreover, super rice varieties show more positive response to the higher rates of nitrogen as compared to non-super rice varieties (Chen *et al.*, 2014). A slight increase in nitrogen dose ($22.5\text{--}30 \text{ kg ha}^{-1}$) is suggested in DSR which is only to meet the rice crop nitrogen requirements as it faces greater losses and lesser availability of nitrogen from the soil as mineralization occurs at early growth stage. In India, 25% more nitrogen is applied in DSR as compared to puddled rice (Varinderpal-Singh *et al.*, 2010; Mahajan and Timsina, 2011). Whereas, in Pakistan, nitrogen requirement of the rice crop cultivated as DSR seems relatively higher (55 kg N ha^{-1}) than the transplanted rice (Jamil *et al.*, 2017) in addition to the recommended dose.

Therefore, present study investigated response of fine aromatic rice Super Basmati variety with objectives to explore appropriate planting techniques in dry direct seeded rice and optimization of nitrogen application dose.

Materials and Methods

Experimental Site, Design and Treatments

The study was conducted at experimental area of Rice Research Institute, Kala Shah Kaku, Pakistan during 2015 and 2016. Randomized complete block design (RCBD) with split plot arrangement was used with three replications having net plot size of $4.5 \text{ m} \times 27 \text{ m}$. Four crop establishing methods *viz.*, broadcasting in well prepared dry soil (BS), drilling in well prepared dry soil (DS), ridges made after broadcasting of dry seed *i.e.*, ridge sowing (RS) and puddled transplanted rice (PTR) were compared. Four nitrogen rates *viz.*, N_0 : Control, N_{133} : 133 kg N ha^{-1} , N_{166} : 166 kg N ha^{-1} and N_{199} : 199 kg N ha^{-1} were applied. The physico-chemical characteristics of experimental soil are given in Table 1.

Crop sowing: Super Basmati was used as test variety. Dry seeding was done in first and second year on 14 June using rice seeding drill (DS), manual broadcasting (BS) and after manual broadcasting ridges were made by the ridger (RS). However, in PTR, seedlings transplanting was done on 18 July manually using thirty days old seedlings in both the years. Seed at the rate of 30 kg ha^{-1} in case of DSR method and 15 kg ha^{-1} by nursery raising for transplanting was used.

Weed control: Bispyribac sodium (Clover) 20% SC at the rate of 250 g ha⁻¹ was sprayed after 20 days of seeding in moist soil conditions and its application was repeated at 38 days after seeding as weeds again germinated in the field. Weed control in PTR was done by using Butachlor 60% EC at the rate of 2000 mL ha⁻¹ at 5 days after transplanting (DAT) in standing water.

Fertilizer management: Fertilizers phosphorus and potash at the rate of 85 and 62 kg ha⁻¹, respectively was used in the study. Full dose of phosphorus, potash, and 1/3rd nitrogen (N) were applied during last cultivation while remaining 2/3 was applied in two splits 35 and 55 days after transplanting or seeding.

Irrigation management: Drilling, ridge making and broadcasting of dry seed was done in a well-prepared dry seed bed following immediate flood irrigation and re-irrigated after 4 days to enhance germination and then irrigated with almost 4-5 days interval at early growth stage and 5-7 days at later stages of development. In case of PTR, continuous flooding of water was done for 30 days after transplanting (DAT) and then irrigated at about one-week interval. For insects and disease control, recommended plant protection measures were adopted to save the crop from loss. The crop was harvested at full physiological maturity and threshed manually.

Data Recording

Plant height of 05 randomly selected plants from each of the experimental unit was recorded from soil surface to the tip of the main panicle with the help of a meter rod and then averaged. The numbers of tillers was recorded at harvest from an area of 1 m² from three different places in each plot and averaged to calculate the tiller density m⁻². Five randomly selected panicles from each plot at harvest time were taken to determine the panicle length (cm), filled and unfilled grain density per panicle. 1000-grain weight of the normal kernels was recorded in grams using digital balance. After harvesting and threshing, the clean rough rice was air dried, bulked and weighed. Moisture content in grain was determined by using LKB-PRODUK TERAB-Sweden Grain Moisture Meter. The grain weight was adjusted at 14% moisture content and the yield of clean rough rice was expressed in tons ha⁻¹.

Sterility percentage, harvest Index (HI) and cost-benefit ratio (CBR) were determined by following equations:

$$\text{Sterility \%} = \frac{\text{unfilled spikelets per panicle}}{\text{total spikelets per panicle}} \times 100$$

$$\text{HI \%} = \frac{\text{Grain yield}}{\text{Total Biological Yield}} \times 100$$

$$\text{CBR} = \frac{\text{total income}}{\text{total cost of production}}$$

Agronomic efficiency for nitrogen was determined by following this equation:

$$\text{AE} = \frac{\text{GYNA} - \text{GYNO}}{\text{NRN}}$$

Where:

AE: Agronomic efficiency

GYNA: grain yield (kg ha⁻¹) with nitrogen addition

GYNO: grain yield (kg ha⁻¹) without nitrogen addition

NRN: Rate of nitrogen addition (kg ha⁻¹)

For planting methods agronomic efficiency was determined as follow

$$\text{AE} = \frac{\text{X Yield} - \text{PTR Yield}}{\text{PTR Yield}} \times 100$$

Where, X is the DSR by broadcast, drill or ridge sowing and PTR is puddled transplanted rice.

Statistical Analysis

The data collected was statistically analyzed using computer statistical package *STATISTIX 8.1*. Least significance difference (LSD) test at $P \leq 0.05$ was used to compare the treatment means.

Results

The cropping years had a significant effect on filled grain density per panicle, sterility percentage, 1000 grain weight, grain yield, straw yield, grand income, net profit and cost-benefit ratio and non-significant results observed for plant height, tiller density m⁻², panicle length and harvest index (Table 2). Therefore, the results are presented years wise. Different crop establishing techniques showed a significant effect ($P \leq 0.05$) on plant height in both years. The maximum height was recorded in DS that was similar with RS with an increment of 6 and 4%, 8 and 11% during the first and second year respectively (Table 4). The tiller density m⁻² was also affected significantly ($P \leq 0.05$) and 24, 20 and 15% more tillers were observed in RS, DS and BS, respectively over PTR and in second year increase of 22, 17 and 4% was observed. In case of panicle length (cm) 13, 5, 2% in 2015 and 14, 12, 5% in 2016 longer panicles were found in RS, DS and PTR than BS, respectively (Table 5). Number of filled grains per panicle (Table 4) was 19% and 13% more in RS and DS than PTR in 2015 and 22 and 9% in 2016. For the sterility percentage, BS, DS and RS recorded 26, 11 and 1.5% increase over PTR in 2015 with a similar trend observed in 2016. The thousand grain weight was non-significantly affected by planting techniques during both years (Table 5), however, 5% heaviest grains were achieved in RS than PTR during 2015 and 2% in 2016. Grain yield (t/ha⁻¹) varied significantly ($P \leq 0.05$) among different planting techniques in both years (Table 5) with maximum for RS that was similar with DS and BS indicating 20, 14 and 7% increase

Table 1: Physico-chemical properties of experimental field

Parameter	Soil depth	
	0-6 inch	6-12 inch
Texture	Clay loam	Clay loam
Organic matter (%)	0.39	0.25
Soil pH	8.31	8.14
EC (dS m ⁻¹)	1.42	0.89
SAR (m mol L ⁻¹) ^{1/2}	7.23	7.14
Saturation (%)	41	36
Nitrogen (%)	0.47	0.26
Available P (ppm)	5.5	5.2
Available K (ppm)	94	76

Table 2: Analysis of Variance of **cropping years, crop establishing techniques** and nitrogen application on different parameters of fine aromatic rice during 2015-16 (Total *df* = 95)

Sources	<i>df</i>	PH	TD	PL	FG	St	TGW	GY	SY	HI	GI	NP	BCR
Y	1	0.39 ^{NS}	10292 ^{NS}	55.21 ^{NS}	72.25 ^{**}	88.34 [*]	15.64 [*]	5.05 [*]	8.66 [*]	32.82 ^{NS}	328297 [*]	32830 ^{**}	1.36 [*]
PM	3	265.82 [*]	22216 ^{**}	46.35 ^{**}	1252.42 ^{**}	97.42 ^{**}	2.18 ^{**}	3.31 ^{**}	1.85 ^{NS}	92.04 ^{**}	210811 ^{**}	341957 ^{**}	1.36 ^{**}
Y×PM	3	14.28 ^{NS}	1329 ^{NS}	3.46 ^{NS}	29.69 [*]	8.39 [*]	0.50 [*]	0.32 ^{**}	0.64 ^{NS}	18.56 ^{NS}	20373 ^{**}	20372 ^{**}	0.05 ^{**}
NL	3	1619.69 ^{**}	137100 ^{**}	326.75 ^{**}	8560.73 ^{**}	546.91 ^{**}	25.63 [*]	19.88 ^{**}	38.38 ^{**}	251.68 ^{**}	1294226 ^{**}	1086675 ^{**}	3.73 ^{**}
Y×NL	3	23.14 ^{NS}	1312 ^{NS}	7.10 ^{NS}	61.68 ^{**}	3.41 ^{**}	0.83 ^{**}	0.32 ^{**}	1.20 [*]	0.86 ^{NS}	20696 ^{**}	20696 ^{**}	0.06 ^{**}
PM×NL	9	27.01 ^{**}	399 ^{NS}	9.98 ^{NS}	21.51 ^{**}	3.29 ^{**}	0.08 ^{NS}	0.16 ^{**}	0.28 ^{NS}	3.92 ^{NS}	10096 ^{**}	10095 ^{**}	0.03 ^{**}
Y×PM×NL	9	27.95 ^{**}	229 ^{NS}	3.62 ^{NS}	9.08 ^{NS}	3.13 ^{**}	0.10 ^{NS}	0.20 ^{**}	0.49 ^{NS}	4.27 ^{NS}	13067 ^{**}	13067 ^{**}	0.04 ^{**}

Y = Years; PM = Planting methods; NL = Nitrogen levels; PH = Plant height (cm); TD = Tiller density m⁻²; PL = Panicle length (cm); FG = Filled grain density per panicle; St = Sterility (%); TGW = 1000-grain weight (g); GY = Grain yield (t ha⁻¹); SY = Straw yield (t ha⁻¹); HI = Harvest Index (%); GI = Grand Income (\$ ha⁻¹); NP = Net profit (\$ ha⁻¹); BCR = Benefit-cost ratio; NS= Non Significant; *= Significant; **= highly significant

Table 3: Analysis of Variance of **crop establishing techniques** and nitrogen application on different parameters of fine aromatic rice during 2015 and 2016 (Total *df* = 47)

Sources	<i>df</i>	PH	TD	PL	FG	St	TGW	GY	SY	HI	GI	NP	BCR
2015													
PM	3	101.19 ^{**}	12268.70 [*]	20.37 ^{**}	533.31 ^{**}	35.10 ^{**}	2.33 ^{NS}	0.82 ^{**}	0.48 ^{NS}	28.40 [*]	51860 ^{**}	98106 ^{**}	0.45 ^{**}
NL	3	693.99 ^{**}	82207.50 [*]	142.65 ^{**}	4476.03 ^{**}	315.33 ^{**}	17.20 ^{**}	12.51 ^{**}	26.75 ^{**}	122.01 ^{**}	815629 ^{**}	698269 ^{**}	2.36 ^{**}
PM×NL	9	32.71 [*]	76.70 ^{NS}	9.42 ^{NS}	9.69 ^{**}	5.61 ^{**}	0.11 ^{NS}	0.18 [*]	0.58 ^{NS}	5.88 ^{NS}	11656 [*]	11659 [*]	0.04 ^{NS}
2016													
PM	3	178.91 ^{**}	11276.90 ^{**}	29.44 ^{NS}	748.80 ^{**}	70.71 ^{**}	0.35 ^{NS}	2.81 ^{**}	2.00 ^{**}	82.19 ^{**}	179324 ^{**}	264223 ^{**}	0.97 ^{**}
NL	3	948.83 ^{**}	56203.80 ^{**}	191.20 ^{**}	4146.39 ^{**}	234.99 ^{**}	9.27 ^{**}	7.69 ^{**}	13.46 ^{**}	130.53 ^{**}	499293 ^{**}	409102 ^{**}	1.43 ^{**}
PM×NL	9	22.24 ^{**}	551.70 ^{NS}	4.81 ^{NS}	20.91 ^{**}	0.81 ^{NS}	0.07 ^{NS}	0.18 ^{**}	0.19 ^{NS}	2.31 ^{NS}	11504 ^{**}	11503 ^{**}	0.04 ^{**}

PM = Planting methods; NL = Nitrogen levels; PH = Plant height (cm); TD = Tiller density m⁻²; PL = Panicle length (cm); FG = Filled grain density per panicle; St = Sterility (%); TGW = 1000-grain weight (g); GY = Grain yield (t ha⁻¹); SY = Straw yield (t ha⁻¹); HI = Harvest Index (%); GI = Grand Income (\$ ha⁻¹); NP = Net profit (\$ ha⁻¹); BCR = Benefit-cost ratio; NS= Non Significant; *= Significant; **= highly significant

Table 4: Effect of planting methods and nitrogen rates on agronomic and yield traits

Treatment/Parameters	Plant height (cm)		Tiller density m ⁻²		Panicle length (cm)		Filled grain density per panicle		Sterility (%)	
	2015	2016	2015	2016	2015	2016	2015	2016	2015	2016
BS	115.73	115.63	350.50	308.48	23.30	24.10	72.95	73.63	17.81	19.79
DS	119.83	119.40	365.00	348.02	24.44	27.02	81.13	80.63	15.67	18.91
RS	116.90	118.78	377.00	360.50	26.29	27.50	85.95	90.62	14.30	16.39
PTR	112.82	110.96	304.00	296.67	23.83	25.32	72.10	74.20	14.10	14.46
LSD (P≤ 0.05)	3.070	3.443	38.520	19.026	1.089	3.009	3.007	6.219	1.009	0.849
Nitrogen Levels										
N ₀	105.37	104.26	229.42	230.52	19.80	20.13	49.52	53.15	22.93	23.73
N ₁₃₃	117.02	114.70	359.25	332.38	24.01	26.73	85.23	82.17	14.39	16.65
N ₁₆₆	121.00	121.83	396.33	370.94	26.51	28.60	92.35	95.69	11.36	13.52
N ₁₉₉	121.90	123.98	411.50	379.83	27.55	28.47	85.03	88.06	13.19	15.64
LSD (P≤ 0.05)	2.942	2.228	19.276	29.055	1.994	2.284	2.421	2.391	0.614	1.707

BS: Broadcast sowing; DS: Drill sowing; RS: Ridge sowing; PTR: Puddled transplanted rice; N₀: control; N₁₃₃: 133 kg N ha⁻¹; N₁₆₆: 166 kg N ha⁻¹; N₁₉₉: 199 kg N ha⁻¹

over the PTR. Straw yield showed a non-significant response during both the years with a 6, 2 and 6% increase in RS, BS and DS respectively over PTR in first year and with similar trend were recorded in second year (Table 5). Different planting methods affected HI (%) significantly, with highest in RS and DS showing an increase of 9 and 8%

respectively than BS and PTR with lower HI (Table 5). Highest agronomic efficiency was of 20 and 51% observed in RS and followed by DS (14 and 35%) and BS (7 and 25%) during 2015 and 2016, respectively as compared to PTR (Table 5). Economic analysis for both the years (Table 4) indicated that the highest net profit and benefit-cost ratio

Table 5: Effect of planting methods and nitrogen rates on yield traits and agronomic efficiency (AE)

Treatment	1000-grain weight (g)		Grain yield (t ha ⁻¹)		Straw yield (t ha ⁻¹)		Harvest Index (%)		Agronomic efficiency	
	2015	2016	2015	2016	2015	2016	2015	2016	2015	2016
Planting methods										
BS	20.46	21.38	3.26	2.84	6.97	6.00	31.22	31.82	7.13	25.39
DS	20.37	21.37	3.46	3.05	6.69	6.12	33.61	32.92	13.99	34.77
RS	21.23	21.61	3.64	3.42	6.98	6.80	33.95	32.97	19.89	51.15
PTR	20.26	21.19	3.04	2.26	6.58	5.90	31.04	27.44	-	-
LSD (P≤0.05)	0.736	0.280	0.422	0.261	0.583	1.000	2.075	2.747	-	-
	Nitrogen levels									
N ₀	19.06	20.30	1.91	1.79	4.78	4.82	28.29	27.00	-	-
N ₁₃₃	20.21	21.17	3.38	2.84	6.66	6.00	33.69	32.04	11.09	8.85
N ₁₆₆	21.33	21.70	4.23	3.61	7.55	6.68	35.84	34.98	14.22	10.96
N ₁₉₉	21.72	22.38	3.88	3.33	8.23	7.31	32.00	31.13	10.68	7.74
LSD (P≤0.05)	0.431	0.258	0.308	0.147	0.544	0.483	1.364	1.214	-	-

BS: Broadcast sowing; DS: Drill sowing; RS: Ridge sowing; PTR: Puddled transplanted rice N₀: control; N₁₃₃: 133 kg N ha⁻¹; N₁₆₆: kg N ha⁻¹; N₁₉₉: 199 kg N ha⁻¹

Table 6: Interaction of planting methods and nitrogen levels on plant height (cm), tiller density m⁻², panicle length (cm), filled grain density per panicle and sterility percentage during 2015 and 2016

Treatment/parameters		Plant height (cm)		Tiller density m ⁻²		Panicle length (cm)		Filled grain density per panicle		Sterility (%)	
Planting methods	Nitrogen levels	2015	2016	2015	2016	2015	2016	2015	2016	2015	2016
BS	N ₀	98.87	103.53	235.67	229.83	16.73	17.60	47.87	50.87	26.14	26.49
	N ₁₃₃	118.73	113.40	355.00	290.25	24.13	26.40	78.40	77.00	16.83	19.31
	N ₁₆₆	121.60	122.27	397.33	352.75	25.67	27.20	85.40	85.93	13.41	16.13
	N ₁₉₉	123.73	123.33	414.00	361.08	26.67	25.20	80.13	80.70	14.86	17.24
DS	N ₀	111.27	105.73	239.67	249.00	22.00	22.93	50.73	52.60	25.40	24.82
	N ₁₃₃	118.20	119.93	382.00	357.00	22.43	27.00	89.33	84.60	13.16	17.97
	N ₁₆₆	124.33	125.40	411.67	387.67	26.07	28.73	96.00	97.47	11.00	15.62
	N ₁₉₉	125.53	126.53	426.67	398.42	27.27	29.40	88.47	87.87	13.14	17.22
RS	N ₀	105.67	104.07	252.33	247.92	23.73	21.07	56.53	62.07	19.84	22.40
	N ₁₃₃	120.53	115.53	386.33	385.58	24.53	27.33	94.60	90.00	14.57	15.90
	N ₁₆₆	121.53	126.13	426.33	402.00	27.97	30.73	100.47	108.87	10.57	12.21
	N ₁₉₉	119.87	129.40	443.00	406.50	28.93	30.87	92.20	101.53	12.21	15.06
PTR	N ₀	105.67	103.69	190.00	195.33	16.73	18.93	42.93	47.07	20.34	21.20
	N ₁₃₃	110.60	109.93	313.67	296.67	24.93	26.20	78.60	77.09	13.00	13.44
	N ₁₆₆	116.53	113.53	350.00	341.33	26.33	27.73	87.53	90.50	10.49	10.15
	N ₁₉₉	118.47	116.67	362.33	353.33	27.33	28.40	79.33	82.13	12.57	13.04
LSD (P≤0.05)		5.933	5.088	50.792	53.726	3.617	4.955	5.145	2.471	1.460	2.245

BS: Broadcast sowing; DS: Drill sowing; RS: Ridge sowing; PTR: Puddled transplanted rice N₀: control; N₁₃₃: 133 kg N ha⁻¹; N₁₆₆: kg N ha⁻¹; N₁₉₉: 199 kg N ha⁻¹

were found in RS which was followed by DS and BS, whereas, minimum were observed in PTR (Table 6).

Nitrogen (N) had a significant ($P \leq 0.05$) effect on plant height (cm), tiller density m⁻², panicle length (cm), filled grain number per panicle, sterility (%), 1000 grain weight (g), grain yield (t ha⁻¹), straw yield (t ha⁻¹) and HI (%) in both years (Table 3). In case of plant height (cm), the N₁₉₉ and N₁₆₆ applied expressed similar tallest plants and response was followed by N₁₃₃. These N levels expressed 16, 15, 11% (2015) and 19, 17, 10% (2016) more height over the N₀ treatment (Table 4). The production of tillers m⁻² increased with increasing N level. It was observed that N₁₉₉, N₁₆₆ and N₁₃₃ resulted in 80, 72, 57 (2015) and 65, 61, 44% (2016) more tiller density m⁻² than the plots without N (Table 4). Different N levels had significant ($P \leq 0.05$) impact on panicle length which increased by 39, 34 and 21% for N₁₉₉, N₁₆₆ and N₁₃₃ over control in 2015 and similar trend was observed during the second year of experimentation (Table 4). Different N rates showed a significant ($P \leq 0.05$) behavior on filled grain density per panicle (Table 4) and maximum number was observed in N₁₆₆. Nonetheless, N₁₉₉, N₁₆₆, N₁₃₃ resulted in 72, 87, 72 (2015) and 66 80, 55% (2016) more filled grain number

per panicle respectively than control. The sterility percentage significantly ($P \leq 0.05$) varied under both years (Table 4). In 2015 and 2016 42, 50, 37% and 43, 34, 30% less sterility was observed with N₁₉₉, N₁₆₆ and N₁₃₃ respectively over control. The 1000-grain weight was also significantly ($P \leq 0.05$) affected by different levels of N (Table 5) and increase in grain weight was found with increments of N doses. For N₁₉₉, N₁₆₆ and N₁₃₃, an increase of 14, 12, 6 and 10, 7, 4% was recorded in 2015 and 2016, respectively, over control. Grain yield (t ha⁻¹) is the most crucial and important parameter which also significantly ($P \leq 0.05$) affected by varying N doses (Table 5). It was observed that during 2015, an increase of 122% for N₁₆₆ over control. Moreover, 103 and 77% increase of grain yield was found for N₁₉₉ and N₁₃₃ over control treatment respectively. In addition, N₁₆₆ resulted in 25 and 9% more yield than N₁₃₃ and N₁₉₉. In second year, treatments N₁₆₆, N₁₉₉, N₁₃₃ produced 102, 86, 59% more yield than N₀, even N₁₆₆ better performed (27 and 8%) than N₁₉₉ and N₁₃₃. Straw yield (t ha⁻¹) was significantly ($P \leq 0.05$) altered by different N doses and maximum was produced by N₁₉₉ ($P \leq 0.05$), 72 and 51% more than the control treatment in 2015 and 2016, respectively (Table 5). Harvest index (%)

Table 7: Interaction of planting methods and nitrogen levels on 1000-grain weight (g), grain yield (t ha⁻¹), straw yield (t ha⁻¹) and harvest Index (%) during 2015 and 2016

Treatment/Parameters		1000-grain weight (g)		Grain yield (t ha ⁻¹)		Straw yield (t ha ⁻¹)		Harvest Index (%)	
Planting methods	Nitrogen levels	2015	2016	2015	2016	2015	2016	2015	2016
BS	N ₀	19.12	20.25	1.65	1.86	4.62	4.88	26.21	27.64
	N ₁₃₃	20.02	21.32	3.47	2.78	7.42	5.74	31.89	32.87
	N ₁₆₆	21.17	21.78	3.87	3.54	7.45	6.50	34.14	35.28
	N ₁₉₉	21.54	22.18	4.03	3.16	8.38	6.88	32.63	31.49
DS	N ₀	18.76	20.24	2.02	1.86	4.88	4.84	29.35	27.95
	N ₁₃₃	19.97	21.14	3.50	3.19	6.48	6.09	34.90	34.46
	N ₁₆₆	21.07	21.64	4.33	3.58	7.57	6.38	36.41	35.99
	N ₁₉₉	21.67	22.47	4.00	3.56	7.82	7.17	33.79	33.29
RS	N ₀	19.93	20.45	2.45	1.95	5.45	5.08	30.89	27.88
	N ₁₃₃	20.73	21.33	3.45	3.26	6.72	6.42	33.93	33.77
	N ₁₆₆	22.08	22.08	4.45	4.33	7.35	7.38	37.64	37.00
	N ₁₉₉	22.17	22.55	4.22	4.13	8.42	8.33	33.35	33.22
PTR	N ₀	18.44	20.25	1.52	1.48	4.18	4.54	26.72	24.53
	N ₁₃₃	20.11	20.87	3.12	2.13	6.02	5.75	34.04	27.07
	N ₁₆₆	21.00	21.31	4.27	2.98	7.85	6.46	35.17	31.67
	N ₁₉₉	21.49	22.32	3.25	2.46	8.28	6.85	28.23	26.51
LSD ($P \leq 0.05$)		1.045	0.526	0.424	0.364	0.881	1.300	3.133	3.447

BS: Broadcast sowing; DS: Drill sowing; RS: Ridge sowing; PTR: Puddled transplanted rice N₀: control; N₁₃₃: 133 kg N ha⁻¹; N₁₆₆: 166 kg N ha⁻¹; N₁₉₉: 199 kg N ha⁻¹

Table 8: Economics of different planting techniques and nitrogen levels and their impact on grand income, net profit and benefit cost ratio for 2015 and 2016

Treatments		Total cost (\$ ha ⁻¹)	2015			2016		
Planting techniques	Nitrogen levels		Grand Income (\$ ha ⁻¹)	Net profit (\$ ha ⁻¹)	Benefit-cost ratio	Grand Income (\$ ha ⁻¹)	Net profit (\$ ha ⁻¹)	Benefit-cost ratio
BS	N ₀	475.62	430.27	-45.35	-0.10	484.94	9.32	0.02
	N ₁₃₃	506.76	895.22	388.46	0.77	716.95	210.19	0.41
	N ₁₆₆	514.48	1036.18	521.70	1.01	909.36	394.88	0.77
	N ₁₉₉	522.21	999.78	477.57	0.91	816.47	294.26	0.56
DS	N ₀	496.28	522.97	26.69	0.05	484.12	-12.16	-0.02
	N ₁₃₃	527.42	899.96	372.54	0.71	820.59	293.18	0.56
	N ₁₆₆	535.14	1112.47	577.32	1.08	919.90	384.75	0.72
	N ₁₉₉	542.87	1030.09	487.23	0.90	916.94	374.07	0.69
RS	N ₀	488.68	633.48	144.80	0.30	507.07	18.39	0.04
	N ₁₃₃	519.82	888.36	368.54	0.71	838.87	319.05	0.61
	N ₁₆₆	527.54	1140.80	613.25	1.16	1111.23	583.69	1.11
	N ₁₉₉	535.27	1086.57	551.30	1.03	1065.25	529.98	0.99
PTR	N ₀	548.91	395.27	-153.64	-0.28	382.96	-165.95	-0.30
	N ₁₃₃	580.05	802.33	222.28	0.38	679.46	99.41	0.17
	N ₁₆₆	587.77	1096.89	509.12	0.87	770.54	182.76	0.31
	N ₁₉₉	595.50	1044.39	448.89	0.75	641.21	45.71	0.08
LSD ($P \leq 0.05$)		-	164.41	147.08	0.275	93.892	76.236	0.174

PKR is Pakistan's currency. US\$1=PKR 130. Paddy market price=250 \$ t⁻¹. Straw market price =3.85 \$ t⁻¹. Grand income= [(grain yield× paddy market price t⁻¹) + (Straw yield× straw market price t⁻¹). Net benefit= (grand income - total production cost). Benefit-cost ratio =Net profit/production cost

Abbreviations: DS= DSR-drill sowing, RS= DSR-ridge sowing, BS=DSR-broadcast, PTR = conventional transplanting, N₀=Control; N₁₃₃ = 133 kg N ha⁻¹; N₁₆₆= 166 kg N ha⁻¹; N₁₉₉=199 kg N ha⁻¹

was significantly ($P \leq 0.05$) affected by different levels of N in both years. In the first year, 27, 13 and 19%, more HI with N₁₆₆, N₁₃₃ and N₁₉₉ was achieved over control respectively. Similar response was observed during 2016 (Table 5). The highest agronomic efficiency (AE) was observed with the N application of N₁₆₆ in both the years. AE for N₁₃₃ and N₁₉₉ was 22% and 25% lower than N₁₆₆ in first year and 19 and 29% in second year (Table 5). Economic analysis revealed that maximum profit and cost-benefit ratio were achieved in N₁₆₆ but for control treatment income didn't even touch the total investments (Table 6).

In case of combined effect of both factors under investigation (planting method and N levels) significant ($P \leq 0.05$) behavior was observed regarding the parameters plant height, filled grain density per panicle, sterility

percentage, grain yield (t ha⁻¹), grand income and net profit. However, all other parameters were non-significant ($P \leq 0.05$) in the first year of study. In the second year, plant height, filled grain density per panicle, grain yield, grand income, net profit and benefit-cost ratio had also significant ($P \leq 0.05$) impact while other components remained non-significant ($P \leq 0.05$). During first year, maximum plant height (Table 6) and panicle length (Table 6) was recorded in DS×N₁₉₉ and minimum in DS×N₀. However, in case of number of filled grains per panicle (Table 6) and grain yield (Table 7), highest values were achieved in BS×N₁₆₆ and lowest were recorded for PTR×N₀. In case of sterility percentage (Table 6), lowest in BS×N₁₆₆ and maximum in DS×N₀ was found and HI% (Table 7) showed a contrary response to this. Number of tillers m⁻²

(Table 6), 1000-grain weight (Table 7) and straw yield (Table 7) were highest in BS×N₁₉₉ and lowest in PTR×N₀. On examining the interaction of both the factors under investigation, it becomes clear that RS × N₁₆₆ resulted in highest net profit and cost-benefit ratio and lowest was recorded in PTR × N₀ (Table 8).

Discussion

Different planting techniques had significant effect on rice growth and development. Maximum plant height in DS might be attributed to better soil condition and more root proliferation leading increased plant height. Jamil *et al.* (2017) reported more plant height in DSR ridge sowing than drill and broadcast methods. More tiller density per unit area in RS was owed to more surface area available with looser and porous soil, ultimately enabling plants to uptake nutrient and water more efficiently, to enhance tillering (Zhang *et al.*, 2003). Baloch *et al.* (2007) reported more tillers density in DSR ridge over parachute technology, PTR and DSR on flat. The effect of different planting methods was non-significant on panicle length as reported in DSR for broadcast and line sowing methods (Das *et al.*, 2015). More grain filling and less sterility in RS might be due to better vegetative growth on ridges that promoted higher assimilates contribution during the grain filling stage (Song *et al.*, 2009). Thousand grain weight was not much influenced by planting techniques; however, comparatively heavier grains were recorded in RS might be due to more water retention in ridges for nutrient transportation during physiological maturity (Yuan-zhi, 2015). Zhang *et al.* (2003) also reported more thousand grain weight in ridge cultivation over conventional rice cultivation. Grain yield and HI (%) are the most crucial components greatly influenced under varying plant establishment techniques and associated to other allometric traits like more plant population, higher filled grain number per panicle, 1000 grain weight and lowest sterility. This might be due to higher flag leaf chlorophyll content, slowed leaf aging, higher net photosynthetic rate (Ting *et al.*, 2013), higher root biomass and root vitality (Dan-Ying *et al.*, 2008), enhanced tillering, increased leaf area index and ultimately contributing to higher grain yield and HI (Feng *et al.*, 2010). Lower yield in PTR was seem to be suffered by subsurface plough pan hindering root development and eventually crop growth and grain yield. Yuan-zhi (2015) and Jamil *et al.* (2017) also reported higher grain yield in ridge cultivation over PTR, broadcast and drill sowing. Any of the crop establishment DSR methods were either more profitable as compared to PTR which is practiced by common farmers in Indo-Pakistan (Singh *et al.*, 2004; Ali *et al.*, 2012).

N is more volatile in nature and less stable in water, hence its efficiency is reduced due to volatilization and leaching (Ladha *et al.*, 2005; Xu *et al.*, 2013). Therefore, N fertilization management has a key role for the productivity enhancement of rice. Thus, optimum supply of N is

prerequisite need for enhancing farm yield. Higher dose of N probably promoted cellular activity giving more vegetative growth resulting in taller plants and more tillers. Manzoor *et al.* (2006); Pramanik and Bera (2013) reported 150 and 175 kg N ha⁻¹ optimum N levels above and below of that significant yield losses were observed. However, filled grain density per panicle increased up to a certain level above that fell down and it might be due to excessive vegetative growth because of accelerated metabolic activity offering more accumulation in shoots (Hu *et al.*, 2007; Huang *et al.*, 2008; Yoseftabar, 2013; Hirzel *et al.*, 2014). Increase in grain weight and grain yield with increments in N dose was observed up to a certain level after that it declined (Chen *et al.*, 2014; Singh *et al.*, 2015). Moreover, grain yield increases only up to particular level (Khan *et al.*, 2006; Manzoor *et al.*, 2006) above which more straw is produced (Yadi *et al.*, 2012). In case of HI, results of present study are in conformity with Feng *et al.* (2010) who reported alternate wetting and drying performed better than submerged conditions.

Conclusion

Among all crop establishment methods, RS with 166 kg N ha⁻¹ was the best combination producing maximum grain yield attributes, grain yield, and more economic returns under alkaline soil conditions and can be recommended for rice growers.

References

- Abou-Khalifa, A.A.B., 2012. Evaluation of some rice varieties under different nitrogen levels. *Adv. Appl. Sci. Res.*, 3: 1144–1149
- Ahmed, S., T.H. Awan, M. Salim and B.S. Chauhan, 2015. Economics of nitrogen and integrated weed management in dry seeded rice. *J. Anim. Plant Sci.*, 25: 1675–1684
- Alam, M.M., J.K. Ladha, K.S. Rahaman, H.R. Foyjunnessa, A.H. Khan and R.J. Buresh, 2005. Leaf color chart for managing nitrogen fertilizer in lowland rice in Bangladesh. *Agron. J.*, 97: 949–959
- Ali, A., O. Erenstein and D.B. Rahut, 2014. Impact of direct rice-sowing technology on rice producers' earnings: empirical evidence from Pakistan. *Dev. Stud. Res.*, 1: 244–254
- Ali, M.A., J.K. Ladha, J. Rickman and J.S. Lales, 2006. Comparison of different methods of rice establishment and nitrogen management strategies for lowland Rice. *J. Crop Improv.*, 16: 31–32
- Ali, R.I., N. Iqbal, M.U. Saleem and M. Akhtar, 2012. Effect of different planting methods on economic yield and grain quality of rice. *Intl. J. Agric. Appl. Sci.*, 4: 28–34
- Anisuzzaman, M., M. Salim and M.A.H. Khan, 2010. Effect of seed rate and nitrogen level on the yield of direct seeded hybrid boro rice. *J. Agro-For. Environ.*, 4: 77–80
- Anonymous, 2018. *Economic survey of Pakistan*. Govt. Pakistan Finance Division Economic Advisory Wing Islamabad
- Aslam, M., S. Hussain, M. Ramzan and M. Akhtar, 2008. Effect of different stand establishment techniques on rice yields and its attributes. *J. Anim. Plant Sci.*, 18: 80–82
- Baloch, M.S., I.U. Awan, G. Hassan and M. Zubair, 2007. Studies on plant population and stand establishment techniques for increasing productivity of rice in Dera Ismail Khan, Pakistan. *Rice Sci.*, 14: 118–124

- Bhushan, L., K. Jagdish, J.K. Ladha, R.K. Gupta, S. Singh, A. Tirol- Padre, Y.S. Saharawat, M. Gathala and H. Pathak, 2007. Saving of water and labor in a rice-wheat system with no-tillage and direct seeding technologies. *Agron. J.*, 99: 1288–1296
- Bouman, B.A.M. and T.P. Tuong, 2001. Field water management to save water and increase its productivity in irrigated lowland rice. *Agric. Water Manage.*, 49: 11–30
- Cabangon, R.J., T.P. Tuong and N.B. Abdullah, 2002. Comparing water input and water productivity of transplanted and direct seeded rice production systems. *Agric. Water Manage.*, 57: 11–31
- Chauhan, B.S., 2012. Weed ecology and weed management strategies for dry-seeded rice in Asia. *Weed Technol.*, 26: 1–13
- Chauhan, B.S., A.S.K. Abeysekara, M.S. Wickramaratne, S.D. Kalatunga and U.B. Wickrame, 2014. Effect of rice establishment methods on weedy rice (*Oryza sativa* L.) infestation and grain yield of cultivated rice (*Oryza sativa* L.) in Sri Lanka. *Crop Prot.*, 55: 42–49
- Chen, S., D. Wang, C. Xu, C. Ji and X. Zhang, 2014. Responses of super rice (*Oryza sativa* L.) to different planting methods for grain yield and nitrogen-use efficiency in the single cropping season. *Plos One*, 9: 1–9
- Dan-Ying, W., H. Bo, Z. Xiu-Fu, S. Guo-Sheng, X. Chun-Mei and F. Guan-Fu, 2008. Influence of rhizosphere oxygen concentration on rice root growth. *Acta Agron. Sin.*, 34: 803–808
- Das, G.C., S.C. Samanta, P. Biswas, N.K. Saha and J. Bhattacharya, 2015. Effects of sowing methods on yield attributes and yield of aus rice under the tidal ecosystem. *J. Biosci. Agric. Res.*, 4: 1–9
- Dawe, D., 2005. Increasing water productivity in rice-based systems in Asia-Past trends, current problems, and future prospects. *Plant Prod. Sci.*, 8: 221–230
- Farooq, M., K.H.M. Siddique, H. Rehman, T. Aziz, A. Wahid and D. Lee, 2011. Rice direct seeding experiences and challenges. *Soil Till. Res.*, 111: 87–98
- Farooq, M., S.M.A. Basra and S.A. Asad, 2008. Comparison of conventional puddling and dry tillage in rice-wheat system. *Paddy Water Environ.*, 6: 397–404
- Feng, Z., W. Dan-Ying, X. Chun-Mei, Z. Wei-Jian, L. Feng-Bo, M. Hai-Jun and Z. Xiu-Fu, 2010. Response of morphological, physiological and yield characteristics of rice (*Oryza sativa* L.) to different oxygen-increasing patterns in rhizosphere. *Acta Agron. Sin.*, 36: 303–312
- Gill, J.S., S.S. Walia and R.S. Gill, 2014. Direct seeded rice: An alternative rice establishment technique in north-west India—a review. *Intl. J. Adv. Res.*, 2: 375–386
- Hanjar, M.A. and M.E. Qureshi, 2010. Global water crisis and food security in an era of climate change. *Food Policy*, 35: 365–377
- Hirzel, J., A. Pedrero and K. Cordero, 2014. Effect of nitrogen rates and split nitrogen fertilization on grain yield and its components in flooded rice. *Chil. J. Agric. Res.*, 71: 437–444
- Hu, R.J. Cao, J. Huang, S. Peng, X. Zhong, Y. Zou, J. Yang and R.J. Buresh, 2007. Farmer participatory testing of standard and modified site-specific nitrogen management for irrigated rice in China. *Agric. Sci.*, 94: 331–340
- Huang, J., F. He, K. Cui, R.J. Buresh, B. Xu, W. Gong and S. Peng, 2008. Determination of optimal nitrogen rate for rice varieties using a chlorophyll meter. *Field Crops Res.*, 105: 70–80
- Iqbal, M.A., 2014. Productivity and quality of direct seeded rice under different types of mulches and planting patterns. *Amer.-Eur. J. Agric. Environ. Sci.*, 14: 1240–1247
- Jamil, M., S.S. Hussain, M.A. Qureshi, S.M. Mehdi, M.Q. Nawaz and Q. Javed, 2017. Rice yield improvement through various direct seeding techniques on moderately salt affected soil. *J. Anim. Plant Sci.*, 27: 848–854
- Jat, M.C., M.K. Gathala and J.K. Ladha, 2009. Evaluation of precision land leveling and double zero-till systems in rice-wheat rotations: water use, productivity, profitability and soil physical properties. *Soil Till. Res.*, 105: 112–121
- Khan, A.W., R.A. Mann, M. Saleem and A. Majeed, 2012. Comparative rice yield and economic advantage of foliar KNO₃ over soil applied K₂SO₄. *Pak. J. Agric. Sci.*, 49: 481–484
- Khan, U., B. Mishra, P. Pachauri and Y. Kumar, 2006. Effect of integrated nitrogen management on yield and nitrogen nutrition of irrigated rice (*Oryza sativa* L.). *Ind. J. Agric. Sci.*, 76: 176–180
- Kumar, V. and J.K. Ladha, 2011. Direct Seeding of Rice: Recent Developments and Future Research Needs. *Adv. Agron.*, 111: 297–413
- Laary, J.K., W. Dogbe, P.O. Boamah and J. Agawini, 2012. Evaluation of planting methods for growth and yield of “digang” rice (*Oryza sativa* L.) under upland condition of Bawku, upper east region, Ghana. *ARPN J. Agric. Bio. Sci.*, 7: 814–819
- Ladha, J.K., H.P. Pathak, T.J. Krupnik, J. Six and C.V. Kessel, 2005. Efficiency of fertilizer nitrogen in cereal production: retrospect and prospect. *Adv. Agron.*, 87: 86–156
- Mahajan, G., B.S. Chauhan, J. Timsina, P.P. Singh and K. Singh, 2012. Crop performance and water- and nitrogen-use efficiencies in dry-seeded rice in response to irrigation and fertilizer amounts in northwest India. *Field Crops Res.*, 134: 59–70
- Mahajan, G. and J. Timsina, 2011. Effect of nitrogen rates and weed control methods on weeds abundance and yield of direct-seeded rice. *Arch. Agron. Soil Sci.*, 57: 239–250
- Mann, R.A., S. Ahmad, G. Hassan and M.S. Baloch, 2007. Weed management in direct seeded rice crop. *Pak. J. Weed Sci. Res.*, 13: 219–226
- Manzoor, Z., T.H. Awan, M.A. Zahid and F.A. Faiz, 2006. Response of rice crop to different nitrogen levels. *J. Anim. Plant Sci.*, 16: 52–55
- Pasha, M.L., D. Bhadru, L. Krishna and R.B.M. Naik, 2011. Comparative performance of different rice planting methods under Nagarjuna Sagar Project Left Canal Command Area (Sri Lal Bahadur Shastri Canal) of Nalgonda district. *Crop Res.*, 44: 1–4
- Pramanik, K. and A.K. Bera, 2013. Effect of seedling age and nitrogen fertilizer on growth, chlorophyll content, yield and economics of hybrid rice (*Oryza sativa* L.). *Intl. J. Agron. Plant. Prod.*, 4: 3489–3499
- Rana, M.M., M.A. Al Mamun, A. Zahan, M.N. Ahmed and M.A.J. Mridha, 2014. Effect of planting methods on the yield and yield attributes of short duration Aman Rice. *Amer. J. Plant Sci.*, 5: 251–255
- Saharawat, Y.S., B. Singh, R.K. Malik, J.K. Ladha, M. Gathala, M.L. Jat, R.J. Sanjitha and K. Jayakaran, 2010. Evaluation of different planting techniques for economic feasibility in Rice. *Elec. J. Environ. Agric. Land Food. Chem.*, 9: 150–153
- Sanjitha Rani, T. and K. Jayakaran, 2010. Evaluation of different planting techniques for economic feasibility in Rice. *Electronic J. Environ. Agric. Food Chem. (EJEAFChE)*, 9: 150–153
- Sarkar, R.K.D., Sanjukta and S. Das, 2003. Yield of rainfed lowland rice with medium water depth under anaerobic direct seeding and transplanting. *Trop. Sci.*, 43: 192–198
- Sharma, R.P., S.K. Pathak and R.C. Singh, 2007. Effect of nitrogen and weed management in direct-seeded rice (*Oryza sativa* L.) under upland conditions. *Ind. J. Agron.*, 52: 114–119
- Singh, D.K., P.C. Pandey, Priyanker, A. Qureshi and S. Gupta, 2015. Nitrogen management strategies for direct seeded aerobic rice (*Oryza sativa* L.) grown in mollisols of Uttarakhand (India). *Intl. J. Appl. Pure Sci. Agric.*, 1: 130–139
- Singh, S., R.P. Tripathi, P. Shrama and R. Kumar, 2004. Effect of tillage on root growth, crop establishment and economics of rice. *Ind. J. Agric. Sci.*, 74: 300–304
- Song, C., C. Shenguan and C. Xin, 2009. Genotypic differences in growth and physiological responses to transplanting and direct seeding cultivation in rice. *Rice Sci.*, 16: 143–150
- Sudhir, Y., M.S. Gill and S.S. Kukal, 2007. Performance of direct seeded basmati rice in loamy sand in semi-arid subtropical India. *Soil Till. Res.*, 97: 229–238
- Tabbal, D.F., B.A.M. Bouman, S.I. Bhuiyan, E.B. Sibayan and M.A. Sattar, 2002. On-farm strategies for reducing water input in irrigated rice: case studies in the Philippines. *Agric. Water Manage.*, 56: 93–112
- Ting, H., L. Zhun, Y. Lin, H. Hui, L. Jian-xia, Z. Hua-bin, C. Yang and H. Huang, 2013. Effects of ridge-bed cultivation on physiological characteristics of flag leaf and grain yield during grain filling. *J. Human Agric. Uni. (Nat. Sci.)*, 1: 1–6

- Tuong, L., 2008. Studies on direct-seeding adaptability of Cambodian rice cultivars and development of cultivars with good eating quality. *PhD thesis*, Science of Plant and Animal Production, United Graduate School of Agricultural Science, Tokyo University of Agriculture and Technology, Japan
- Varinderpal-Singh, B.S., H.S.T. Yadvinder-Singh and R.K. Gupta, 2010. Need based nitrogen management using the chlorophyll meter and leaf colour chart in rice and wheat in South Asia: a review. *Nutr. Cycl. Agroecosyst.*, 88: 361–380
- Weerakoon, W.M., M.M. Mutunayake, C. Bandara, A.N. Rao, D.C. Bhandari and J.K. Ladha, 2011. Direct-seeded rice culture in Sri Lanka: lessons from farmers. *Field Crops Res.*, 121: 53–63
- Xu, J., L. Liao, J. Tan and X. Shao, 2013. Ammonia volatilization in gemmiparous and early seedling stages from direct seeding rice fields with different nitrogen management strategies: a pots experiment. *Soil Till. Res.*, 126: 169–176
- Yadi, R., S. Dastan and E. Yasari, 2012. Role of zinc fertilizer on grain yield and some qualities parameters in Iranian rice genotypes. *Ann. Biol. Res.*, 3: 4519–4527
- Yoseftabar, S., 2013. Effect of nitrogen and phosphorus fertilizer on spikelet structure and yield in rice (*Oryza sativa* L.). *Intl. J. Agric. Crop Sci.*, 5: 1204–1208
- Yuan-zhi, Y., 2015. Effects of ridge tillage on photosynthesis and root characters of rice. *Chilean J. Agric. Res.*, 35–41
- Zhang, R.P., H.Y. Dai, G.Z. Cai, X.J. Zhu and Z.Q. Hu, 2009. Correlation and path analysis on the yield characters of japonica giant embryo rice under different cultivation regimes. *J. Xichang College (Nat. Sci. Ed.)*, p. 8
- Zhang, X.F., D.Y. Wang and G.S. Shao, 2003. Effects of rice ridge cultivation on grain yield and quality and its physiological and ecological mechanisms. *Chin. J. Rice Sci.*, 4

(Received 24 April 2019; Accepted 15 July 2019)

IMPRESS