



Full Length Article

Effect of Nitrogen Concentration on the Growth and Fatty Acid Content of *Mortierella alpina*

Lingfei Li* and Na Jin

College of Food Science and Technology, Yunnan Agricultural University, Kunming 650201, China

*For correspondence: lingfeili@163.com

Received 21 April 2020; Accepted 07 May 2020; Published _____

Abstract

The effects of nitrogen sources and nitrogen concentration on the growth and fatty acid contents of the fungus *Mortierella alpina* were investigated in this study. The poorest nitrogen source (sodium nitrate) and the best nitrogen source (urea) from six nitrogen sources were selected and used for cultivation of *M. alpina* D36 in three nitrogen concentrations (0.05, 0.1 and 0.2 M) for 12 days. Dynamic changes in biomass (dry cell weight), yield, as well as fatty acid composition, proportion and content were determined during fermentation. The results showed that the growth of *M. alpina* was markedly influenced by nitrogen source and nitrogen concentration. With sodium nitrate as nitrogen source (low nitrogen), biomass was significantly higher than that from the medium- and the high-nitrogen concentrations, and the maximum biomass concentration was 6.42 g L⁻¹. However, with urea as nitrogen source, biomass concentration at the medium and high nitrogen concentrations were higher than that at low nitrogen concentration, and the maximum biomass concentration was 18.19 g L⁻¹. Besides, the effect of nitrogen concentration on fatty acid accumulation in *M. alpina* varied as a function of nitrogen source. Using sodium nitrate as nitrogen source, nitrogen concentration had very little effect on the content and yield of total fatty acids and polyunsaturated fatty acids (arachidonic acid and eicosapentaenoic acid). However, when urea was used as nitrogen source, the contents of fatty acids at low and medium nitrogen concentrations were significantly higher than those obtained at high nitrogen concentration. However, the fatty acid yield at low nitrogen concentration was not high due to lower biomass concentration. In conclusion, nitrogen source and nitrogen concentration greatly affect the growth and fatty acid accumulation in *M. alpina*, with medium urea concentration being the most conducive condition. © 2020 Friends Science Publishers

Keywords: Fatty acids; *Mortierella alpina*; Nitrogen concentration; Sodium nitrate; Urea

Introduction

Mortierella alpina, a fungus widely distributed in the soil, has a strong ability to produce lipids which account for up to 50% of its dry weight (Wang *et al.* 2013). At present, *M. alpina* is used commercially for the production of polyunsaturated fatty acids (PUFAs), especially arachidonic acid (Kikukawa *et al.* 2018). Besides, *M. alpina* produces many other polyunsaturated fatty acids, including linoleic acid, γ -linolenic acid, dihomo- γ -linolenic acid and eicosapentaenoic acid (Sakuradani 2010; Tang *et al.* 2018). These polyunsaturated fatty acids are very beneficial to human health, in that they are essential for normal physiological function (Wiktorowska-Owczarek *et al.* 2015; Madsen *et al.* 2019). Therefore, due to the oil-rich property and safety of *M. alpina* (Streekstra 1997; Nisha *et al.* 2009), it is considered a very promising producer of arachidonic acid, and it has attracted the attention of many researchers.

Although *M. alpina* has been used in the industrial production of arachidonic acid, optimization of fermentation conditions so as to increase yield is still the most direct and

effective way for arachidonic acid production (Gu *et al.* 2018; Zhang *et al.* 2019). It has been proposed that when microorganisms are in a suitable environmental conditions with sufficient nutrition, various cellular metabolisms are active and in a state of balance. At this stage, cell metabolism mainly serves to promote cellular growth, while the lipid content in the cell is maintained at a low level. However, changes in environmental conditions impair the metabolic balance in the cell, a situation which is beneficial for the accumulation of some metabolites (Ratray 1984). Many factors affect the production of arachidonic acid through *M. alpina* fermentation. These factors include carbon source, nitrogen source, amount of dissolved oxygen, pH, and temperature. Nitrogen is a necessary nutrient element for the growth of microorganisms. Nitrogen source has significant effects on the growth of oleaginous microorganisms and the accumulation of lipids (Baky *et al.* 2020; Feng *et al.* 2020). Nitrogen limitation is a common strategy for inducing lipid synthesis in lipid-producing microorganisms (Janssen *et al.* 2019; Tossavainen *et al.* 2019). It has been shown that the yield of mono-unsaturated

fatty acids in *Saccharomyces cerevisiae* was higher under nitrogen-limited conditions than that under non-nitrogen limited conditions (Tang and Chen 2014). Previous studies have revealed that oil-producing microorganisms begin to accumulate vast amounts of lipids only when the nitrogen source is exhausted and the carbon source is sufficient (Granger et al. 1993; Raimondi et al. 2014).

At present, research on the effect of nitrogen sources on lipid accumulation in *M. alpina* focus mainly on the selection of nitrogen sources and the optimization of fermentation conditions (Lu et al. 2011; Asadi et al. 2018; Lu et al. 2019). Studies have shown that the available nitrogen sources for *M. alpina* are inorganic and organic nitrogen sources, and the direction and yield of metabolites are influenced by different types of nitrogen sources (Stressler et al. 2013; Yu et al. 2018). A study on the effects of eight nitrogen sources on the characteristics of fatty acids in *M. alpina* showed that organic nitrogen sources were more favorable for cell growth and total lipid accumulation, than inorganic nitrogen sources, with urea being the most economical nitrogen source for industrial production of arachidonic acid (Lu et al. 2011). However, there are limited reports on the effect of nitrogen concentration on lipid accumulation in *M. alpina*. Moreover, different strains of *M. alpina* may present distinct demands for nitrogen. In this study, the effects of six nitrogen sources (ammonium chloride, ammonium nitrate, ammonium sulfate, sodium nitrate, potassium nitrate, and urea) on the growth and fatty acid accumulation of *M. alpina* were investigated. *Mortierella alpina* D36 was fermented with sodium nitrate and urea as nitrogen sources under different nitrogen concentrations (0.05, 0.1 and 0.2 M). Dynamic changes in biomass concentration, nitrogen residue, fatty acid content, and yields of mycelia were measured during the fermentation. This could provide a reference for the industrial production of polyunsaturated fatty acids using *M. alpina*.

Materials and Methods

Fungal strain

Mortierella alpina D36 was previously isolated from the soil by the Laboratory of Food Microbiology, Yunnan Agricultural University, and identified using morphological and ITS sequences. This fungal strain was maintained on potato dextrose agar (PDA) slants at 4°C.

Fermentation methods

The mycelia of *M. alpina* D36 were aseptically picked from the PDA slant medium and transferred to the inoculum medium comprised of glucose (30 g L⁻¹), yeast extract (6 g L⁻¹), KH₂PO₄ (3 g L⁻¹), NaNO₃ (3 g L⁻¹) and MgSO₄·7H₂O (0.5 g L⁻¹). The fungus was cultured with shaking at 175 rpm in Erlenmeyer flasks for 3 days at 20°C. Then, 10–15 mycelium pellets were transferred to the fermentation

medium containing glucose (50 g L⁻¹), KH₂PO₄ (3.8 g L⁻¹), MgSO₄·7H₂O (0.5 g L⁻¹), and a specific nitrogen source (any of ammonium chloride, ammonium nitrate, ammonium sulfate, sodium nitrate, potassium nitrate, and urea). The amount of each nitrogen source was calculated based on 0.1 M nitrogen. Four replicates were carried out with each nitrogen source. The fungus was cultured in Erlenmeyer flasks for 12 days at 20°C, with shaking at 175 rpm.

The formula of media

Different sodium nitrate concentrations were used to replace the nitrogen source in the above fermentation medium at concentrations of 4.25, 8.5 and 17 g L⁻¹. Similarly, the nitrogen source in the fermentation medium was replaced with urea at concentrations of 1.5, 3.0 and 6.0 g L⁻¹. The nitrogen concentrations in the medium were equivalent to 0.05 M, 0.1 M, and 0.2 M, respectively.

Determination of biomass concentration in medium

The mycelia from each fermentation medium were harvested using vacuum filtration, and washed three times with distilled water, followed by freeze-drying to a constant weight. Biomass concentration was expressed in terms of dry cell weight (DCW) per liter of fermentation medium.

Determination of nitrogen residue in media

The nitrogen content of the medium was determined using alkaline potassium persulfate digestion in combination with ultraviolet spectrophotometry (Sattayatewa et al. 2011).

Analysis of fatty acids

Dry mycelia (50 mg) were added to 1 mL of toluene, 2 mL of 1% (v/v) sulphuric acid in methanol, and 1 mL of heptadecanoic acid (1 mg mL⁻¹ in hexane). The mixture was placed in a 50°C-water bath overnight. Then, 5 mL of 5% (w/v) NaCl was added, followed by two extractions with hexane. The hexane layer was evaporated with N₂, and the residue was dissolved in 1 mL of hexane for GC-MS analysis (Agilent 7890A/5975C). The GC-MS conditions and methods used were in line with those reported in a previous report (Gu et al. 2018). The quantity of each fatty acid component was calculated from the peak area on the chromatogram using C_{17:0} as the internal standard.

Statistical analysis

Statistical analysis was performed with S.P.S.S. 19.0. Differences amongst multiple groups were determined using one-way analysis of variance (ANOVA), followed by Duncan's multiple range test. Values of *P* < 0.05 were taken as indicative of statistical significance.

Results

Effects of six nitrogen sources on biomass concentration of *M. alpina*

The nitrogen sources had significant effects on the biomass concentration of *M. alpina* (Fig. 1). The highest biomass concentration was 16.9 g L^{-1} when urea was used as the nitrogen source, followed by biomass concentration from use of ammonium nitrate (14.6 g L^{-1}). However, when the other four nitrogenous compounds were used as nitrogen sources, the biomass concentrations were all below 5 g L^{-1} , with sodium nitrate producing the lowest biomass concentration (2.6 g L^{-1}). Therefore, urea was the best nitrogen source for *M. alpina*, while sodium nitrate was the worst nitrogen source.

Effect of nitrogen concentration on biomass of *M. alpina*

With sodium nitrate or urea as the nitrogen source, the biomass concentration of *M. alpina* was gradually increased time-dependently (Fig. 2). However, when sodium nitrate was used as the nitrogen source, there were no significant differences in fungal biomass under different nitrogen concentrations on the third day of fermentation. As the fermentation proceeded, *M. alpina* grew fastest under low sodium nitrate concentration (Fig. 2A). Furthermore, the highest biomass was 6.42 g L^{-1} , which was significantly higher than those under medium and high sodium nitrate concentrations. There was no statistical difference in biomass concentration between the medium and high concentrations at each time point of the fermentation, except for the 12th day.

Using urea as nitrogen source, the biomass values under the medium concentration condition on the 3rd, 6th, and 9th days of fermentation were significantly higher than those under the low and high concentrations (Fig. 2B). However, on the 12th day of fermentation, biomass under high urea concentration condition surpassed that under medium concentration condition, and reached the highest value of 18.19 g L^{-1} . Moreover, at the same fermentation time, the biomass resulting from use of organic nitrogen urea as the nitrogen source was significantly higher than that resulting from the use of inorganic nitrogen sodium nitrate as the nitrogen source (Fig. 2A, Fig. 2B). At the 12th day of fermentation, biomass concentration with urea as the nitrogen source was 2.83 times higher than that produced with sodium nitrate as nitrogen source. Therefore, urea was more suitable for the growth of *M. alpina* than sodium nitrate (Fig. 2).

Residual nitrogen in fermentation media during the growth of *M. alpina*

In this study, nitrogen consumption by *M. alpina* depended on type of nitrogen source and nitrogen concentration in the




Fig. 1: Biomass concentration of *M. alpina* cultivated with different nitrogen sources. Data are expressed as mean \pm SE of four replicates. Data with different letters differ significantly ($P < 0.05$)




Fig. 2: Biomass concentration of *M. alpina* cultivated with different nitrogen concentrations during fermentation. (A) Sodium nitrate as nitrogen source; (B) urea as nitrogen source. Data are expressed as mean \pm SE of four replicates. Data with different letters differ significantly ($P < 0.05$)

media (Fig. 3). When sodium nitrate was the nitrogen source, nitrogen was slowly consumed by *M. alpina*. At the end of the fermentation, the nitrogen in the media with three concentrations of sodium nitrate (0.05, 0.1 and 0.2 M) was not depleted (Fig. 3A). Nitrogen consumption in the media with the low concentration of sodium nitrate was 0.035 M , which was 2.33 times higher than nitrogen consumption at the high concentration of sodium nitrate.

When urea was used as the nitrogen source to cultivate *M. alpina*, nitrogen content in the media decreased sharply with time (Fig. 3B). At the low urea concentration of 0.05 M, nitrogen in the media was depleted on the 9th day. At the medium urea concentration of 0.1 M, the nitrogen consumption was 0.093 M at the end of fermentation, whereas, at a high nitrogen concentration of 0.2 M, the nitrogen consumption was 0.14 M at the end of the fermentation. Correlation analyses showed that, irrespective of whether sodium nitrate or urea was used as the nitrogen source, the biomass concentration of *M. alpina* was significantly positively correlated with nitrogen consumption (Pearson correlation coefficient $r = 0.816$, $n = 48$, $P \leq 0.001$ for sodium nitrate; $r = 0.903$, $n = 48$, $P \leq 0.001$ for urea).

Effect of nitrogen concentration on the proportion of fatty acids in *M. alpina*

M. alpina D36 was cultured with different concentrations of nitrate or urea for 12 days. The fatty acid profiles of *M. alpina* cultured with different nitrogen concentrations are shown in Table 1. At different concentrations of the same nitrogen source, the fatty acid profiles did not differ much, except for the lack of lignoceric acid (C24:0) in the medium- and high-concentration sodium nitrate media. Monounsaturated fatty acid C24:1 was increased in mycelia when urea was used as the nitrogen source, when compared with sodium nitrate. Arachidonic acid was the most typical and the most abundant fatty acid in *M. alpina*.

With sodium nitrate as the nitrogen source, the level of arachidonic acid in total fatty acids fluctuated between 19.13 and 33.77%. At low sodium nitrate concentration, arachidonic acid level gradually decreased with fermentation time, but it was highest (27.54%) on the third day. Nevertheless, at the medium and high concentrations of sodium nitrate, the level of arachidonic acid first increased, and then decreased. Moreover, the arachidonic acid at the middle concentration of sodium nitrate reached a maximum level of 33.77% on the 6th day of fermentation. At the high concentration of sodium nitrate, the arachidonic acid level was only 19.55% on the 12th day of fermentation (Table 1).

With urea as nitrogen source, arachidonic acid level fluctuated between 22.02 and 34.32%. Unlike sodium nitrate, the levels of arachidonic acid at the low, medium, and high concentrations of urea slowly increased with fermentation time. Moreover, arachidonic acid level was higher at the low and medium concentrations of urea than at the high concentration of urea (Table 1).

The levels of saturated fatty acids, monounsaturated fatty acids and polyunsaturated fatty acids in total fatty acids are shown in Fig. 4 (A–C). In most cases, polyunsaturated fatty acids in *M. alpina* accounted for more than 50% of total fatty acids, and even exceeded 60% at low urea concentration. Overall, nitrogen source and nitrogen concentration had no significant effects on the proportion of polyunsaturated fatty acids (Fig. 4C).




Fig. 3: Residual nitrogen in the fermentation media during the growth of *M. alpina*. (A) Sodium nitrate as the nitrogen source; (B) urea as the nitrogen source. Data with different letters differ significantly ($P < 0.05$)

Effect of nitrogen concentration on the fatty acid contents in *M. alpina*

Fatty acids contents in mycelia *M. alpina* cultivated under different nitrogen concentrations for 12 days are shown in Table 2. In most cases, the total fatty acid content (Fig. 5A) and the content of each fatty acid component (Table 2) increased gradually with increase in fermentation time. When sodium nitrate was used as the nitrogen source, nitrogen concentration had little effect on the levels of total fatty acids, polyunsaturated fatty acids, arachidonic acid, and eicosapentaenoic acid. However, when urea was used as nitrogen source, the contents of fatty acids at the low and medium nitrogen concentrations were significantly higher than those at the high concentration (Fig. 5A–D). Moreover, the fatty acid contents were significantly higher when urea was used as nitrogen source than when sodium nitrate was used as nitrogen source. Therefore, urea was more suitable for fatty acid production by *M. alpina* than sodium nitrate.

Effect of nitrogen concentration on the fatty acid yields in *M. alpina*

The effects of nitrogen source and nitrogen concentration on yields of fatty acids were significant (Fig. 6A–D). At the same nitrogen concentration, the yields of total fatty acids,




Fig. 4: Proportion of fatty acids in total fatty acids of *M. alpina* D36 cultivated at different nitrogen concentrations during fermentation. (A) Proportion of saturated fatty acids in total fatty acids; (B) Proportion of monounsaturated fatty acids in total fatty acids; (C) Proportion of polyunsaturated fatty acids in total fatty acids. Data with different letters differ significantly ($P < 0.05$)

polyunsaturated fatty acids, arachidonic acid, and eicosapentaenoic acid, increased gradually as the fermentation continued. Compared with sodium nitrate, the yield of fatty acids using urea as nitrogen source was significantly higher, mainly due to the much higher biomass concentration of *M. alpina* when urea was used as nitrogen source. With sodium nitrate as nitrogen source, the yield of fatty acids decreased with increasing nitrogen concentration, which was consistent with the trend in biomass concentration. The yield of arachidonic acid was 190.87 mg L⁻¹ at low sodium nitrate concentration on the 12th day, which was 1.78 times higher than the corresponding yield at medium concentration, and 3.97 times higher than the yield at high concentration.

With urea as the nitrogen source, the yield of fatty acids was highest at the medium nitrogen concentration. On the 12th day, the yields of polyunsaturated fatty acids and arachidonic acid at the medium urea concentration were 1866.66 and 1090.97 mg L⁻¹, respectively, which were 1.53 and 1.66 times higher than the corresponding yields at low urea concentration, respectively. Moreover, the low and medium urea concentrations were conducive for the accumulation of eicosapentaenoic, another essential polyunsaturated fatty acid (Fig. 6D).

Discussion

Nitrogen sources are essential for microbial growth, but not all nitrogen sources are suitable for the growth and oil accumulation of *M. alpina*. This study compared the effects of six nitrogen sources on the growth of *M. alpina* D36, and found that the organic nitrogen urea was the most beneficial for the growth of mycelia, followed by an inorganic nitrogen i.e. ammonium nitrate, while the most unfavorable nitrogen source was sodium nitrate. Moreover, urea was more conducive to accumulation of fatty acids in *M. alpina* than sodium nitrate. These results are consistent with the findings of other researchers who reported that organic nitrogen sources were more conducive for *M. alpina* growth and oil accumulation than inorganic nitrogen sources (Lu *et al.* 2011; Nisha and Venkateswaran 2011). Singh and Ward (1997) used 1% (w/v) corn steep instead of sodium nitrate, for cultivation of *M. alpina*, and found that the production of arachidonic acid improved significantly. Similar results were found in other organisms. Certik *et al.* (1999) compared lipogenesis and activities of lipogenic enzymes in the fungus *Cunninghamella echinulata* as a function of different inorganic and organic nitrogen sources, and found that organic nitrogen enhanced lipid accumulation. A study

Table 1: Levels of fatty acids of *M. alpina* D36 grown under different nitrogen concentrations during fermentation

Nitrogen source	Nitrogen concentration	Fermentation time (days)	Saturated fatty acids (%)					Monounsaturated fatty acids (%)			Polyunsaturated fatty acids (%)						
			C _{14:0}	C _{16:0}	C _{18:0}	C _{22:0}	C _{24:0}	C _{18:1}	C _{20:1}	C _{24:1}	LA (C _{18:2})	C _{20:2}	GLA (C _{18:3})	DGLA (C _{20:3})	ARA (C _{20:4})	EPA (C _{20:5})	
Sodium nitrate	Low	3	1.09± 0.05bcd	15.83± 0.37bc	8.74± 0.25de	0.81± 0.27d	0.63± 0.12c	18.28± 0.31bcd	0.63± 0.21ab	-	5.65± 0.08b	0.80± 0.33c	6.98± 0.16b	8.66± 0.03bc	27.54± 0.52bc	4.37± 0.22bc	
		6	1.00± 0.08cde	16.24± 0.19bc	10.07± 0.14cde	1.33± 0.05bc	0.75± 0.07b	15.52± 0.22cde	0.75± 0.04ab	-	4.95± 0.10cd	1.14± 0.06bc	5.79± 0.23bcd	8.28± 0.26bc	26.96± 0.48bc	7.23± 0.65a	
		9	0.93± 0.05de	16.82± 0.73abc	10.79± 0.25cd	1.51± 0.05bc	0.79± 0.04b	17.04± 0.83cde	0.73± 0.13ab	-	4.35± 0.09efg	1.09± 0.12bc	5.13± 0.12cdef	7.95± 0.54c	26.03± 1.57bcd	6.84± 0.39a	
	Medium	3	1.04± 0.02bcde	16.90± 0.60abc	10.69± 0.26bcd	1.45± 0.04bc	0.98± 0.06a	18.24± 1.03bcd	0.58± 0.05b	-	4.36± ±0.22efg	0.92± 0.06c	5.52± 0.37cde	8.38± 0.45bc	23.75± 1.82de	7.19± 0.62a	
		6	0.80± 0.07e	16.18± 1.00bc	10.05± 0.82cde	1.50± 0.12bc	-	12.39± 0.72cde	0.71± 0.02b	-	4.44± 0.19bc	1.25± 0.28bc	4.45± 0.28def	9.92± 0.51abc	33.77± 1.26b	4.52± 0.69c	
		9	1.06± 0.07bcd	18.71± 1.24ab	12.19± 0.84ab	2.11± 0.19a	-	14.81± 0.86ef	0.82± 0.05ab	-	3.90± 0.22fgh	1.40± 0.15abc	4.15± 0.44ef	8.80± 0.89bc	27.80± 1.74def	4.25± 0.63abc	
	High	3	1.18± 0.07bc	19.48± 1.01a	12.19± 1.07ab	2.19± 0.11a	-	18.69± 1.07bc	0.94± 0.06a	-	3.69± 0.13h	1.80± 0.23a	4.04± 0.39f	8.77± 0.62bc	21.72± 1.47def	5.28± 0.63abc	
		6	1.93± 0.08a	18.89± 0.60ab	8.32± 0.46e	0.32± 0.07e	-	25.49± 0.50a	0.09± 0.01c	-	6.80± 0.28a	0.13± 0.02d	9.46± 1.18a	6.14± 0.48d	19.13± 0.24f	5.72± 0.92c	
		9	1.16± 0.01bcd	16.31± 0.61bc	9.54± 0.62cde	1.04± 0.14cd	-	17.99± 1.07bcde	0.71± 0.18ab	-	5.86± 0.373b	1.39± 0.29abc	6.41± 0.46bc	9.00± 0.55bc	25.37± 1.21bcd	5.24± 0.63abc	
	Urea	Low	3	1.21± 0.11bc	15.32± 0.94c	11.59± 0.83ab	1.45± 1.6bc	-	14.61± 0.91def	0.79± 0.13ab	-	4.96± 0.10de	1.29± 0.05abc	4.47± 0.03def	11.39± 0.45a	25.98± 2.08bcd	7.09± 0.72a
			6	1.26± 0.09b	18.42± 1.04abc	13.11± 0.41a	1.76± 0.13ab	-	21.00± 2.63b	0.71± 0.04ab	-	3.86± 0.16gh	1.54± 0.18abc	3.95± 0.73f	9.12± 1.04bc	19.55± 2.28ef	5.26± 0.50ab
			9	0.66± 0.01e	11.64± 0.21a	6.37± 0.25b	1.35± 0.04f	3.37± 0.19e	12.10± 17.56±	0.49± 0.40±	1.46± 1.46±	3.11± 5.31±	1.27± 0.55±	4.78± 7.85±	8.87± 8.77±	32.55± 22.02±	11.98± 7.88±
Medium		3	0.49± 0.02f	13.66± 0.14b	6.16± 0.16gh	1.45± 0.04c	4.04± 0.13b	12.25± 0.33de	0.49± 0.01bcd	1.34± 0.06abc	3.56± 0.13cd	1.38± 0.08bc	4.27± 0.03h	7.82± 0.17d	32.46± 0.96a	10.64± 0.39ab	
		6	1.55± 0.07b	16.19± 0.20a	9.21± 0.19a	1.26± 0.06de	1.99± 0.11e	20.39± 0.56a	0.84± 0.01a	5.00± 0.08e	1.00± 0.07d	8.18± 0.23ab	9.09± 0.13bc	20.83± 1.40c	3.90± 0.42ef		
		9	0.88± 0.04c	13.60± 0.38b	7.77± 0.16cd	1.58± 0.01b	4.03± 0.16b	15.92± 0.58bc	0.73± 0.01a	3.90± 0.06e	1.22± 0.03c	5.94± 0.13de	9.22± 0.13de	31.17± 0.03ab	3.84± 0.89ab		
High		3	0.83± 0.02cd	12.95± 0.40bc	6.94± 0.03e	1.61± 0.03b	4.44± 0.14ab	14.29± 0.31cde	0.54± 0.02abc	0.89± 0.05e	3.72± 0.07c	1.33± 0.10bc	5.34± 0.06f	8.87± 0.08bc	32.75± 1.12a	5.53± 0.44de	
		6	0.69± 0.06de	13.40± 0.37b	5.85± 0.13h	1.41± 0.02c	3.40± 0.77c	14.42± 0.20cd	0.47± 0.01cde	1.20± 0.07de	3.52± 0.14de	1.45± 0.06ab	4.65± 0.10gh	7.89± 0.03d	34.32± 0.52a	6.51± 0.84cd	
		9	1.72± 0.06a	15.83± 0.28a	8.52± 0.22b	1.21± 0.02ef	1.73± 0.06e	19.84± 0.58a	0.40± 0.03e	1.41± 0.06ab	5.52± 0.10a	0.61± 0.13e	8.46± 0.26a	7.77± 0.53d	20.67± 0.68c	6.29± 0.90cd	
High		3	0.93± 0.06c	12.62± 0.25bcd	8.33± 0.09b	1.42± 0.04c	2.59± 0.12d	17.55± 0.70b	0.56± 0.03ab	1.23± 0.03bcd	3.64± 0.11d	1.23± 0.07bc	6.56± 0.16c	9.70± 0.15a	28.37± 0.97b	5.26± 0.76def	
		6	0.92± 0.04c	12.67± 0.29bcd	7.44± 0.12d	1.59± 0.03b	3.51± 0.06c	20.41± 0.56a	0.57± 0.02a	1.11± 0.04d	3.56± 0.05cd	1.58± 0.02a	5.74± 0.07ef	8.86± 0.12bc	28.40± 0.83b	3.47± 0.28f	
		12	0.65± 0.02e	11.83± 0.65d	6.71± 0.24ef	1.72± 0.07a	4.82± 0.23a	20.22± 2.02a	0.61± 0.01a	0.82± 0.10e	3.97± 0.15c	1.65± 0.02a	6.00± 0.25de	8.57± 0.19c	28.33± 2.49b	4.10± 0.73ef	

^{a-c} Indicates 'not detected'. Data are expressed as mean ± SE of four replicates. Values with different letters differ significantly ($P < 0.05$)

on the effects of sodium nitrite, sodium nitrate, urea, and ammonium chloride on the growth and lipid accumulation of marine algae *Desmodesmus* spp. WC08 showed that the use of urea produced the best growth and lipid accumulation of *Desmodesmus* spp. WC08 (Luo et al. 2016).

In the present study, nitrogen concentration had little effect on the proportions of fatty acids in *M. alpina*. However, it significantly affected the contents and yields of total fatty acids, polyunsaturated fatty acids, arachidonic acid, and eicosapentaenoic acid. When urea was used as nitrogen source, the contents and yields of total fatty acids, polyunsaturated fatty acids, and arachidonic acid at the low and medium nitrogen concentrations were higher than those at the high nitrogen concentration. These findings are consistent with the results obtained in a previous study in

which the effects of low and high nitrogen concentrations on the growth and lipid accumulation of two oleaginous microalgae were investigated (Wu et al. 2015). The results showed that the levels of total lipids, neutral lipids, and total fatty acids produced by these two microalgae at low nitrogen concentration were significantly higher than those produced at the high nitrogen concentration. In the present study, using sodium nitrate as the sole nitrogen source, fungal biomass and total fatty acids, polyunsaturated fatty acids, and arachidonic acid yields decreased with increasing nitrogen concentration, indicating that a high concentration of nitrate could inhibit the growth and fatty acid accumulation of *M. alpina*. Using urea as nitrogen source, the nitrogen was completely consumed or nearly consumed at the low and medium nitrogen concentrations on the 9th

Table 2: Contents of fatty acids in *M. alpina* D36 grown at different nitrogen concentrations during fermentation

Nitrogen source	Nitrogen concentration	Fermentation time	Saturated fatty acids (mg g ⁻¹)					Monounsaturated fatty acids (mg g ⁻¹)			Polyunsaturated fatty acids (mg g ⁻¹)					
			C _{14:0}	C _{16:0}	C _{18:0}	C _{22:0}	C _{24:0}	C _{18:1}	C _{20:1}	C _{24:1}	LA (C _{18:2})	C _{20:2}	GLA (C _{18:3})	DGLA (C _{20:3})	ARA (C _{20:4})	EPA (C _{20:5})
Sodium nitrate	Low	3	0.81±	12.05±	6.54±	0.61±	0.26±	13.17±	0.47±	-	4.23±	0.62±	5.24±	6.51±	20.78±	3.31±
		6	0.04c	0.79cd	0.44d	0.21c	0.02c	0.99bc	0.16def	-	0.31b	0.29e	0.42bcd	0.53c	2.04bc	0.38efg
		9	1.11±	18.20±	11.29±	1.49±	0.85±	17.39±	0.84±	-	5.55±	1.27±	6.50±	9.27±	30.21±	8.12±
		12	0.07bc	0.25abcd	0.18abc	0.06b	0.09b	0.28abc	0.06bc	-	0.05a	0.06bc	0.33bc	0.27abc	0.43ab	0.80ab
	Medium	3	1.10±	19.56±	12.65±	1.77±	0.92±	19.78±	0.82±	-	5.15±	1.31±	6.06±	9.55±	31.23±	8.01±
		6	0.14bc	1.33abcd	1.15abc	0.15b	0.07b	1.24abc	0.05bcd	-	0.58ab	0.22bc	0.68bcd	1.47abc	4.83ab	0.86ab
		9	1.32±	21.47±	13.61±	1.83±	1.25±	23.30±	0.73±	-	5.50±	1.17±	7.00±	10.65±	30.30±	9.18±
		12	0.10abc	1.50abc	1.01ab	0.07b	0.07a	2.48ab	0.06bcde	-	0.18a	0.09bcd	0.37ab	0.80ab	2.46ab	1.08a
	High	3	0.58±	10.34±	6.11±	0.83±	-	9.77±	0.33±	-	3.52±	0.78±	2.95±	6.52±	19.09±	2.46±
		6	0.03c	0.70d	0.41d	0.08c	-	0.71c	0.01f	-	0.37b	0.14de	0.36d	0.64c	2.08bc	0.36g
		9	1.04±	20.44±	12.73±	1.70±	-	15.58±	0.85±	-	5.56±	1.40±	5.31±	11.75±	41.08±	5.06±
		12	0.10bc	1.60abc	1.36abc	0.07b	-	1.25bc	0.12bc	-	0.13a	0.05bc	0.23bcd	0.30a	2.82a	0.62cdef
Urea	Low	3	1.36±	24.40±	15.75±	2.73±	-	19.17±	1.06±	-	5.06±	1.83±	5.41±	11.49±	36.30±	5.61±
		6	0.04abc	1.24ab	0.74a	0.22a	-	0.92abc	0.09a	-	0.33ab	0.19b	0.63bcd	1.35a	3.52a	1.69bcde
		9	1.63±	27.02±	14.24±	3.03±	-	26.16±	1.30±	-	5.17±	2.54±	5.75±	12.39±	30.42±	7.51±
		12	0.04a	1.70a	1.83ab	0.11a	-	2.47a	0.07a	-	0.48ab	0.39a	0.93bcd	1.67a	3.66ab	1.35abc
	Medium	3	1.33±	12.64±	5.88±	0.43±	-	17.43±	0.45±	-	4.01±	0.91±	7.18±	4.52±	13.76±	2.86±
		6	0.10abc	1.07cd	0.81d	0.05c	-	1.82abc	0.10def	-	0.55b	0.22cde	1.07a	1.07c	1.39c	0.29g
		9	1.14±	16.03±	9.38±	1.02±	-	17.69±	0.69±	-	5.77±	1.35±	6.34±	8.86±	25.09±	5.20±
		12	0.04bc	0.46bcd	0.49bcd	0.11bc	-	0.91abc	0.13cde	-	0.31a	0.21bc	0.56bc	0.57abc	1.81bc	0.71cdef
	High	3	1.38±	17.61±	13.69±	1.72±	-	17.18±	0.92±	-	5.39±	1.49±	5.18±	12.29±	27.62±	7.34±
		6	0.27abc	1.61bcd	1.60ab	0.23b	-	1.85abc	0.12bc	-	0.23ab	0.16bc	0.49bcd	0.93a	2.51bc	0.38abcd
		9	1.14±	16.63±	11.91±	1.58±	-	18.69±	0.66±	-	3.55±	1.40±	3.71±	8.52±	18.29±	5.20±
		12	0.09bc	1.29bcd	1.05abc	0.12b	-	1.36abc	0.09cde	-	0.43b	0.20bc	0.80d	1.48abc	3.39c	0.59cdef
Urea	Low	3	1.21±	11.86±	6.53±	0.87±	1.51±	13.62±	0.31±	1.14±	4.08±	0.43±	6.06±	6.84±	17.12±	6.24±
		6	0.07cd	1.18de	0.69f	0.12g	0.32de	1.22gh	0.05cd	-	0.13def	0.28bcd	0.28e	0.45fg	0.24e	1.75f
		9	1.12±	15.08±	9.82±	1.60±	3.17±	16.41±	0.84±	1.61±	7.75±	1.05±	7.69±	11.72±	33.92±	11.87±
		12	0.04cde	0.64cd	0.40de	0.10f	0.24d	2.35fg	0.32ab	0.50cd	0.49a	0.15cd	0.21cde	0.45cd	1.76de	1.01bc
	Medium	3	1.21±	21.54±	11.77±	2.48±	6.22±	22.36±	0.90±	2.71±	5.75±	2.35±	8.82±	16.36±	60.01±	22.19±
		6	0.03cd	1.16b	0.63ab	0.10bc	0.21bc	1.56cde	0.04a	0.15a	0.28abcd	0.14b	0.31abc	0.10a	2.51b	1.41a
		9	1.04±	29.06±	13.10±	3.08±	8.57±	26.05±	1.04±	2.84±	7.06±	2.91±	9.10±	16.64±	69.15±	22.66±
		12	0.03efg	1.28a	0.69a	0.17a	0.39a	1.28bc	0.07a	0.08a	0.51ab	0.10a	0.42ab	0.77a	4.10a	1.42a
	High	3	0.93±	9.82±	5.59±	0.77±	1.21±	12.39±	0.34±	0.51±	3.02±	0.61±	4.95±	5.52±	12.66±	2.36±
		6	0.02fgh	0.48ef	0.29fg	0.07g	0.10g	0.83gh	0.02cd	0.05g	0.12cd	0.07de	0.17gh	0.29ef	1.15f	0.27f
		9	1.10±	17.03±	9.73±	1.98±	5.08±	19.92±	0.71±	0.92±	4.88±	1.53±	7.44±	11.57±	39.22±	4.38±
		12	0.05def	0.52c	0.30de	0.09de	0.39c	0.60def	0.02ab	0.09efg	0.16abcd	0.09c	0.20de	0.51cd	2.49cd	0.53ef
Urea	Low	3	1.44±	22.52±	12.01±	2.78±	7.69±	24.80±	0.93±	1.53±	6.45±	2.33±	9.25±	15.36±	56.29±	9.65±
		6	0.10a	2.07b	0.83ab	0.21ab	0.55ab	2.06bcd	0.10a	0.12cd	0.52abc	0.34b	0.69a	1.05ab	1.71b	1.21cd
		9	1.42±	28.11±	12.17±	2.95±	7.08±	30.13±	0.98±	2.49±	7.30±	3.03±	9.68±	16.49±	71.52±	13.89±
		12	0.04ab	2.65a	0.72ab	0.20a	1.65ab	2.33ab	0.09a	0.19ab	0.36ab	0.31a	0.58a	1.25a	4.84a	2.51b
	Medium	3	0.86±	7.97±	4.27±	0.61±	0.88±	9.93±	0.20±	0.72±	2.77±	0.32±	4.23±	3.93±	10.43±	3.22±
		6	0.04h	0.53f	0.22g	0.05g	0.08e	0.40h	0.03d	0.07fg	0.13d	0.08e	0.18g	0.37f	0.88f	0.60ef
		9	1.27±	17.49±	10.28±	2.20±	4.85±	28.22±	0.79±	1.53±	4.92±	2.18±	7.94±	12.24±	39.15±	4.81±
		12	0.05bc	0.56c	0.37cd	0.10cd	0.18c	1.35ab	0.04ab	0.04cd	0.12abcd	0.08b	0.31bcd	0.36c	0.68cd	0.51ef
	High	3	1.04±	18.95±	10.74±	2.75±	7.70±	32.45±	0.97±	1.31±	6.35±	2.64±	9.61±	13.74±	45.17±	6.49±
		6	0.06efg	1.43bc	0.61abc	0.18ab	0.41ab	3.70a	0.05a	0.14de	0.31abc	0.13ab	0.62a	0.81bc	3.67c	1.04de

“-” Indicates ‘not detected’. Data are expressed as mean ± SE of four replicates. Values with different letters differ significantly ($P < 0.05$)

and 12th days, at which time the mycelia of *M. alpina* grew slowly. However, the fungal biomass increased rapidly at the high nitrogen concentration on the 9th and 12th days, due to sufficient nitrogen content in the media. In other words, the low and medium urea concentrations were not conducive for the growth of *M. alpina* at the later stages of fermentation. It is likely that under nitrogen-restricted conditions, *M. alpina* initiated a stress-resistant physiological response which resulted in inhibition of mycelial protein synthesis, thereby decreasing the levels of enzymes required for mycelial growth and metabolism.

It is noteworthy that the biomass of *M. alpina* reached its maximum on the 12th day of fermentation at the high urea concentration, but the contents of total fatty acids, polyunsaturated fatty acids, and arachidonic acid were

significantly lower than the corresponding contents at the low and medium urea concentrations. These results are consistent with the findings in some studies which showed that oleaginous yeasts and molds accumulated high levels of lipids in nitrogen-restricted media (Arous *et al.* 2016; Janssen *et al.* 2019; Tossavainen *et al.* 2019). The primary function of a nitrogen source is to enhance the synthesis of proteins and nucleic acids. When the nitrogen source is limited, the synthesis of intracellular proteins and nucleic acids is blocked, and the tricarboxylic acid cycle is impeded, thereby suppressing cell proliferation and causing changes in metabolic pathways. Thus, the carbon flow in the media is channeled in the direction of lipid synthesis. A previous study showed that when a bioengineered strain of *Saccharomyces cerevisiae* was cultivated under nitrogen-




Fig. 5: Fatty acid contents of cell dry weight (mg g⁻¹) in *M. alpina* D36 cultivated with different nitrogen concentrations during fermentation. (A) Total fatty acid contents of cell dry weight; (B) polyunsaturated fatty acid contents of cell dry weight; (C) arachidonic acid contents of cell dry weight; (D) eicosapentaenoic acid contents of cell dry weight. Data with different letters differ significantly ($P < 0.05$)

limited conditions, the yields of monounsaturated fatty acids were higher than the corresponding yields under sufficient nitrogen conditions (Tang and Chen 2014). Moreover, citrate level in the *S. cerevisiae* under limited nitrogen condition was much higher than that under ample nitrogen limited condition. The accumulated citrate in cells is cleaved by ATP-citrate lyase (ACL) to acetyl-CoA, thereby providing a key substrate for fatty acid synthesis.

The oil-producing performance of *M. alpina* is closely related not only to the fatty acid content of the fungal mycelia, but also to the biomass concentration. Low biomass reduces the yield of fatty acids. Therefore, this study also used the volumetric productivity of each fatty acid component to evaluate the fatty acid production performance of *M. alpina*. The results showed that with sodium nitrate as the nitrogen source, there was little change

in the fatty acid content of *M. alpina*, but its biomass concentration was significantly lower, when compared with biomass with urea as the nitrogen source. As a result, the volumetric productivities of total fatty acids, polyunsaturated fatty acids, and arachidonic acid were very low when sodium nitrate was used as the nitrogen source. Therefore, urea is more suitable for the growth and accumulation of fatty acids in *M. alpina* than sodium nitrate. In this study, with urea as nitrogen source, low nitrogen concentration (0.05 M) was beneficial to the accumulation of fatty acids, but it was not conducive for the growth of *M. alpina*. Sufficient nitrogen at high concentration (0.2 M) promoted the growth of *M. alpina*, but inhibited the accumulation of fatty acids in mycelia. However, biomass concentration and fatty acid production of *M. alpina* were higher under medium concentration of urea (0.1 M).




Fig. 6: Fatty acid yields at different nitrogen concentrations during fermentation. (A) Total fatty acid yield; (B) polyunsaturated fatty acid yield; (C) arachidonic acid yield; (D) eicosapentaenoic acid yield. Data with different letters differ significantly ($P < 0.05$)

Generally, the inverse relationship between biomass concentration and lipid accumulation is the most significant technical bottleneck faced when using *M. alpina* to produce lipids. If the nitrogen demand can be accurately controlled, the biomass concentration and lipid content of *M. alpina* can be increased, so that the yields of polyunsaturated fatty acids, especially arachidonic acid, can be maximized.

Conclusion

The effects of nitrogen sources and nitrogen concentrations on the mycelial growth and fatty acid accumulation of *M. alpina* were determined in this study. The results suggest that the biomass concentration, and contents and yields of total fatty acids, polyunsaturated fatty acids, and arachidonic

acid are regulated when *M. alpina* is grown under different nitrogen sources and nitrogen concentrations. In this study, urea was more beneficial to the growth and fatty acid accumulation of *M. alpina* than sodium nitrate, and the biomass and fatty acid yields at the medium urea concentration were relatively high. However, the mechanism underlying the effect of nitrogen source and its concentration on the growth and fatty acid synthesis of *M. alpina* is unclear. This needs further research.

Acknowledgements

This work was supported by the Key Project of Yunnan Provincial Agricultural Union (2017FG001-015), and the Natural Science Fund of Yunnan province (2016FB030).

Author Contributions

Lingfei Li designed the study and revised the manuscript. Lingfei Li and Na Jin performed the experiments and analyzed the data. Lingfei Li wrote the first draft of the manuscript.

References

- Arous F, T Mechichi, M Nasri, G Aggelis (2016). Fatty acid biosynthesis during the life cycle of *Debaryomyces etchellsii*. *Microbiology* 162:1080–1090
- Asadi SZ, K Khosravi-Darani, H Nikoipour, H Bakhoda (2018). Production of arachidonic acid and eicosapentaenoic acid by *Mortierella alpina* CBS 528.72 on date waste. *Food Technol Biotechnol* 56:197–207
- Baky HHAE, GSE Baroty, EM Mostafa, M Enas (2020). Optimization growth of *Spirulina (Arthrospira)* platensis in photobioreactor under varied nitrogen concentration for maximized biomass, carotenoids and lipid contents. *Rec Pat Food Nutr Agric* 11:40–48
- Certik M, J Megova, R Horenitzky (1999). Effect of nitrogen sources on the activities of lipogenic enzymes in oleaginous fungus *Cunninghamella echinulata*. *J Gen Appl Microbiol* 45:289–293
- Feng P, Z Xu, L Qin, MA Alam, Z Wang, S Zhu (2020). Effects of different nitrogen sources and light paths of flat plate photobioreactors on the growth and lipid accumulation of *Chlorella* spp. GN1 outdoors. *Bioresour Technol* 301:122762
- Granger LM, P Perlot, G Goma, A Pareilleux (1993). Efficiency of fatty acid synthesis by oleaginous yeasts: prediction of yield and fatty acid cell content from consumed C/N ratio by a simple method. *Biotechnol Bioeng* 42:1151–1156
- Gu X, X Fu, L Li (2018). Effect of temperature and agitation speed on fatty acid accumulation in *Mortierella alpina*. *Intl J Agric Biol* 20:2319–2324
- Janssen JH, RH Wijffels, MJ Barbosa (2019). Lipid production in *Nannochloropsis gaditana* during nitrogen starvation. *Biology* 8; Article 5
- Kikukawa H, E Sakuradani, A Ando, S Shimizu, J Ogawa (2018). Arachidonic acid production by the oleaginous fungus *Mortierella alpina* 1S-4: A review. *J Adv Res* 11:15–22
- Lu H, H Chen, X Tang, Q Yang, H Zhang, YQ Chen, W Chen (2019). Ultra performance liquid chromatography-Q exactive orbitrap/mass spectrometry-based lipidomics reveals the influence of nitrogen sources on lipid biosynthesis of *Mortierella alpina*. *J Agric Food Chem* 67:10984–10993
- Lu J, C Peng, XJ Ji, J You, L Cong, P Ouyang, H Huang (2011). Fermentation characteristics of *Mortierella alpina* in response to different nitrogen sources. *Appl Biochem Biotechnol* 164:979–990
- Luo N, S Zhang, P Liu (2016). Effects of nitrogen and phosphorus on cell growth and lipid accumulation of tropic ocean microalgae strain *Desmodesmus* spp. WC08. *Sci Technol Food Ind* 37:223–227
- Madsen MTB, AA Bjerregaard, JD Furtado, TI Halldorsson, M Ström, C Granström, E Giovannucci, SF Olsen (2019). Comparisons of estimated intakes and plasma concentrations of selected fatty acids in pregnancy. *Nutrients* 11; Article 568
- Nisha A, G Venkateswaran (2011). Effect of culture variables on mycelial arachidonic acid production by *Mortierella alpina*. *Food Bioproc Technol* 4:232–240
- Nisha A, SP Muthukumar, G Venkateswaran (2009). Safety evaluation of arachidonic acid rich *Mortierella alpina* biomass in albino rats-a subchronic study. *Regul Toxicol Pharmacol* 53:186–194
- Raimondi S, M Rossi, A Leonardi, MM Bianchi, T Rinaldi, A Amaretti (2014). Getting lipids from glycerol: new perspectives on biotechnological exploitation of *Candida freyschussii*. *Microb Cell Fact* 13:83–94
- Tratray JBM (1984). Biotechnology and the fats and oils industry - An overview. *J Amer Oil Chem Soc* 61:1701–1712
- Sakuradani E (2010). Advances in the production of various polyunsaturated fatty acids through oleaginous fungus *Mortierella alpina* breeding. *Biosci Biotechnol Biochem* 74:908–917
- Sattayatewa C, M Arnaldos, K Pagilla (2011). Measurement of organic nitrogen and phosphorus fractions at very low concentrations in wastewater effluents. *Water Environ Res* 83:675–683
- Singh A, OP Ward (1997). Production of high yields of arachidonic acid in a fed-batch system by *Mortierella alpina* ATCC 32222. *Appl Microbiol Biotechnol* 48:1–5
- Streekstra H (1997). On the safety of *Mortierella alpina* for the production of food ingredients, such as arachidonic acid. *J Biotechnol* 56:153–165
- Stressler T, T Eisele, J Rost, E-M Haunschild, A Kuhn, L Fischer (2013). Production of polyunsaturated fatty acids by *Mortierella alpina* using submerge and solid state fermentation. *Chem Ing Technol* 85:318–322
- Tang X, WN Chen (2014). Investigation of fatty acid accumulation in the engineered *Saccharomyces cerevisiae* under nitrogen limited culture condition. *Bioresour Technol* 162:200–206
- Tang X, H Chen, T Mei, C Ge, Z Gu, H Zhang, YQ Chen, W Chen (2018). Characterization of an Omega-3 desaturase from phytophthora parasitica and application for eicosapentaenoic acid production in *Mortierella alpina*. *Front Microbiol* 9; Article 1878
- Tossavainen M, U Ilyass, V Ollilainen, K Valkonen, A Ojala, M Romantschuk (2019). Influence of long term nitrogen limitation on lipid, protein and pigment production of *Euglena gracilis* in photoheterotrophic cultures. *Peer J* 7:6624–6642
- Wang H, H Chen, G Hao, B Yang, Y Feng, Y Wang, L Feng, J Zhao, Y Song, H Zhang, YQ Chen, L Wang, W Chen (2013). Role of the phenylalanine-hydroxylating system in aromatic substance degradation and lipid metabolism in the oleaginous fungus *Mortierella alpina*. *Appl Environ Microb* 79:3225–3233
- Wiktorowska-Owczarek A, M Berezinska, JZ Nowak (2015). PUFAs: structures, metabolism and functions. *Adv Clin Exp Med* 24:931–941
- Wu GX, BY Gao, ZW Zhou, XQ Lei, LD Huang, AF Li, CW Zhang (2015). Effects of high and low nitrogen concentration on the growth and lipids accumulation pattern of two oleaginous microalgae. *Microbiol Chin* 42:1442–1452
- Yu Y, L Zhang, T Li, N Wu, L Jiang, X Ji, H Huang (2018). How nitrogen sources influence *Mortierella alpina* aging: from the lipid droplet proteome to the whole-cell proteome and metabolome. *J Proteom* 179:140–149
- Zhang H, Z Wang, Y Feng, Q Cui, X Song (2019). Phytohormones as stimulators to improve arachidonic acid biosynthesis in *Mortierella alpina*. *Enzyme Microb Technol* 131:109381