



**Full Length Article**

## Mechanism of Action of Potato Glycoalkaloids against *Fusarium solani*

Jing He<sup>1,2\*</sup>, Tian-Tian Duo<sup>1</sup>, Wei Chen<sup>1</sup> and Xiao-Yan Zhang<sup>1</sup>

<sup>1</sup>College of Forestry, Gansu Agricultural University, Lanzhou 730070, China

<sup>2</sup>Agronomy College, Gansu Agricultural University, Lanzhou 730070, China

\*For correspondence: hejing268@aliyun.com

Received 09 September 2020; Accepted 12 January 2021; Published 25 March 2021

### Abstract

The antifungal mechanism of potato glycoalkaloids was studied using a sensitive species, *Fusarium solani*. The effects of potato glycoalkaloids extract on the ultrastructure, membrane permeability, contents of reducing sugar, soluble sugar, soluble protein and mycelial fat of *Fusarium solani* were determined. Potato glycoalkaloids significantly affected *F. solani* mycelial morphology, resulting in bubbly mycelial cell walls, incomplete outer layer, discontinuous cell membrane, disorganized structures of mitochondria and other organelles, and visible leakage of cell contents. Investigation of material metabolism indicated that potato glycoalkaloids disrupted selective permeability of mycelial cell membranes; In the treatment group, the soluble protein content increased from 66.50 g mL<sup>-1</sup> to 169.51 g mL<sup>-1</sup> for 0–6 h, the soluble sugar content in extracellular fluid increased from 117.4 g mL<sup>-1</sup> to 132.5 g mL<sup>-1</sup> for 0–24 h, which were much higher than that of the control group, hindered hydrolysis of reducing sugar, affected nutrient absorption and utilization and inhibited decomposition metabolism of mycelia. Thus, potato glycoalkaloids altered the morphology of fungal mycelia, destroyed cell membrane structure, increased mycelial cell membrane permeability, and caused cell contents leakage, resulting in effective inhibition of growth and metabolism of plant pathogenic fungus and so could decrease the occurrence of plant disease. © 2021 Friends Science Publishers

**Keywords:** Antifungal mechanism; *Fusarium solani*; Mycelial morphology; Plant disease; Potato glycoalkaloids

### Introduction

*Fusarium* is one of the most important fungi in nature, and can survive in soil and plants during winter and summer, with wide distribution, diverse hosts, strong resistance, and rapid growth and reproduction (Du 2017). *Fusarium solani* is one of the most common *Fusarium* spp., which infects the host vascular tissue, destroys its conducting tissue, and produces toxins that harm crops during growth, development, and metabolism. To date, *F. solani* is one of the plant pathogens which are difficult to control in production, and can cause root diseases in various economic crops worldwide, such as white mulberry (Zhang 2013), Chinese angelica (Zhao *et al.* 2012), walnut (Zheng *et al.* 2016), medlar (Chen *et al.* 2017), wild pepper (Li *et al.* 2016), and pear (Tang *et al.* 2017). *Fusarium solani* can also cause fruit rot (Rampersad 2010) and deformity (Zhao *et al.* 2018), resulting in wilting and death of crops, affecting yield and quality, and producing huge economic losses. Currently, *F. solani* control depends on chemical fungicides such as carbendazim, mancozeb, thiram, cymoxanil–mancozeb, xinjunan acetate, or pyrazole-kresoxim-methyl (Wang *et al.* 2014; Chen *et al.* 2017; Tang *et al.* 2017). However, long-term usage of chemical fungicides can result in pesticide residues, environmental pollution, and disease resistance,

leading to a series of adverse effects (Yao *et al.* 2017). Therefore, developing alternatives for chemical fungicides using active substances in plants, such as botanical fungicides, is urgently needed.

Plants produce more than 400,000 kinds of bioactive components, such as alkaloids, organic acids, flavonoids, phenols, and plant essential oils, most of which have anthelmintic, insecticidal, antifungal, or antibacterial activities (Yao *et al.* 2017). These bioactive compounds have a broad spectrum of target organisms, are safe for non-target organisms, and are characterized by low toxicity, low residue, easy degradation, and no target resistance (He *et al.* 2006; Yoon *et al.* 2013; Zhang *et al.* 2013a, b). Plant bioactive components are a key research topic for controlling many plant diseases and developing new botanical pesticides, with prospects of wide exploitation and utilization (Li 2017; Tang *et al.* 2018). Potato glycoalkaloids are steroids produced as secondary metabolites by potatoes (Guo *et al.* 2017). The main components of potato glycoalkaloids are  $\alpha$ -solanine and  $\alpha$ -chaconine (Zhao *et al.* 2013), which represent more than 95% of the total glycoalkaloids (Kodamatani *et al.* 2005). Previous studies have shown that potato glycoalkaloids have a wide range of biological activities (Qiao 2017), and can inhibit infection or growth of fungi, bacteria, viruses, and other plant pathogenic

microorganisms, prevent insects from feeding or harming plants (Friedman 2004; Liang *et al.* 2017), exert protective effects on plants, and possess important medicinal value (Zhao *et al.* 2013). Fewell and Roddick (1993) reported that  $\alpha$ -solanine and  $\alpha$ -chaconine in potatoes can inhibit the growth of fungi such as *Alternaria brassicicola*, *Ascobolus crenulatus*, *Rhizoctonia solani*, and *Phoma medicaginis*. Furthermore, Zhao *et al.* (2013) showed that potato glycoalkaloids have high inhibitory effects on *Alternaria porri* and *Cercospora brassicae*. Likewise, Ombra *et al.* (2014) found that potato extract had antibacterial activity against *Bacillus cereus*, *Escherichia coli*, and *Pseudomonas aeruginosa* under in vitro conditions.

Currently, potato glycoalkaloids extract has been noted to have an inhibitory effect on three economic forest pathogenic fungi: *F. solani*, *Capnodium leaophilum*, and *Marssonina juglandis*. Notably, the inhibitory effect on *F. solani* was the strongest (Duo *et al.* 2017). However, the studies on control of *F. solani* disease using plant bioactive compounds as well as the antifungal mechanism of these compounds are limited. Therefore, in the study, the effects of potato glycoalkaloids extract on the ultrastructure, cell membrane permeability, and contents of reducing sugar, soluble sugar, soluble protein, and mycelial fat of *F. solani* were examined, and the inhibition mechanism was preliminarily determined. The results could provide a theoretical basis for prevention and control of economic forest diseases as well as for the development and utilization of plant-derived fungicides.

## Materials and Methods

### Materials

The potato is Qingshu variety No.9 being purchased from supermarket. The fresh potato samples were washed and dried in sunlight for several weeks to turn the potato skin green and allow germination. Subsequently, the green potato skin and buds were dried in a vacuum blast drying oven and pulverized into a powder using a plant pulverizer through an 80-mesh sieve and stored at 0–4°C until further use.

The test strain of *F. solani* was isolated from *Lycium barbarum* root rot in Gansu Province, China (Fang 1998), and its pathogenicity was confirmed based on Koch's postulates. After identification, the strain was stored at 0–4°C until further use. Potato dextrose broth (PDB) and potato dextrose agar (PDA) were employed for fungal cultivation. The PDB comprised 200 g of peeled potato, 20 g of glucose, and 1000 mL of distilled water (neutral pH). The PDA was prepared by adding 17–20 g of agar to the constituents of PDB. All reagents used were of domestic analytical grade and purchased from Gansu Zhongrui Chemical Co. Ltd., China.

### Extraction of potato glycoalkaloids

Potato glycoalkaloids were extracted using an acetic acid

extraction–ammonia precipitation method with slight modifications (Bo *et al.* 2012). In brief, 100 g of the potato sample was mixed with 400 mL of 5% acetic acid, stirred for 60 min (JB-1 magnetic stirrer, Shanghai Leici Xinjing Instrument Co. Ltd., China), and filtered (SHZ-D III circulating water vacuum pump, Gongyi Yuhua Instrument Co. Ltd., China). The residue was extracted twice with 200 mL of 5% acetic acid, and the filtrate was combined, and its pH adjusted to 11 with ammonia. After extracting three times with 200 mL of water-saturated *n*-butanol, the extracts were combined and dried on a rotary evaporator (RE-3000, Shanghai Yarong Biochemical Instrument Factory, China), and the residue mixed with 20 mL of methanol to obtain total glycoalkaloids extract. The mass concentration of the glycoalkaloids extract was 5 g mL<sup>-1</sup>.

### Effect of potato glycoalkaloids on ultrastructure of *F. solani*

The *F. solani* was inoculated onto PDA at a concentration of 0.3036 g mL<sup>-1</sup> (EC<sub>50</sub>), along with 2 g mL<sup>-1</sup> potato glycoalkaloid extract, and incubated at 25°C for 48 h. Subsequently, sterile filter paper strips (0.7 cm × 5 cm) were placed around the colony (covering an area of 5 cm × 5 cm) and incubated at 25°C. After 72 h, the edge of the colony was sampled. The collected sample was fixed by double fixation with glutaraldehyde and citric acid (Zeng 2012), and observed and photographed under a transmission electron microscope (JEM2000EX; JEOL, Japan).

### Effect of potato glycoalkaloids on cell membrane permeability of *F. solani*

The mycelia of *F. solani* were cultured in PDB for 4 days and washed four times with ultrapure sterile water. Then, the washed mycelia were freeze-dried to a constant weight (Labconco freeze drier, USA) and 1 g of the mycelia was transferred into 5 mL of potato glycoalkaloid (EC<sub>50</sub>) extract and incubated at 25°C under constant shaking at 120 r min<sup>-1</sup>. Subsequently, conductivity of the culture broth was measured (DDB-303A digital conductivity meter, Shanghai Yidian Scientific Instrument Co. Ltd., China) hourly during 0–9 h. Finally, the culture broth was boiled in a water bath (HH-S6 digital display thermostat water bath, Jintan Medical Instrument Factory, China) for 10 min and conductivity was determined. The experiment was repeated thrice, with sterile water and methanol as controls. The permeability of cell membrane was expressed as relative permeability (%) = (relative time conductivity value - initial conductivity value) / (kill conductivity value - initial conductivity value) × 100% (Shen 2014).

### Effect of potato glycoalkaloids on soluble mycoprotein of *F. solani*

The effect of potato glycoalkaloids on soluble mycoprotein was determined by Coomassie Brilliant Blue G-250 staining

(Song 2010). In brief, mycelia of *F. solani* were cultured in PDB for 4 days, and washed four times with ultrapure sterile water. Then, the washed mycelia were freeze-dried (Labconco) to a constant weight and 1 g of the mycelia was added to 5 mL of potato glycoalkaloids (EC<sub>50</sub>) extract, and sampled at 0, 2, 4, 6, and 8 h. The collected samples were centrifuged (D-37520 centrifuge, Heraeus Biofuge, Germany), and the absorbance of samples was recorded at 595 nm (Jenway 6505 UV/Vis UV Spectrophotometer; Gaonan Instrument (Shenzhen) Co. Ltd. China). The protein concentration was calculated according to the protein standard curve, and the experiment was repeated thrice, with sterile water and methanol as controls.

#### **Effect of potato glycoalkaloids on soluble sugar of *F. solani***

The effect of potato glycoalkaloids on soluble sugar in *F. solani* was determined by anthrone colorimetry (Yao and Xu 1992). In brief, *F. solani* mycelia were cultured in PDB for 4 days, and then washed four times with ultrapure sterile water. Then, the washed mycelia were freeze-dried (Labconco) to a constant weight and 1 g of mycelia mixed with 5 mL of potato glycoalkaloids (EC<sub>50</sub>) extract, and sampled at 0, 1, 2, 4, 6, 8, 10 and 12 h. Subsequently, collected samples were subjected to centrifugation (6000 rpm; 5 min), heated with anthrone reagent, and cooled to room temperature, and the absorbance of the samples measured at 620 nm. The soluble sugar content was calculated according to the glucose standard curve, and the experiment was repeated thrice, with sterile water and methanol as controls.

#### **Effect of potato glycoalkaloids on reducing sugar of *F. solani***

The effect of potato glycoalkaloids on reducing sugar in *F. solani* was determined by 3, 5-dinitrosalicylic acid (DNS) method (Chen 2002). In brief, *F. solani* mycelia were cultured in PDB for 4 days, and washed four times with ultrapure sterile water. Then, the washed mycelia were freeze-dried (Labconco) to a constant weight and 1 g of mycelia was added to 5 mL of potato glycoalkaloid (EC<sub>50</sub>) extract, and sampled hourly during 0–8 h. Subsequently, 1 mL of the collected samples was respectively subjected to centrifugation (6000 rpm; 5 min), and 0.5 mL of the supernatant was mixed with 1.5 mL of distilled water and 1.5 mL of DNS reagent to determine the absorbance at 520 nm. The reducing sugar content was calculated according to the standard curve, and the experiment was repeated thrice, with sterile water and methanol as controls.

#### **Effect of potato glycoalkaloids on fat content of *F. solani***

The oil weight method was employed for sample processing and fat content determination (Li 1987). The *F. solani*

mycelia were cultured in PDB for 4 days and washed four times with ultrapure sterile water. Then, the washed mycelia were freeze-dried (Labconco) to a constant weight and 1 g of the mycelia was added to 5 mL of potato glycoalkaloids (EC<sub>50</sub>) extract, and sampled at 24, 48, and 72 h. Subsequently, the collected samples were filtered (SHZ-D III circulating water vacuum pump) and rinsed with redistilled water four times; the obtained wet hyphae were dried at 60–80°C for 4 h, smashed with a mortar and pestle, and filtered through a mesh sieve to obtain dried powder. The procedure was repeated thrice, and sterile water and methanol were used as controls. For fat content determination, the Soxhlet extractor was cleaned, heated in a blast drying oven at 105°C for 20 min, cooled to room temperature, and weighed (m). Then, 12-cm quantitative filter paper was weighed (m<sub>1</sub>) and made into a bucket, and 2 g of the dried sample powder were added to the filter paper bucket and weighed (m<sub>2</sub>). The difference between the two masses indicated the quality of the dried sample powder (m<sub>2</sub>-m<sub>1</sub>). Subsequently, the dried sample was soaked in petroleum ether overnight, and heated in a thermostat water bath for 65°C. Reflux extraction was performed for 12 h using a Soxhlet fat extractor, and then the extract was heated at 100°C for 8 h, cooled to room temperature, and weighed (m'). Crude fat was calculated as follows: crude fat (%) = (m'-m)/(m<sub>2</sub>-m<sub>1</sub>) × 100%.

#### **Statistical analysis**

All the above measurements were repeated at least three times. The results were expressed as mean value ± standard errors calculated by Microsoft Excel 2007. The variance was examined using SPSS 19.0 and the difference investigated by employing Duncan's new complex range method (Zhang 2013).

## **Results**

#### **Effect of potato glycoalkaloids on ultrastructure on *F. Solani***

Once the cell wall or cell membrane is destroyed, the cell cannot maintain its inherent structure in morphology, and the mycelium will be twisted, deformed, or even broken. In this study, the controls without potato glycoalkaloids treatment presented thin and uniform cell walls, complete structure, hyphae surrounded by a continuous outer layer, intact internal tissue structure, no extravasation of cell contents, normal development, and clearly visible vacuoles, mitochondria, and other organelles (Fig. 1A). However, after treatment with EC<sub>50</sub> potato glycoalkaloids, the fungal cell walls became thinner and irregular, cell internal structure was disrupted, and some membrane structures were incomplete (Fig. 1B). Furthermore, after treatment with 2 g mL<sup>-1</sup> potato glycoalkaloids, cell walls of mycelia had a bubble shape, outer layer components were altered,


cell wall structure was incomplete, the outer cell membrane was discontinuous, structure of mitochondria and other organelles was not obvious, and extracellular inclusions were exuded (Fig. 1C). In contrast, following treatment with methanol, the cell wall structure was clear, mitochondria and ribosomes were clearly visible, and the vacuolar structure was intact, indicating that methanol did not affect antibacterial activity of potato glycoalkaloids (Fig. 1D). These findings indicated that potato glycoalkaloids destroyed the cell surface morphology of *F. solani*, severely damaging the cytoplasm, mitochondria, and other organelles, as well as increasing the cell membrane permeability and so causing leakage of cell contents.

### Effect of potato glycoalkaloids on cell membrane permeability of *F. solani*


Selective permeability is the most basic function of the cell membrane. If the cell loses this function, the cell will die. In this study, the Fig. 2 shows the effects of potato glycoalkaloids on the cell membrane permeability of *F. solani*. The relative cell membrane permeability increased with time in both the control and treatment groups. However, the relative cell membrane permeability was significantly higher for the treatment compared with the control group. In both the control and treatment groups, the relative cell membrane permeability rapidly increased during 1–4 h, with a greater increase for the treatment than the control group. After 4 h, the difference in relative cell membrane permeability quickly broadened between the two groups, indicating that the *F. solani* cell membrane damage caused by potato glycoalkaloids at EC<sub>50</sub> concentration intensified at 4 h. Subsequently, the relative cell membrane permeability of the control group gradually decreased during 4–9 h, whereas that of the treatment group increased by 25.2% at 9 h, and remained stable at 8 and 9 h.

### Effect of potato glycoalkaloids on soluble protein of *F. solani*

Material metabolism is divided into anabolism and catabolism, both of which maintain dynamic balance in normal life activities. When this balance is broken, it will cause diseases of the organism and even death. In this study, the soluble protein leakage in the treatment group exhibited an upward trend with time, which significantly differed from that in the control group (Fig. 3). During 0–6 h, soluble protein leakage in the treatment group increased from 66.50 to 169.51  $\mu\text{g mL}^{-1}$ , which was much higher than that in the control group. However, during 6–24 h, the soluble protein leakage in the treatment group gradually decreased with time, possibly due to consumption of soluble protein by newly formed cells of *F. Solani* for growth. After 24 h, the soluble protein leakage in the treatment and control groups gradually increased and stabilized.



**Fig. 1:** Effect of potato glycoalkaloids on ultrastructure of *F. solani* (2550 $\times$ )  
 A: Control 1 (sterile water); B: Treatment 1 (EC<sub>50</sub>); C: Treatment 2 (2 g mL<sup>-1</sup>); D: Control 2 (methanol).  
 CW: Cell wall; V: Vacuole; M: Mitochondrion



**Fig. 2:** Effect of potato glycoalkaloids on cell membrane permeability of *F. solani*

### Effect of potato glycoalkaloids on soluble sugar of *F. solani*

Soluble sugar leakage in the treatment group significantly differed from that in the control with time (Fig. 4). During 0–24 h, soluble sugar leakage in the treatment group increased from 117.4 to 132.5  $\mu\text{g mL}^{-1}$ , which was much higher than that in the control group. However, after 24 h, leakage in the treatment group decreased with time owing to consumption of soluble sugar in the extracellular fluid by newly formed *F. solani* cells for growth.

### Effect of potato glycoalkaloids on reducing sugar content of *F. solani*

The reducing sugar content in the extracellular fluid

significantly differed between the treatment and control group with time (Fig. 5). In both groups, the reducing sugar content in the extracellular fluid sharply decreased during 0–2 h, but gradually reduced during 2–72 h, indicating that potato glycoalkaloids significantly decreased the absorption and utilization of reducing sugar by *F. solani*.


### Effect of potato glycoalkaloids on fat content of *F. solani*

The fat content in the extracellular fluid of both the treatment and control groups increased during 0–48 h, with significantly higher fat leakage for the treatment compared with the control group (Fig. 6). The fat content in the extracellular fluid of the treatment group increased by 54.20 and 52.07%, compared with sterile water and methanol control groups at 48 h, respectively, indicating that potato glycoalkaloids damaged the cell structure and caused fat content leakage from *F. solani* plasma membrane. However, after 48 h, fat content in the extracellular fluid of both treatment and control groups showed a downward trend, possibly resulting from a weakening effect of potato glycoalkaloids or a self-remediation mechanism of *F. solani*.


### Discussion

Plant bioactive substances mainly target the cell membrane of fungi by altering membrane stability and causing damage to the membrane structure and extravasation of inclusions, ultimately resulting in fungicidal or fungistatic effect (Zhou *et al.* 2014). Previous studies have shown that glycoalkaloids predominantly combine with sterols in the fungal cell membrane, forming a complex that destroys membrane integrity and causes loss of normal membrane function (Sun 2014). In the present study, transmission electron microscopy showed that *F. solani* morphology was distorted after potato glycoalkaloids treatment, and that some cell walls were blurred or even lost. Additionally, structure of the cell and vacuolar membranes was destroyed and the organelles distorted. These observations indicated that potato glycoalkaloids can affect the surface morphology of *F. solani*, leading to incomplete membrane structure and causing serious damage to cytoplasm and mitochondria, consistent with the effect of pyrolin on *Monilinia fructicola* (Wu *et al.* 2009), ethyl acetate extract of amaranth on *Xanthomonas citri* (Liao *et al.* 2017), and water-soluble chitosan on the ultrastructure of *Fusarium* (Jia *et al.* 2016).

Electrical conductivity can indirectly reflect cell membrane permeability, and a higher electrical conductivity of a culture broth signifies enhanced electrolyte leakage and increased damage to cell membranes (Shang 2017). Peng *et al.* (2017) showed that extract of *Cynanchum atratum* could enhance the cell membrane permeability of *Penicillium italicum*, while Zhang *et al.* (2008) revealed that the extract of *Xanthium sibiricum* induced changes in membrane permeability of *Botrytis cinerea*, resulting in increased



**Fig. 3:** Effect of potato glycoalkaloids on soluble protein leakage of *F. solani*



**Fig. 4:** Effect of potato glycoalkaloids on soluble sugar leakage of *F. solani*


conductivity of the culture broth. Furthermore, Liu *et al.* (2018a) found that the total saponins and total ginsenoside from ginseng stem and leaf can induce changes in the permeability of the mycelium membrane of *F. solani*, respectively, leading to increased conductivity of the culture broth. Liu *et al.* (2018b) showed that limonene can increase the cell membrane permeability of *Pseudomonas aeruginosa* and destroy its cell morphology and integrity, thus effectively inhibiting its growth. Similarly, in the present study, the cell membrane permeability of *F. solani* increased after treatment with potato glycoalkaloids, and transmission electron microscopy revealed leakage of intracellular contents and destruction of cell membrane structure. These results showed that the cell membrane and integrity of *F. solani* were destroyed by potato glycoalkaloids, which directly affected the physiological functions of the cell membrane, such as exchange of intracellular and extracellular substances and regulation of cell growth, leading to disturbance in fungal metabolism.

It must be noted that plant bioactive compounds also affect the morphology and structure of fungal mycelia, causing deformity, kinking, swelling, and lysis. As a result, the mycelial soluble protein and soluble sugar can leak into


the culture medium (Zhou *et al.* 2011; Fan *et al.* 2015; Zhang *et al.* 2016). In fungi, soluble protein mainly comprises various enzymes involved in metabolism. During growth, the fungi secrete proteins that penetrate the cell membrane into the thallus fluid through osmosis. Thus, changes in the content of these proteins reflect alteration in the total cellular metabolism (Liu *et al.* 2016). Sugar catabolism provides the energy needed for fungal growth, and inhibition of the absorption and utilization of sugar could lead to lack of energy required by the fungi, affecting growth and propagation of the thallus (Liu *et al.* 2016). The total lipid content in the fungal cell membrane affects cell membrane fluidity, and a decrease in the total lipid content may lead to a reduction in cell membrane fluidity (Shang 2017). Thus, one approaches to achieve fungicidal or fungistatic effect is to inhibit fungal metabolism (Zhou *et al.* 2014). In the present study, the contents of total sugar, protein, and fat in *F. solani* initially increased and then decreased with time after treatment with potato glycoalkaloids. However, the content of reducing sugar decreased in *F. solani* treated with potato glycoalkaloids, but significantly increased in control cells without glycoalkaloids treatment. Similar findings were also reported by Wu (2008), who showed that the contents of total sugar, reducing sugar, protein, and fat in *B. cinerea* treated with propamidine were significantly higher than those in control *B. cinerea* without treatment, indicating that plant bioactive compounds inhibited catabolism in fungi. Biological catabolic systems are complex, and disturbance in a certain catabolic link can obstruct the entire catabolic process, threatening life of the organism. However, self-remedial mechanisms can overcome the blocked metabolic processes to continue life activities (Zeng 2012). Therefore, it is possible that self-remedial mechanisms of *F. solani* allowed the fungal cells to thrive after treatment with potato glycoalkaloids, resulting in an initial increase and subsequent decrease in contents of total sugar, protein, and fat with treatment duration.

## Conclusion

Results in the present study showed that potato glycoalkaloids significantly affected the morphological structure of *F. solani*. Treatment with potato glycoalkaloids resulted in bubbly and undulated mycelial cell walls, incomplete outer structure, discontinuous cell membrane, disordered structure of mitochondria and other organelles, and visible extracellular contents. The material metabolism analysis demonstrated that potato glycoalkaloids destroyed the selective permeability of fungal cell membranes and causing extravasation of large quantities of internal lipids, proteins, and sugars. This hindered the hydrolysis of reducing sugars, affecting the absorption and utilization of nutrients, and ultimately inhibiting catabolism in fungi. However, knowledge of the specific antifungal mechanism of potato glycoalkaloids is limited. Therefore, further



**Fig. 5:** Effect of potato glycoalkaloids on reducing sugar content of *F. solani*



**Fig. 6:** Effect of potato glycoalkaloids on fat content of *F. solani*

research on the effects of potato glycoalkaloids on fungal respiratory metabolism (*e.g.*, related enzymes activities) and energy metabolism (*e.g.*, inhibition of electron transport and oxidative phosphorylation, and information expression of nucleic acids and other molecular substances) is necessary for a better understanding of the antifungal mechanism of these compounds and for acquiring comprehensive and systematic theoretical support for the development and utilization of botanical pesticides.

## Acknowledgements

The work was supported by grants from the National Natural Science Foundation of China (32060341), Special Funds for Discipline Construction of Gansu Agricultural University (GAU-XKJS-2018-096), and Supporting Funds for Youth Mentor of Gansu Agricultural University (GAU-QDFC-2018-07) and Postdoctoral Research Projects (2015GSZYZZ002). We thank International Science Editing (<http://www.internationalscienceediting.com>) for editing this manuscript.

## Author Contributions

JH planned the experiments, JH and TTD interpreted the

results, JH, TTD and WC contributed to writing and XYZ statistically analyzed the data and prepared illustrations.

## Conflicts of Interest

The authors declare that they have no conflict of interest.

## Data Availability

The data will be made available on acceptable requests to the corresponding author.

## Ethics Approval

Not applicable.

## References

- Bo LL, XY Sun, H Lin, MJ Dai, JP Niu, JW Zhang, D Wang (2012). Extraction of glycoalkaloids from potatoes. *J Anhui Agric Sci* 40:1448–1491
- Chen W, J He, TT Duo, RJ Xue, HJ Xu, HJ Yang (2017). Laboratory toxicity test of several fungicides against root rot pathogens of *Lycium barbarum*. *J Gansu Agric Univ* 52:109–113
- Chen YQ (2002). Determination of reducing sugar and total sugar content. In: *Methods and Techniques of Biochemistry Experiment*, pp:97–99. Science Press, Beijing, China
- Du B (2017). Classification and research status of *Fusarium*. *J Taiyuan Univ* 35:68–70
- Duo TT, J He, W Chen, JW Zhang, SH Feng, HY Ma, J Du (2017). Antibacterial activity of potato glycoalkaloids and their effects on cell membrane permeability. *J Gansu Agric Univ* 52:50–54
- Fan MC, YJ Sun, WX Li, XK Kou (2015). Screening and mechanism study of antibacterial effect of natural plant extracts. *J Chin Inst Food Sci Technol* 15:180–185
- Fang ZD (1998). Isolation and culture of fungi. In: *Research Methods of Plant Disease*, pp:122–145. China Agric Press, Beijing, China
- Fewell AM, JG Roddick (1993). Interactive antifungal activity of the glycoalkaloids  $\alpha$ -solanine and  $\alpha$ -chaconine. *Phytochemistry* 33:323–328
- Friedman M (2004). Analysis of biologically active compounds in potatoes (*Solanum tuberosum*), tomatoes (*Lycopersicon esculentum*), and jimson weed (*Datura stramonium*) seeds. *J Chromatogr A* 1054:143–155
- Guo HX, JJ Zhang, R An, Y Qiao, WH Shi, J Shi, JW Zhang (2017). Study on the regulation of glycosidic alkaloid synthesis in green tissues of potato aboveground parts. *J Hortic* 44:1105–1115
- He J, ZQ Ma, X Zhang (2006). An overview of botanical pesticides. *J N W Agric For Univ* 34:79–85
- Jia RX, M Qiu, JY Huang, Y Zhou (2016). Antimicrobial mechanism of chitosan and its derivatives against *Fusarium*. *Chin J Food* 16:70–75
- Kodamatani H, K Saito, N Niina, S Yamazaki, Y Tanaka (2005). Simple and sensitive for determination of glycoalkaloids in potato tubers by high-performance liquid chromatography with chemiluminescence detection. *J Chromatogr A* 1100:26–31
- Li C (1987). Separation of volatile. In: *Principles and Techniques of Food Analysis*, 1<sup>st</sup> Edition, pp: 12–30. Sci Technol Lit Press, Beijing, China
- Li CB (2017). *Bioactivity of Ten Plant Extracts against Pathogenic Bacteria in Tea Leaves*. Huazhong Agricultural University, College of Science, Wuhan, Hubei, China
- Li SJ, TH Zhu, TM Cen, S Han (2016). Screening, identification, colonization and control effect of *Bacillus* spp. against root rot of *Zanthoxylum bungeanum*. *J N W Univ Agric For Sci Technol (Nat Sci Edn)* 44:114–122
- Liang KH, LG Lu, DZ Zhu, H Zhu (2017). Research progress of potato glycoalkaloids. *Food Res Dev* 38:195–199
- Liao SL, W Yin, XL Liao, Y Zhang (2017). Study on the inhibition and mechanism of ethyl acetate extract from Amaranth on Citrus canker. *J Hunan Agric Univ (Nat Sci Edn)* 43:544–550
- Liu FF, AH Zhang, FJ Lei, YH Xu, LX Zhang (2018a). Inhibitory effect of total saponins from ginseng stem and leaf on *Fusarium oxysporum* and its bacteriostatic mechanism. *J Jilin Agric Univ* 40:85–91
- Liu X, JN Wang, WX Chen, RH Chen, GF Zhang (2018b). The antibacterial activity and mechanism of limonene against *Pseudomonas aeruginosa*. *Food Industr Sci Technol* 39:1–59
- Liu JY, QQ Zhao, ZY Zhang, H Den, N Qi, H Li (2016). Preliminary study on the antibacterial activity and mechanism of sterols in seed oil of *Xanthoceras sorbifolia* L. *Chin Oil* 41:29–33
- Ombra MN, F Fratianni, T Granese, F Cardinale, A Cozzolino, F Nazzaro (2014). *In vitro* antioxidant, antimicrobial and anti-proliferative activities of purple potato extracts (*Solanum tuberosum* cv. vitelotte noire) following simulated gastro-intestinal digestion. *Nat Prod Res* 14:1–5
- Peng X, YH Chen, JY Chen (2017). Antifungal mechanism of *Cynanchum atratum* Bunge extracts on *Penicillium italicum* and control effect on blue mold of navel orange. *J Plant Pathol* 47:398–405
- Qiao Y (2017). *Identification and Functional Analysis of miRNAs Induced by Photoinduced Glucoside Alkaloid Metabolism in Potato*. Gansu Agricultural University, College of Agronomy, Lanzhou, Gansu, China
- Rampersad S (2010). First report of *Fusarium solani* causing fruit rot of sweet pepper in Trinidad. *Plant Dis* 94:1375
- Shang CY (2017). *Study on the Antibacterial Mechanism of  $\beta$ -pinene on Penicillium Citrinum*. Huazhong Agricultural University, College of Food Science and Technology, Wuhan, Hubei, China
- Shen Y (2014). *Study on the Screening and Identification of Endophytic Actinomycetes from Maize and the Mechanism of Inhibition of Maize*. Northeast Agricultural University, College of Agronomy, Harbin, Heilongjiang, China
- Song L (2010). *Inhibition and Mechanism of BIT against Botrytis Cinerea*. Northwest University, College of Life Science, Xi'an, Shaanxi, China
- Sun PZ (2014). *Antimicrobial Activity of Potato Alpha-solanine against Botrytis Cinerea In vitro*. Inner Mongolia Agricultural University, College of Food Science and Engineering, Hohhot, Inner Mongolia, China
- Tang XH, LY Zhong, T Tang, W Fli, X Li, L Jiang, G Zhao, JL Zhao (2018). Inhibition of extracts from six kinds of perfume plants on *Rhizoctonia solani* of buckwheat. *J Jiangxi Agric Univ* 40:89–95
- Tang XM, TY Lin, SS Zhou, GF Li, P Liu, ZF Ye, ZY Wu, DS Wang, LW Zhu (2017). Isolation and identification of pathogenic bacteria of pear root rot and screening of effective control agents. *J Nanj Agric Univ* 40:76–83
- Wang YX, YK Xu, ZM Li (2014). Selection of 8 fungicides on main pathogens of seedling blight of *Acanthopanax senticosus*. *Jilin Agric* 24:12
- Wu FL (2008). *Preliminary Study on the Antibacterial Mechanism of Propane Oxime against Botrytis Cinerea*. Northwest A&F University, College of Food Science and Engineering, Yangling, Shaanxi, China
- Wu ZY, Y Wang, QJ Ai (2009). Inhibitory effect of pyrolysin on peach brown rot pathogen and its bacteriostatic mechanism. *Chin Agric Sci* 42:2784–2792
- Yao HW, KK Ge, JL Pan, WL Liu, L Ma (2017). Screening of antibacterial activity of extracts from 29 plants such as *Syringa*. *J N E For Univ* 45:35–39
- Yao LH, Q Xu (1992). Simplified study on the determination of total sugar content in food by anthrone colorimetry. *Food Industr* 3:40–42
- Yoon MY, B Cha, JC Kim (2013). Recent trends in studies on botanical fungicides in agriculture. *Plant Pathol J* 29:1–9
- Zeng R (2012). *Study on the Antibacterial Active components, antibacterial mechanism and the effect on Citrus Preservation and Preservation of Impatiens balsamina*. Nanchang University, College of Life Science and Food Engineering, Nanchang, Jiangxi, China
- Zhang L (2013). Analysis of variance. In: *The Application of SPSS19.0 (Chinese version) in Biostatistics*, 3<sup>rd</sup> edn, pp:16–35. Xiamen University Press, Xiamen, China

- Zhang W, JJ Liao, GL Zhu, H Zhang, XL Duan, SS Zhu, M Yang (2013a). Study on the inhibitory activity of 8 plant volatiles and extracts against root rot on *Panax notoginseng*. *Chin Agric Sci Bull* 29:197–201
- Zhang X, ZF Wu, W Li, ZQ Ma, JT Feng (2013b). New progress in the development and application of botanical pesticides-Introduction to special biological activities. *Pestic Sci Manage* 34:24–31
- Zhang XH, J He, HM Shen (2008). Inhibitory effect of *Xanthium sibiricum* extract against *Botrytis cinerea* and its bacteriostatic mechanism. *J Grass Ind* 3:99–104
- Zhang XM, QL Xue, R Xiao, LZ Zhao, Q Lin, YJ Hu (2016). Bacteriostatic mechanism of extracts from mistletoe against *Staphylococcus aureus*. *Mod Food Sci Technol* 32:82–88
- Zhao QF, L Liu, QX Li, HY Li (2012). Study on the inhibition effects of 15 plant aqueous extracts to the growth of *Angelica sinensis* root rot fungi. *J N E Norm Univ (Nat Sci Edn)* 48:66–69
- Zhao XS, SY Li, DJ He, J Wang (2013). Study on the structure-activity relationship of potato glycoalkaloids against fungal activity. *Food Ind Sci Technol* 34:159–163
- Zhao YL, W Wang, TT Pei, G Tao (2018). Isolation and identification of the pathogen of mango malformation in Guizhou. *S W Chin J Agric Sci* 31:494–499
- Zheng L, J Zhang, WJ Ma, Y Pen, TH Zhu (2016). Effects of *Fusarium solani* on physiological indexes of resistance of different varieties of walnut. *Guangxi Plants* 36:651–657
- Zhou L, BY Yun, YJ Wang, MJ Xie (2011). Antimicrobial mechanism of emodin against *Staphylococcus aureus*. *Chin J Biochem Mol Biol* 27:1156–1160
- Zhou MJ, CP Wan, JY Chen (2014). Screening of high-efficiency antimicrobial agents of Chinese herbal medicine against Citrus green mold and their antimicrobial mechanism. *Mod Food Sci Technol* 30:144–149