



**Full Length Article**

## Potential Inhibitory Effects of *Rhizopus oligosporus* on the Growth of *Aspergillus flavus* FNCC6109 in Corn Seeds

IBG Darmayasa<sup>1</sup>, AAK Darmadi<sup>1</sup>, Arofi<sup>1</sup>, IW Suanda<sup>2</sup> and IK Widnyana<sup>3\*</sup>

<sup>1</sup>Microbiology Laboratory Faculty of Mathematics and Natural Sciences, Udayana University Denpasar, Indonesia

<sup>2</sup>Department of Biology Education, Faculty of Teacher Training and Education, University of PGRI Mahadewa, Indonesia

<sup>3</sup>Department of Agrotechnology, Faculty of Agriculture, Mahasaraswati Denpasar University, Indonesia

\*For Correspondence: widnyanaketut@gmail.com

Received 08 April 2022; Accepted 28 July 2022; Published 25 August 2022

### Abstract

Corn (*Zea mays* L.) has the highest carbohydrate content after rice but is contaminated easily by *Aspergillus flavus* with its Aflatoxin, which decreases corn quality. Chemical control of *A. flavus* has side effects, so there is a need for safe control techniques, namely biological control. This study aimed to determine the potential of the culture filtrate of *Rhizopus oligosporus* in the control of *A. flavus* FNCC6109 on corn kernels carried out *in vitro* and *in vivo*. *In vitro* testing was carried out using the dual culture method after *R. oligosporus* incubated for 3, 4 and 5 days. While the *in vivo* treatment was carried out by giving culture filtrate to corn kernels with a concentration of 0, 10, 20, 30, 40 and 50% (v/v) into corn kernels. *In vitro* tests were carried out by measuring the diameter of the colonies of *A. flavus* FNCC6109 on PDA media that give culture filtrate, while *in vivo* testing was determined by the plating method. The results showed that the culture filtrate of *R. oligosporus* significantly ( $P \leq 0.05$ ) inhibited *A. flavus* FNCC6109 both *in vitro* and *in vivo*. This study concluded that *R. oligosporus* could inhibit the growth of *A. flavus* with the highest inhibition percentage at  $67.27 \pm 2.70$  with a 5-day incubation period, and the lowest was at  $61.27 \pm 5.13$  with a 3-day incubation period. © 2022 Friends Science Publishers

**Keywords:** *Aspergillus flavus*; Biological control; Corn; *Rhizopus oligosporus*; *Zea mays*

### Introduction

Maize or corn is a food ingredient with the second-highest carbohydrate content after rice in Indonesia and comes third in the world after wheat. Maize or corn is used as a raw ingredient for the animal feed manufacturing industry. The need for corn in Indonesia was increased by 1.11% per year in 2010–2014. The official statistics for the province of Bali reported that corn production in 2015 was 40.603 tons, a decrease compared to 2013. The decline was caused by several factors, such as the decreasing of land areas, the farming transitions into horticultural crops, and lack of water supply (CBS 2014).

Corn as a raw and feed ingredient must have good quality. The most frequent problem in public and among corn farmers is the contamination of aflatoxin compounds produced by *A. flavus* (Montalbano *et al.* 2021). This fungus grows easily in tropical areas. In Indonesia, the fungus *A. flavus* has the potential to grow well and produce aflatoxins because supported by optimal levels of humidity, rainfall, and temperature for the growth of *A. flavus*. Frequent contamination of aflatoxins can easily be found in whole grains such as corns and peanuts. Several causes of aflatoxins contamination are the post-harvest handling,

harvest storage, and processing of products made from corn. These factors are common in corn farmers and companies that use corn as raw material (Rahayu *et al.* 2010). The contamination of aflatoxin compounds in corn kernels results in losses to corn farmers and health problems in animals and humans, so the control and prevention of *A. flavus* are needed.

The prevention of aflatoxins contamination in raw and feed ingredients can be conducted more effectively by inhibiting the growth of *A. flavus*. It can be physically conducted through reduction of water content in the materials, lowering of temperature, and modification of storage conditions. It can be chemically conducted by giving disinfectants acidic and alkaline substances. Chemical control requires a lot of money regarding the high cost of the chemicals needed. In addition, at the farmers' level, it is difficult to do so; therefore it is necessary to do the control biologically. The biological control of fungal pathogens can be achieved by using antagonistic bacteria such as PGPR (Sharf *et al.* 2021), and various antagonistic fungi such as species of *Trichoderma* (Ali *et al.* 2020), *Aspergillus* (Khan and Javaid 2022a, b) and *Penicillium* species (Khan and Javaid 2022c). According to Mauro *et al.* (2018), biological prevention is more effective than physical and chemical

prevention. Given the relatively fast microbial growth with a short generation time, it can be produced on a large scale.

*Rhizopus oligosporus* is a nonpathogenic fungus. This fungus is likely used for the fermentation process of raw ingredients becoming products that have high nutritional value, such as the fermentation of soybeans using the *Rhizopus* spp. yeast becoming into tempeh. Nursadin and Supriyanto (2012) reported that *Rhizopus* spp. has a high ability to compete and able to inhibit the growth of pathogenic fungi. The results of previous studies showed that *Rhizopus* sp. could inhibit *Fusarium oxysporum* by 60%. Therefore, in the present study, *R. oligosporus* was evaluated for its potential to control the growth of *A. flavus* FNCC6109.

## Materials and Methods

### Research location

The research of the potential of *R. oligosporus* filtrate in controlling the growth of *A. flavus* FNCC6109 was conducted in the Microbiology Laboratory, Department of Biology, Faculty of Mathematics and Natural Sciences Udayana University, Indonesia.

### Isolation of *R. oligosporus* in tempeh

The isolation of *R. oligosporus* was conducted by aseptically taking as much as one loop of the part of the colony suspected of being *R. oligosporus* which grew on the surface of the tempeh. This part of the colony was then placed right in the middle of Petri dishes which already contained PDA media. The Petri dishes were incubated at 28°C temperature for four days. The growth of fungal colonies on Petri dishes was observed macroscopically and microscopically by referring to the Pitt and Hocking (1997) identification book. The fungi that showed the characteristics of *R. oligosporus* were then being re-isolated until getting a pure culture.

### Regeneration of *A. flavus* FNCC6109

The *A. flavus* FNCC6109 isolate was obtained from the stock culture of the Microbiology Laboratory, Department of Biology FMIPA, Udayana University. The stock culture was rejuvenated by taking the hyphal flakes using a needle and then implanted right in the middle of the Petri dish that contained PDA media, then being incubated at 28°C for four days. The growing colonies of *A. flavus* FNCC6109 were used for the next testing stage.

### Inhibition test of *R. oligosporus* against the growth of *A. flavus* FNCC6109

The inhibition of *R. oligosporus* was conducted *in vitro* using the dual culture method. The procedures started with taking the culture of *R. oligosporus* and *A. flavus* FNCC

6109 with a 5 mm diameter cork borer. Both colonies were grown side by side with a distance of 3 cm in Petri dishes containing PDA media, then being incubated at 28°C and measured the diameter for four days. Subsequently, the same control was conducted; however, only one type of fungi was grown. The antagonism effect of *R. oligosporus* against *A. flavus* FNCC6109 could be calculated with the PIRG (percentage inhibition of radial growth) (Singh and Vijay 2011):

$$\text{PIRG (\%)} = \frac{R1 - R2}{R1} \times 100\%$$

PIRG: Percentage Inhibition of Radial Growth

R1: Colony area of *A. flavus* FNCC6109 without the antagonist (control)

R2: Colony area of *A. flavus* FNCC6109 with the antagonist (dual culture)

### Inhibition test of *R. oligosporus* culture filtrate against the growth of *A. flavus* FNCC 6109

A bottle of 100 mL PDB (potato dextrose broth) was prepared and then inoculated with the *R. oligosporus* that had the potential of inhibiting *A. flavus* FNCC 6109. Then, it was incubated for 3, 4 and 5 days at 28°C. Once the incubation period was over, the inhibition of the *R. oligosporus* culture filtrate was tested *in vitro* against the growth of *A. flavus* FNCC6109. This test started off with preparing the culture filtrate of *R. oligosporus* and took as much as 1 mL of it and placed on the Petri dish, which would be poured with PDA media afterward, then left to solidify. Next, the *A. flavus* FNCC6109 colony was taken using a 5 mm diameter cork borer and incubated for four days. After that, control was made by growing colonies of *A. flavus* FNCC6109 on PDA media without being given any culture filtrates. The effect of culture filtrate could be determined by measuring the colony area of *A. flavus* FNCC6109 using the formula:

$$\text{Inhibition (\%)} = \frac{L1 - L2}{L1}$$

L1: Colony area of *A. flavus* FNCC6109 without the antagonist (control)

L2: Colony area of *A. flavus* FNCC6109 with control

### Inhibition of *R. oligosporus* culture filtrate against *A. flavus* FNCC 6109 in corn kernels

A sterile Petri dish was prepared and contained with solid PDA media; colonies of *A. flavus* fungi were planted right in the middle of the Petri dish with a diameter of 5 mm and then incubated at a temperature of 28°C for four days. The grown colonies dripped with sterile water as much as 5 mL, then rubbed it on the colonies' surface using a spatula. The liquid which contained the spores was then taken using a Pasteur pipette and then collected into a sterile bottle. To determine the density of the spores in the suspension,

calculations were made using a hemacytometer.

The design used to determine the inhibition of the *R. oligosporus* culture filtrate on the growth of *A. flavus* FNCC 6109 on corn seeds was a Completely Randomized Design (CRD). Types of treatment of *R. oligosporus* filtrate on the growth of *A. flavus* FNCC6109 on corn seeds were as follows: a) Corn kernels without treatment (control), b) Corn kernels + 5 mL suspension of *A. flavus*, c) Corn kernels + *A. flavus* + 10% *R. oligosporus* filtrate, d) Corn kernels + *A. flavus* + 20% *R. oligosporus* filtrate, e) Corn kernels + *A. flavus* + 30% *R. oligosporus* filtrate, f) Corn kernels + *A. flavus* + 40% *R. oligosporus* filtrate, and g) Corn kernels + *A. flavus* + 50% (v/v) *R. oligosporus* filtrate.

The giving treatments were conducted by preparing as much as 100 g of corn kernels for each treatment which was placed in 8 sterile containers. The corn kernels were sprayed with as much as 15 mL of *R. oligosporus* filtrate and then adding 5 mL of *A. flavus* FNCC6109 spores were. After being given the treatment, they were stored for 15 days at 28°C. After the incubation period ended, the total population of *A. flavus* FNCC6109 was calculated using the plating method. To obtain the representative data, all treatments were repeated four times.

#### Data analysis

The quantitative data were analyzed with ANOVA. If the data obtained have a significant difference at the test level of 5% ( $P \leq 0.05$ ), it would be followed by the Duncan test to determine the differences in each treatment.

## Results

#### The isolation and identification of *R. oligosporus*

The macroscopic characteristics of the isolated *R. oligosporus* colonies in tempeh showed the color of white to the grayish, diameter of 4 cm in the four days incubation period, cottony texture, no zoning, no radial and growing zone. Whereas on the microscopic observation through 40 × 10 magnification on the *R. oligosporus* showed round or elliptical sporangium which had no insulation on the hyphae, there was a stolon that connected two sporangiophores and had rhizoid (Fig. 1). Macroscopic observations showed that *R. oligosporus* had a white to grayish colony color, textured like cotton, had no radial lines, had a growing zone, and had no zonation.

#### *In vitro* inhibition of *A. flavus* FNCC6109 by *R. oligosporus*

Inhibition of *R. oligosporus* against *A. flavus* FNCC6109 using the dual culture method has obtained the average colony area of *A. flavus* FNCC6109 as much as 5.77 ± 0.773 cm<sup>2</sup>. In comparison, it had as much as 13.00 ± 1,154 cm<sup>2</sup> of colony area in 4 days incubation under control. In

Table 1, it can be seen that the average percentage of inhibition power of *R. oligosporus* on the growth of *A. flavus* FNCC6109 was 56.08 ± 10.103% on PDA media with four days incubation period. The colony area of *A. flavus* FNCC6109 on treatments seemed smaller than those with no treatments (Fig. 2). There were allegations on the treatment of the *R. oligosporus* to be able to suppress the growth of *A. flavus* FNCC6109.

#### *In vitro* culture filtrate test of *R. oligosporus* against *A. flavus* FNCC6109

The inhibition of the culture filtrate of *R. oligosporus*, which tested against the growth of *A. flavus* FNCC6109 *in vitro* showed that the culture filtrate had the ability to inhibit *A. flavus* FNCC6109. In Fig. 3, it can be seen that the colonies of *A. flavus* FNCC6109 could not grow well in the presence of the culture filtrate of *R. oligosporus*, while on the control, colonies of *A. flavus* FNCC6109 seemed bigger.

In numbers, the inhibition test of the *R. oligosporus* culture filtrate against *A. flavus* FNCC6109 can be seen in Table 1. The five-day incubation period of the culture filtrate of *R. oligosporus* on PDA media had the highest percentage of inhibitory, which was 67.27 ± 2.70%, while the three days and four days incubation percentages were 61.26 ± 5.13% and 64.07 ± 7.04%. The culture filtrate of *R. oligosporus* showed positive results in inhibiting *A. flavus* FNCC6109 *in vitro*, presumably due to the presence of an active compound or enzyme capable of suppressing the growth of *A. flavus* FNCC6109. Meanwhile, *A. flavus* FNCC6109 grown on PDA media without culture filtrate suspension showed good growth of *A. flavus* FNCC6109 with a larger colony area.


#### The population of *A. flavus* FNCC6109 in corn kernels added with *R. oligosporus*

The corn kernels that gave the treatment of adding *R. oligosporus* culture filtrate had less population of *A. flavus* FNCC6109 colonies than the corn kernels that had not been given the culture filtrate. Table 2 represented several treatments of the culture filtrate concentration added to the corn kernels, which showed the varied population numbers of *A. flavus* FNCC6109. The calculation showed that at a concentration of 50%, the culture filtrate of *R. oligosporus* had the highest inhibitory ability with the average number of *A. flavus* FNCC6109 colonies  $3 \times 10^4$  CFU g<sup>-1</sup>, whereas in treatment B, which was only given a suspension of *A. flavus* FNCC6109 spores had a larger average population, namely  $42 \times 10^4$  CFU g<sup>-1</sup> after 15 days of the incubation period. This showed that the culture filtrate of *R. oligosporus* was able to suppress the population of *A. flavus* FNCC6109 in corn kernels.

Table 2 showed the lower the concentration of *R. oligosporus* culture filtrate added into the corn kernels, the

**Table 1:** Average and percentage of *R. oligosporus* inhibition against *A. flavus* FNCC6109 on PDA media at a certain incubation period at 28°C

Treatment	Inhibition (cm <sup>2</sup> or %)
Control (cm <sup>2</sup> )	13.00 ± 1.154
<i>R. oligosporus</i> (cm <sup>2</sup> )	5.77 ± 0.77
Not incubated (%)	56.08 ± 10.103
Three days incubation (%)	61.27 ± 5.13
Four days incubation (%)	64.07 ± 7.05
Five days incubation (%)	67.27 ± 2.70



**Fig. 1:** A. Colony morphology of *R. oligosporus* on PDA media with an incubation period of 3 days at a temperature of 28°C  
 B. Microscopic structure of *R. oligosporus* under a binocular microscope at 400× magnification (arrow S = Sporangium; SP = Sprongiophore; R = Rhizoid; Sn = Stolon)

higher the population of *A. flavus* FNCC6109 after an incubation period of 15 days. Whereas in treatment A (corn kernels without culture filtrate of *R. oligosporus* and *A. flavus* FNCC 6109), there were neither *R. oligosporus* nor *A. flavus* FNCC 6109 found. This proved that the corn kernels used in this study were free of contaminants from the two fungi.

## Discussion

Microscopically it forms non-septate hyphae, has stolons and rhizoids, and the shape of the sporangium is spherical. These characteristics were in accordance with the description stated by Yuliansih (2007) and Dolatabadi *et al.* (2014) in which *R. oligosporus* macroscopically and microscopically have white to grayish colonies, have rhizoid like roots, the hyphae are not insulated, single sporangiophores, the sporangium is round or elliptical and have stolons. The difference between the genus of *Rhizopus* and other fungi is that the non-insulated hyphae have rhizoids and a distinctive sporangium shape. Firmansyah (2007) had also isolated and identified *R. oligosporu* spp. in tempeh. The results obtained three species those were *R. oligosporus*, *R. stolonifer* and *R. oryzae*. The same research had been conducted by Virgianti (2015), which showed that the isolated *R. oligosporus* colonies in tempeh macroscopically had the characteristics of having white to grayish color and grew like cotton. Rahmawati *et al.* (2013) reported the results of the isolation and identification of

molds in corns and found the presence of *R. oryzae* and *R. stolonifer*. Furthermore, McKelvey and Murphy (2017) stated the existence of *R. oligosporus* in cornflour had a role in producing cellulose, xylanase, and protease enzyme activities.


The results of research by Nursadin and Supriyanto (2012) showed that *R. oligosporus* was able to inhibit *Fusarium oxysporum* by 60%. Furthermore, it was conveyed that *R. oligosporus* could be used as a competitor because it has a very high competitive ability and very fast growth. The same thing was also conveyed by Anuragi and Sharma (2016) that *R. oligosporus* could be used as a biocontrol agent because of its ability to inhibit *Fusarium oxysporum* with an inhibitory percentage of 56.76%. Moreover, Adebola and Amadi (2010) reported that *R. oligosporus* could inhibit the growth of the pathogen *Phytophthora palmivora* with a 76% inhibition rate in 7 days incubation period. The inhibition mechanism of *R. oligosporus* against the growth of *A. flavus* FNCC6109 other than by suppressing it was suspected that *R. oligosporus* produced metabolites that could inhibit *A. flavus* FNCC6109. *R. oligosporus* was also able to produce enzymes and bioactive compounds that are antimicrobial and antifungal. Virgianti (2015) reported that isolated *R. oligosporus* from local tempeh was capable of producing antimicrobial bioactive compounds. The bioactive compounds produced were able to inhibit enteric pathogenic bacteria and have different inhibition zones.




**Table 2:** The population of *A. flavus* FNCC6109 colonies in corn kernels added with culture filtrate of *R. oligosporus* before and after the incubation period

Treatment	Total Population of <i>A. flavus</i> FNCC6109 (CFU g <sup>-1</sup> )		Enhancement of <i>A. flavus</i> FNCC6109 (%)
	Population Before the Incubation (T <sub>0</sub> )	Population After the Incubation (T <sub>15</sub> )	
A	0.00	0.00 <sup>a</sup> ± 0.00	0
B	21 × 10 <sup>4</sup>	44.7 × 10 <sup>4b</sup> ± 0.068 × 10 <sup>4</sup>	53
C	14 × 10 <sup>4</sup>	18.0 × 10 <sup>4c</sup> ± 0.062 × 10 <sup>4</sup>	22.2
D	11 × 10 <sup>4</sup>	13.7 × 10 <sup>4cd</sup> ± 0.051 × 10 <sup>4</sup>	19.7
E	9 × 10 <sup>4</sup>	10.3 × 10 <sup>4de</sup> ± 0.134 × 10 <sup>4</sup>	14.4
F	6 × 10 <sup>4</sup>	6.7 × 10 <sup>4df</sup> ± 0.127 × 10 <sup>4</sup>	10.4
G	5 × 10 <sup>4</sup>	5.3 × 10 <sup>4f</sup> ± 0.196 × 10 <sup>4</sup>	5.7

T<sub>15</sub> is the average value with three replications and with different letter notations in the same column indicating a significantly different average value ( $P \leq 0.05$ ) based on the Duncan test after conducting the analysis of variance



**Fig. 2:** Inhibition of *R. oligosporus* against *A. flavus* FNCC6109 on PDA media with an incubation period of 4 days and at a temperature of 28°C (K = Control; P = Treatment; A = *A. flavus* FNCC6109; R = *R. oligosporus*)



**Fig. 3:** Inhibition of *R. oligosporus* culture filtrate against *A. flavus* FNCC6109 *in vitro* on PDA media with an incubation period of 4 days and at a temperature of 28°C (Information on the figure KF = filtrate culture *R. oligosporus*.; A = *A. flavus* FNCC6109)

The culture filtrate of *R. oligosporus* at five days incubation period had the highest inhibitory ability in inhibiting *A. flavus* FNCC6109. During the five days incubation period, the enzymes produced by *R. oligosporus* were thought to have optimum enzyme activity. This is supported by the statement of Pujiati *et al.* (2017) that the levels of crude protein cellulase enzymes from *R. oligosporus* in sugarcane bagasse substrate incubated for 3 to 12 days had an increase in the amount of protein content. The increase in protein content also indicated the activity of the cellulase enzyme has increased. This happened due to the increasing length of fermentation time. *R. oligosporus* was able to produce  $\beta$ -glucanase enzymes. The  $\beta$ -glucanase

enzyme is an extracellular enzyme capable of hydrolyzing carbohydrates of the glucan group. This meets the statement of Ravindran *et al.* (2018) that *R. oligosporus* is capable of producing  $\beta$ -glucanase enzymes. Glucans are one important component in making the cell walls of fungi in general. Glucans can be hydrolyzed by the  $\beta$ -glucanase enzyme produced by several types of fungi, namely *R. oligosporus* and *Trichoderma* spp. The  $\beta$ -glucanase enzyme produced also has an important role in the self-defense mechanism against pathogenic fungal attacks. Research conducted by Lorito *et al.* (1994) stated that the  $\beta$ -glucanase enzyme showed antifungal activity by hydrolyzing the glycan structures present in the cell walls of pathogenic fungi. The

structure of glucans is known to be mostly found at the tip of the hyphae, so the pathogenic fungal hyphae are not able to grow properly in the presence of the  $\beta$ -glucanase enzyme. Furthermore, it was also confirmed by Budiarti and Widyastuti (2011) that the  $\beta$ -glucanase enzyme was able to influence the growth of hyphae, where the hyphae experienced swelling and necrosis.

Based on the statistical test, it showed the effect of the concentration treatment of *R. oligosporus* culture filtrate, which was given into the corn kernels during an incubation period of 15 days against *A. flavus* FNCC6109 in corn kernels. The average population of *flavus* FNCC6109 in treatment B (without culture filtrate) was  $44.7 \times 10^4$  CFU g<sup>-1</sup>, which was significantly different ( $P \leq 0.05$ ) with treatment G (corn kernels with *A. flavus* and culture filtrate of *R. oligosporus* with a concentration of 50% (v/v). The concentration treatment of *R. oligosporus* culture filtrate given in corn kernels was able to decrease the population of *A. flavus* FNCC6109. It was proven by the decreasing tendency in the population of *A. flavus* FNCC6109; the higher the concentration of the culture filtrates given, the lower the population of *A. flavus* FNCC6109. This is thought to be *R. oligosporus* experiencing very fast growth and ability to compete in obtaining nutrients. In addition, it is also suspected that the performance of enzymes and other metabolites played a role in damaging the spore wall components of *A. flavus* FNCC6109 so that the spores of *A. flavus* FNCC6109 were not able to grow properly. This statement is supported by Calestino *et al.* (2006), who stated that *R. oligosporus* is able to produce  $\beta$ -glucanase enzyme, which is thought to be able to damage the spore walls of *A. flavus* FNCC6109. Furthermore, *R. oligosporus* is able to produce volatile compounds, which are thought to affect in inhibiting the growth of *A. flavus* FNCC6109. This is supported by the statement of Huang *et al.* (2019) that *R. oligosporus* is able to produce volatile compounds such as ethanol, isobutyl alcohol, and three methyl butanol, where these compounds are thought to be able to inhibit the growth of *A. flavus*. Similar results were stated by Gunaser *et al.* (2017) that *R. oligosporus* produces volatile compounds. The isolation results of the volatile compounds by *R. oligosporus* obtained in fermented soybeans and barley, volatile compounds, namely ethanol, acetone ethyl acetate, 2-butanone, 2-methyl-1-butanol.

Based on the presumable performance of the enzymes and bioactive compounds produced by *R. oligosporus*, it can be attempted that *R. oligosporus* could be used as a biocontrol agent in inhibiting the growth of pathogenic fungi. This statement is supported by Monk *et al.* (2020) that the inhibition mechanism of *R. oligosporu* spp. against *A. flavus* and *A. parasiticus* by binding to element C (Carbon) contained in beans, causing *A. flavus* and *A. parasiticus* to lose. The utilization of *R. oligosporus* culture filtrate has been conducted in many industries and can be used as a probiotic agent. Besides its ability to be a

biocontrol in inhibiting the growth of pathogenic fungi and degrading mycotoxin compounds, *R. oligosporus* is thought to increase the quality of food quality as reported by Maryana *et al.* (2016), which stated that *R. oryzae* culture could increase the protein content in liquid tofu waste. The results obtained during fermentation for 48 h could increase the protein content by 0, 47%. Suarti and Budijanto (2021) stated that the use of *R. oligosporus* in concentrate feed with fermentation process was able to increase weight and reduce phytic acid levels. Phytic acid is a phosphorus compound that can bind mineral components such as iron, calcium, and zinc so that it cannot be absorbed directly by the body. According to Bhavsar and Khire (2014) that giving *R. oligosporus* culture filtrate in concentrate, feed is thought to be able to produce phytase enzymes that affect in breaking down the phytic acid.

Based on the ability of the culture filtrate tested on corn kernels, it generally had a positive correlation, both *in vitro* and *in vivo*, in inhibiting the growth of *A. flavus* FNCC6109. So it can be seen that the culture filtrate of *R. oligosporus* has the potential in inhibiting the growth of *A. flavus* FNCC6109 in corn kernels with a decrease in the population of *A. flavus* FNCC6109 after an incubation period of 15 days.

## Conclusion

The *R. oligosporus* filtrate inhibited *A. flavus* FNCC6109 with the highest percentage of inhibition was 61.92, and the lowest was 44.42. The highest inhibition rate was  $67.27 \pm 2.70$  with an incubation period of 5 days, and the lowest was  $61.27 \pm 5.13$  with an incubation period of 3 days. The filtrate concentration of 50% suppresses *A. flavus* FNCC6109 with the lowest population average of  $55.3 \times 10^4 \pm 0.196$  CFU g<sup>-1</sup> after an incubation period of 15 days. Further research needs to be conducted on the enzymes and active compounds produced by *R. oligosporus*, as well as testing the decrease in aflatoxin content produced by *A. flavus* FNCC6109 in corn kernels.

## Acknowledgments

The authors would like to thank the Head of the Microbiology Laboratory, Department of Biology, Faculty of Mathematics and Natural Sciences, and the Joint Laboratory of the Faculty of Mathematics and Natural Sciences, Udayana University.

## Author Contributions

IBGD, as the head of the research, was in charge of designing research and isolation of *R. oligosporus* and *A. flavus* FNCC6109; AAKD and Arofi were in charge of preparing tools, media, and conducting dual culture tests; IWS performed fungal growth measurements and tabulated data; and IKW performs data analysis, translation, and publication.

## Conflicts of Interest

The authors declare no conflict of interest

## Data Availability

Data presented in this study will be available on a fair request to the corresponding author.

## Ethics Approval

Not applicable in this paper

## References

- Adebola MO, JE Amadi (2010). Antagonistic activities of *Paecilomyces* and *Rhizopus* species against the cocoa black pod pathogen (*Phytophthora palmivora*). *Afr Sci* 11:235–239
- Ali A, A Javaid, A Shoaib, IH Khan (2020). Effect of soil amendment with *Chenopodium album* dry biomass and two *Trichoderma* species on growth of chickpea var. Noor 2009 in *Sclerotium rolfsii* contaminated soil. *Egypt J Biol Pest Cont* 30:1–9
- Anuragi M, TK Sharma (2016). Biocontrol of chickpea wilt disease by *Fusarium oxysporum* f. spp. *Ciceri* with *Rhizosphere mycoflora*. *Flora Fauna* 22:201–209
- Bhavsar K, JM Khire (2014). Current research and future perspectives of phytase bioprocessing. *RSC Adv* 4:26677–26691
- Budiarti SW, SM Widyastuti (2011). The effect of antifungal activity of  $\beta$ -1, 3-glucanase *Trichoderma reesei* on root fungi of *Ganoderma philippii*. *J Widya Res* 14:455–460
- CBS – Central Bureau of Statistics (2014). *Bali in Numbers*. Statistical Yearbook of Indonesia (in Bahasa), Indonesia
- Celestino K, R Cunha, C Felix (2006). Characterization of a  $\beta$ -glucanase produced by *Rhizopus microsporus* var. *microsporus*, and its potential for application in the brewing industry. *BMC Biochem* 7:23
- Dolatabadi S, G Walther, AHGGVD Ende, GSD Hoog (2014). Diversity and delimitation of *Rhizopus microsporus*. *Fung Div* 64:145–163
- Firmansyah R (2007). *Isolation, Identification and Production of Mycelia Rhizopus spp. Low Nucleic Acid Levels*. Essay Biology Department, FMIPA Bogor Agricultural Institute, Bogor, Indonesia
- Guneser O, A Demirkol, YK Yuceer, SO Togay, MI Hosoglu, M Elibol (2017). Production of flavor compounds from olive mill waste by *Rhizopus oryzae* and *Candida tropicalis*. *Braz J Microbiol* 48:275–285
- Huang ZR, WL Guo, WB Zhou, L Li, JX Xu, JL Hong, XC Lv (2019). Microbial communities and volatile metabolites in different traditional fermentation starters used for hong qu glutinous rice wine. *Food Res Intl* 121:593–603
- Khan IH, A Javaid (2022a). Antagonistic activity of *Aspergillus versicolor* against *Macrophomina phaseolina*. *Braz J Microbiol* 53:1–9
- Khan IH, A Javaid (2022b). Biocontrol *Aspergillus* species together with plant biomass alter histochemical characteristics in diseased mungbean plants. *Microsc Res Tech* 85:2953–2964
- Khan IH, A Javaid (2022c). DNA cleavage of the fungal pathogen and production of antifungal compounds are the possible mechanisms of action of biocontrol agent *Penicillium italicum* against *Macrophomina phaseolina*. *Mycologia* 114:24–34
- Lorito M, CK Hayes, AD Pietro, SL Woo, GE Harman (1994). Purification, characterization, and synergistic activity of a glucan 1, 3- $\beta$ -glucosidase and *n*-acetyl- $\beta$ -glucosaminidase from *Trichoderma harzianum*. *J Phytopathol* 84:398–405
- Maryana L, S Anam, AW Nugrahani (2016). Production of single cell protein from *Rhizopus oryzae* culture with tofu liquid waste medium. *J Pharm* 2:132–137
- Mauro A, E Garcia-Cela, A Pietri, PJ Cott, P Battilani (2018). Biological control products for aflatoxin prevention in Italy: Commercial field evaluation of atoxigenic *Aspergillus flavus* active ingredients. *Toxins* 10:30–43
- McKelvey SM, RA Murphy (2017). Biotechnological use of fungal enzymes. *Fungi Biology and Applications*, pp:201–225. Kavanagh K (Ed.). John Wiley & Sons, Inc., Hoboken, New Jersey, USA
- Monk BC, AA Sagatova, P Hosseini, YN Ruma, RK Wilson, MV Keniya (2020). Fungal lanosterol 14 $\alpha$ -demethylase: a target for next-generation antifungal design. *Biochim Biophys Acta Prot Proteom* 1868:140206
- Montalbano S, F Degola, J Bartoli, F Bisceglie, A Buschini, M Carcelli, D Feretti, S Galati, L Marchi, N Orsoni, G Pelosi, M Pioli, FM Restivo, D Rogolino, M Scaccaglia, O Serra, G Spadola, GCV Viola, I Zerbini, C Zani (2021). The AFLATOX@ Project: Approaching the development of new generation, natural-based compounds for the containment of the mycotoxigenic phytopathogen *Aspergillus flavus* and aflatoxin contamination. *Intl J Mol Sci* 22:4520–4538
- Nursadin IS, Supriyanto (2012). Screening of lignocellulolytic acidophilic antagonist fungi and peat to *Fusarium* wilt. *J Plant Trop Land* 2:27–34
- Pitt JI, AD Hocking (1997). *Fungi and Food Spoilage*. Printed in Great Britain at the University Press, Cambridge, UK
- Pujati A, Sulistyarsi, MW Ardhi (2017). Analysis of cellulase enzyme crude protein levels from *Rhizopus* spp. on sugarcane bagasse substrate isolated from clove gardens Madiun. *J Biota* 3:26–30
- Rahayu ES, S Raharjo, AA Rahmianna (2010). Aflatoxin contamination in maize production in East Java. *J Tech Agric* 8:1–16
- Rahmawati, RD Hariyadi, P Hariyadi, D Fardiaz, N Richana (2013). Isolation and identification of microorganisms during spontaneous fermentation of maize. *J Tech Food Ind* 24:33–39
- Ravindran R, SS Hassan, GA Williams, AK Jaiswal (2018). A review on bioconversion of agro-industrial wastes to industrially important enzymes. *Bioengineering* 5:93–122
- Sharf W, A Javaid, A Shoaib, IH Khan (2021). Induction of resistance in chili against *Sclerotium rolfsii* by plant growth promoting rhizobacteria and *Anagallis arvensis*. *Egypt J Biol Pest Cont* 31:1–11
- Suarti B, S Budijanto (2021). Bio-active compounds, their antioxidant activities, and the physicochemical and pasting properties of both pigmented and non-pigmented fermented de-husked rice flour. *AIMS Agric Food* 6:49–65
- Singh PK, K Vijay (2011). Biological control of *Fusarium* with of *Chrysanthemum* with *Trichoderma* and botanicals. *J Agric Technol* 7:1603–1613
- Virgianti DA (2015). Antagonist test of tempe mushroom (*Rhizopus* spp.) against enteric pathogenic bacteria. *J Biosfera* 32:163–168
- Yuliansih RR (2007). *Effect of Drying Temperature of Acid Soybeans on the Quality of Tempe Prepared as Tempe Kit*. Essay. FTP. UGM. Yogyakarta, Indonesia