

**Running title: Improving Stress Tolerance in Wheat Using Thiourea  
Potential of Foliar Applied Thiourea in Improving Salt and High Temperature Tolerance of  
Bread Wheat (*Triticum aestivum*)**

**Freeha Anjum<sup>1</sup>, Abdul Wahid<sup>2\*</sup>, Muhammad Farooq<sup>3</sup> and Shahzad M.A. Basra<sup>4</sup>**

<sup>1</sup>*Biology Department, Government Islamia College for Women, Faisalabad, Pakistan*

<sup>2</sup>*Department of Botany, University of Agriculture, Faisalabad-38040, Pakistan*

<sup>3</sup>*Department of Agronomy, University of Agriculture, Faisalabad-38040, Pakistan*

<sup>4</sup>*Department of Crop Physiology, University of Agriculture, Faisalabad-38040, Pakistan*

\*For correspondence: drawahid2001@yahoo.com

### **Novelty statement**

Thiourea has been rarely exploited for improving wheat growth under stress conditions. These results show that foliar application of thiourea to wheat was effective in substantially improving root growth at young stage while economic yield at maturity both under high temperature and salinity stresses. We found that improved root growth with foliar-applied thiourea was a major factor in bringing about these changes.

### **Abstract**

In this research, we investigated the possible role of foliar spray of 10 mM thiourea in improving the salinity tolerance threshold ( $EC_e$ ) and high temperature tolerance of five wheat (*Triticum aestivum* L.) varieties at seedling, pre-anthesis and grain filling stages of growth. Data were recorded for changes in length and dry weight of shoot and root and leaf area at seedling and pre-anthesis stages, while yield and yield components were recorded at maturity. Results revealed great varietal differences in the salinity and high temperature tolerance at all growth stages. Foliar applied thiourea improved  $EC_e$  (6–11%) and high temperature tolerance (4–10%) in these varieties at different stages. Correlations of growth and yield attributes substantiated the effectiveness of foliar spray of thiourea in promoting root growth at seedling and pre-anthesis stages, and grain filling and harvest index at maturity. Among the varieties, thiourea was more effective in enhancing high temperature tolerance of the most high temperature sensitive (S-24) and  $EC_e$  of most salt sensitive (MH-97) varieties. In crux, foliar spray with thiourea has a great potential to mitigate the adverse effects of salinity and high temperature and may be recommended for field use in the areas, where moderate salinity or high temperature limit wheat production.

**Keywords:** Correlations; Foliar spray; Grain yield; Root growth; Stress tolerance; Wheat

### **Introduction**

Increased soil salinity and ambient temperature are amongst the major factors affecting plant productivity. Tolerance to salinity and high temperature involve the simultaneous expression of a number of genes, leading to varied synthesis of metabolites. Both these stresses have some common adversaries on plants, such as altered morphological development, reductions in gas exchange attributes, assimilate partitioning, altered patterns of protein synthesis, changes in the activities of antioxidants and disruption of membrane functions (Zhu, 2002; Sairam and Tyagi, 2004; Pandey *et al.*, 2011; Taiz and Zeiger, 2006). Salinity of soils is increasing rapidly all over the world because of irrigation of crops with low quality water, thus leading to net accumulation of ions in the root zone (Murillo-Amador *et al.*, 2002; Pitman and Lauchli, 2002). High temperature damages field grown crops and limits growth and yield by acting as a dehydrative force and aggravates the prevailing stress conditions (Wahid *et al.*, 2007).

The plant stress tolerance can be improved with the exogenous use of stress alleviating chemicals (Wahid and Shabbir, 2005; Farooq *et al.*, 2009a, b). Among the stress alleviating compounds, thiourea is an important molecule with two functional groups; ‘thiol’ is important to oxidative stress response and ‘imino’ partly fulfils the N requirement. It is highly water soluble and easily absorbed in the living tissues. Although role of thiourea in abiotic stress tolerance is scarcely investigated, available reports show that it relieves salinity induced seed dormancy in *Allenrolfea occidentalis* at lower concentration (Gul and Weber, 1998). Under conditions of water stress or high temperature, external use of thiourea can increase K<sup>+</sup> uptake by chickpea and reduce ABA biosynthesis (Aldasoro *et al.*, 1981). Foliar application of thiourea improved net photosynthesis and chlorophyll content in drought stressed clusterbean (*Cyamopsis tetragonoloba*) plants (Garg *et al.*, 2006). Recently, Srivastava *et al.* (2010) reported that thiourea treatment coordinately bioregulates different signaling and effector mechanisms and alleviates the adverse effects of high salinity in salt-treated *Brassica juncea* seeds.

Wheat (*Triticum aestivum* L.) has been ranked as tolerant to salinity with a salinity tolerance threshold (EC<sub>e</sub>) value between 6 and 8.6 dS m<sup>-1</sup> (Carter, 1981; Farooq *et al.*, 2009a). While being a winter crop, it is also sensitive to episodes of high temperature (Yang *et al.*, 2002). In view of the changing climatic conditions mainly related to global warming, high temperature aggravates the salinity status of the field at crucial stages like germination and reproductive growth periods, leading to hampered field stand in the former case and diminished final grain yield in the

latter case. We predict that foliar application of thiourea solution can enhance  $EC_e$  and high temperature tolerance of wheat. Thus, the objectives of this study were to explore the tendency of foliar-applied thiourea in improving  $EC_e$  and heat tolerance and sorting out some mechanism of thiourea action in salinity and high temperature stress tolerance of wheat at three growth stages.

## Materials and Methods

### Experimental Details and Treatments

**Experimental material:** Experiments were performed in the net-house of the Department of Botany, University of Agriculture, Faisalabad, Pakistan during November-April of the years 2004-2005 and 2005-2006. Seed of wheat (*Triticum aestivum* L.) varieties, Inqlab-91, MH-97, BK-2002 and SH-2002 was obtained from Wheat Breeder, Ayub Agricultural Research Institute (AARI), Faisalabad, while that of S-24 from Department of Botany. Six seeds of each variety were directly sown in glazed pots (45 cm high with 25 cm diameter from the neck and 18 cm from bottom) containing thoroughly washed sand (20 kg) and kept in the net-house. After germination, three uniform seedlings were maintained in each pot. Plants were supplemented with half-strength nutrient solution (Hoagland and Arnon, 1950) throughout their growth periods.

**Treatments:** For the determination of salinity tolerance threshold ( $EC_e$ ), wheat varieties were exposed to 0, 60, 80, 100 and 120 mM levels of NaCl at the rate of 20 mM NaCl per day separately at seedling (20 day old plants), pre-anthesis (60 day old plants) and grain filling stages (80 day old plants). For high temperature environment, plants were shifted to open door plexiglass fitted canopies with a light transmission index of about 0.8. The canopies had 7–10°C higher temperature as compared to ambient (Fig. 1). Average photosynthetically active radiations on the leaf surface ranged between 855 to 975  $\mu\text{mol m}^{-2} \text{s}^{-1}$  (measured with Infra Red Gas Analyzer, LCA-4, Analytical Development Company, Hoddesdon, England) and relative humidity 55–60% and 35–40% during winter and summer months respectively. Pots were protected from rainfall by covering with transparent plastic sheet. Plants from all the treatments were divided into three groups, which were foliar sprayed with none (group-I), sprayed with distilled water (group-II) and sprayed with 10 mM thiourea solution (group-III) were sprayed with a sprayer. Effective concentration, 10 mM,

of thiourea for foliar spray was pre-optimized after conducting series of experiments (data not shown).

### **Plant Harvesting**

The plants were harvested 15 days after applying treatment at each stage. At seedling stage, the determinations were made for length and dry weight of shoot and root and leaf area per plant. At pre-anthesis stage, in addition to measurements made at seedling stage, flag leaf area was also measured. At both these stages total and flag leaf areas were determined of intact plants as maximum leaf length and maximum leaf width  $\times$  0.68 (correction factor). Dry weight was determined after drying the harvested plants at 70°C for seven days. The plants receiving high temperature and salinity stresses at grain filling stage were harvested at full maturity to determine grain yield and yield components. The seeds were extracted from the spikes to determine grain yield per plant.

### **Assessment of Stress Tolerance**

Salinity tolerance threshold ( $EC_e$ ) values of wheat varieties were determined with the equation of Maas and Hoffman (1977):

$$Y_r = 100 - B (EC_e - A)$$

Where  $Y_r$  is the relative yield; slope  $B$  represents the decrease in yield and threshold  $A$  is the lowest salinity ( $EC_e$  in  $dS\ m^{-1}$ ), which significantly reduces yield. Net change in the  $EC_e$  due to foliar thiourea application was calculated in percentage based on the individual characters, as well as average for each variety computed.

For determination of any change in the high temperature tolerance with or without thiourea application, the values were computed as percent change in individual growth and yield attributes over respective controls and averages calculated for each variety. Net change was computed as difference of thiourea-sprayed versus non-sprayed treatments.

### **Statistical Analysis**

In the absence of any striking differences in the plants from group-I and group-II (not sprayed and sprayed with water, respectively), the comparison and derivations were made between the group-II and group-III plants (foliar sprayed with water and 10 mM thiourea solution). Design of the experiments was completely randomized with three replications. The data for both the years was averaged to determine comparative responses of varieties in terms of changes in  $EC_e$  and high temperature tolerance separately with or without thiourea spray. Data were statistically analyzed using COSTAT computer software (COHORT, Monterey, California).

## Results

### Changes in Stress Tolerance with Thiourea

**Salinity tolerance:** Wheat varieties indicated significant differences for  $EC_e$  and improvement in it after foliar spray with thiourea at seedling ( $P < 0.05$ ), pre-anthesis ( $P < 0.01$ ) and maturity ( $P < 0.05$ ) stages. At seedling stage, S-24 showed  $EC_e$  of 8.55, followed by BK-2002 and SH-2002 (7.76 and 7.73, respectively), whilst Inqlab-91 (7.10) and MH-97 (6.85) were the most sensitive. Although enhanced in all the varieties with thiourea, maximum improvement in  $EC_e$  was recorded in SH-2002, BK-2002, MH-97 (9.67, 9.60 and 9.09%, respectively) followed by Inqlab-91 (8.45%), whereas S-24 (7.42%) displayed the minimum  $EC_e$  (Table 1).

At pre-anthesis stage, S-24 indicated the greatest average  $EC_e$  ( $9.66 \text{ dS m}^{-1}$ ) followed by SH-2002, BK-2002 and Inqlab-91, while MH-97 fared very poorly, displaying  $EC_e$  of  $6.63 \text{ dS m}^{-1}$ . Foliar applied thiourea although improved  $EC_e$  in all the varieties, BK-2002 excelled the other varieties by showing an enhancement in  $EC_e$  of 11.16%, followed by MH-97 and Inqlab-2002 (Table 1). At maturity stage,  $EC_e$  assessed on the basis of grain yield and yield components revealed that except MH-97, showing the lowest  $EC_e$ , all the varieties indicated improved  $EC_e$  but did not differ ( $P > 0.05$ ) from each other. Foliar spray with thiourea improved the  $EC_e$  of all the varieties, and the maximum enhancement was noted in SH-2002 and MH-97, followed by S-24 and BK-2002, while Inqlab-91 was at the bottom (Table 1).

**High temperature tolerance:** All the varieties indicated a decline in growth and yield attributes at prevailing high temperature, but foliar applied thiourea significantly ( $P < 0.01$ ) improved the high

temperature tolerance at seedling, pre-anthesis and maturity stages. A comparison of varieties showed that S-24 and BK-2002 indicated greatest decline in the growth attributes followed by SH-2002, MH-97 and Inqlab-91. Foliar application of 10 mM thiourea improved growth of all the varieties under high temperature and brought about net improvement in the order of BK-2002 > S-24 > MH-97 > SH-2002 > Inqlab-91 (Table 2).

At pre-anthesis stage, the wheat varieties indicated substantial reduction in high temperature tolerance over respective controls and S-24 was the most affected showing an average decline of ~22%, while MH-97 was minimally influenced (~12%). Foliar application of thiourea curtailed the reductions in growth attributes and as a result the varieties indicating average net improvement were in the rank: S-24 (9.48) > MH-97 (6.53%) > BK-2002 (6.50%) > BK-2004 (6.38%) > SH-2002 (5.12%), respectively (Table 2). Towards maturity, data recorded for changes in grain yield and yield components due to high temperature revealed that S-24 underwent a greatest reduction (22.86%) followed by BK-2002 (17.31%), Inqlab-91 (16.72%), SH-2002 (16.24%), while MH-97 manifested a lowest reduction (10.47%). Foliar spray with thiourea produced a net improvement in high temperature tolerance of these varieties in the orders: Inqlab-91 > S-24 > SH-2002 > BK-2002 > MH-97, respectively (Table 2).

## Discussion

Although some reports show improvement in plant growth by exogenous use of thiourea (Burman *et al.*, 2004; Garg *et al.*, 2006), none is available on the enhancement in the EC<sub>e</sub> and high temperature tolerance in thiourea-sprayed plants. In this study, foliar use of thiourea improved the EC<sub>e</sub> and high temperature tolerance at three growth stages, but the extent of improvement was differential for the stress treatments as well as varieties at three growth stages. Foliar applied thiourea was highly more effective in improving the EC<sub>e</sub> of salt sensitive varieties (MH-97, BK-2002 and SH-2002) at seedling, pre-anthesis and maturity (Table 1). Likewise, S-24 being less tolerant of high temperature indicated substantially improved growth and yield after foliar spray with thiourea at all phenological stages (Table 2).

Results of this study indicated that improved EC<sub>e</sub> and high temperature tolerance of thiourea-applied plants was primarily linked to improved growth and yield parameters (see Materials and methods section), on the basis of which the data presented in Table 1 and 2 were

derived. This implied that the primary action of thiourea is to improve net assimilation efficiency of plant by alleviating salinity and high temperature damage to photosynthetic area, which is crucial for stress tolerance (James *et al.*, 2002; Wahid, 2007). Improved leaf area, particularly flag leaf area with thiourea application proved important for providing greater surface area for the production and partitioning of photoassimilates towards vegetative and reproductive growth (Gibberd *et al.*, 2002). It is noteworthy that such improvements were better discernible in case of sensitive varieties, irrespective of the prevailing stress condition, which definitely signifies the tendency of foliar use of thiourea in improving stress tolerance of wheat.

Both salinity and high temperature have a common facet of oxidative damage (Desikan *et al.*, 2005; Wahid *et al.*, 2007). Looking at the structure of thiourea, both ‘imino’ and ‘thiol’ functional groups have great implications in abiotic stress tolerance. With foliar spray imino group provides a ready source of nitrogen and thiol has a great role in alleviating oxidative stress damage on the physiologically more important mesophyll tissue. It was evident from the results that foliar applied thiourea was effective in improving salt- (6–11%) and high-temperature tolerance (4–10%) of wheat varieties due to its abiotic stress-mitigating properties. Prolific root system is important to stress tolerance and improved final yield (Flowers, 2004; Farooq *et al.*, 2009b). In line with this, a pronounced increase in root growth of salt- and high temperature-sensitive varieties was directly associated to grain yield, which is an important manifestation of thiourea action in improving wheat growth of more sensitive varieties at various growth stages. Thus, in view of being water soluble, readily absorbable in the tissues and ability to ameliorate stress effects, field use of thiourea is beneficial in achieving better wheat yield from relatively warm and saline areas.

## **Conclusion**

Although with substantial varietal difference, thiourea can be effectively used to produce salinity and high temperature tolerance in wheat. Further studies are needed to find the possible mechanism(s) of the growth improvement as noted in the present study.

## **Acknowledgements**


The first author acknowledges the financial grant from University of Agriculture, Faisalabad, Pakistan

## References

- Aldasoro, J.J., A. Matilla and G. Nicolás, 1981. Effect of ABA, fusicoccin and thiourea on germination and K<sup>+</sup> and glucose uptake in chickpea seeds at different temperatures. *Physiol. Plant.*, 53: 139–145
- Burman, U., B.K. Garg and S. Kathju, 2004. Interactive effects of thiourea and phosphorus. *Biol. Plant.*, 48: 61–65
- Carter, D.L., 1981. *Salinity and Plant Productivity*, Vol. 1, pp: 117–133. CRC Handbook of Agricultural Productivity, CRC press, New York, USA
- Desikan, R., K. Hancock and S. Neill, 2005. Reactive oxygen species as signalling molecules. *In: Antioxidants and Reactive Oxygen Species in Plants*, pp: 169–196. N. Smirnoff (ed). Blackwell Publishing, New Delhi, India
- Farooq, M., N. Kobayashi, A. Wahid, O. Ito and S.M.A. Basra, 2009a. Strategies for producing more rice with less water. *Adv. Agron.*, 101: 351–387
- Farooq, M., A. Wahid, N. Kobayashi, D. Fujita and S.M.A. Basra, 2009b. Plant drought stress: effects, mechanisms and management. *Agron. Sustain. Dev.*, 28: 185–212
- Flowers, T.J., 2004. Improving crop salt tolerance. *J. Exp. Bot.*, 55: 307–319
- Garg, B.K., U. Burman and S. Kathju, 2006. Influence of thiourea on photosynthesis, nitrogen metabolism and yield of clusterbean (*Cyamopsis tetragonoloba* (L.) Taub.) under rainfed conditions of Indian arid zone. *Plant Growth Regul.*, 48: 237–245
- Gibberd, M.R., N.C. Turner and R. Storey, 2002. Influence of saline irrigation on growth, ion accumulation and partitioning and leaf gas exchange of carrot (*Daucus carota* L.). *Ann. Bot.*, 90: 715–724
- Gul, B. and D.J. Weber, 1998. Effect of dormancy relieving compounds on the seed germination of non-dormant *Allenrolfea occidentalis* under salinity stress. *Ann. Bot.*, 82: 555–560
- Hoagland, D.R. and D.I. Arnon, 1950. *The Water Culture Method for Growing Plants Without Soil*. Circular 347, University of California, College of Agriculture, Berkeley, USA
- James, R.A., A.R. Rivelli, R. Munns and S. Von Caemmerer, 2002. Factors affecting CO<sub>2</sub> assimilation, leaf injury and growth in salt stressed durum wheat. *Funct. Plant Biol.*, 29: 1393–1403

- Maas, E.V. and J.G. Hoffman, 1977. Crop salt tolerance—current assessment. *J. Irrig. Drain. Div. Amer. Soc. Civil Eng.*, 103: 1309–1313
- Murillo-Amador, B., E. Troyo-Diequez, R. Lopez-Aguilar, A. Lopez-Cortes, C.L. Tinoco-Ojanguri, H.G. Jones and C. Kaya, 2002. Matching physiological traits and ion concentrations associated with salt tolerance with cowpea genotypes. *Aust. J. Agric. Res.*, 53: 1243–1255
- Pandey, R.K., D. Kumar and K.M. Jadhav, 2011. Assessment of determinants for reducing HCN content in Sorghum used for ruminant in Gujarat, India. Available at: <http://www.lrrd.org/lrrd23/3/pand23066.htm> (Accessed: 23 March 2012)
- Pitman, M.G. and A. Lauchli, 2002. Global impact of salinity and agricultural ecosystems. *In: Salinity: Environment–Plants–Molecules*, pp: 3–20. A. Lauchli and U. Luttge (eds.). Kluwer Academic Publishers, Dordrecht, The Netherlands
- Sairam, R.K. and A. Tyagi, 2004. Physiology and molecular biology of salinity stress tolerance in plants. *Curr. Sci.*, 86: 407–421
- Srivastava, A.K., N.K. Ramaswamy, P. Suprasanna and S. F. D'Souza, 2010. Genome-wide analysis of thiourea-modulated salinity stress-responsive transcripts in seeds of *Brassica juncea*: identification of signalling and effector components of stress tolerance. *Ann. Bot.*, 106: 663–674
- Taiz, L. and E. Zeiger, 2006. *Plant Physiology*, 4<sup>th</sup> edition. Sinauer Associates Inc., Publishers, Sunderland, Massachusetts, USA
- Wahid, A., 2004. Analysis of toxic and osmotic effects of sodium chloride on leaf growth and economic yield of sugarcane. *Bot. Bull. Acad. Sin.*, 45: 133–141
- Wahid, A., 2007. Physiological implications of metabolites biosynthesis in net assimilation and heat stress tolerance of sugarcane (*Saccharum officinarum*) sprouts. *J. Plant Res.*, 120: 219–228
- Wahid, A. and A. Shabbir, 2005. Induction of heat stress tolerance in barley seedlings by pre-sowing seed treatment with glycinebetaine. *Plant Growth Regul.*, 46: 133–141
- Wahid, A., S. Gelani, M. Ashraf and M.R. Foolad, 2007. Heat tolerance in plants: An overview. *Environ. Exp. Bot.*, 61: 199–223
- Yang, J., R.G. Sears, B.S. Gill and G.M. Paulsen, 2002. Quantitative and molecular characterization of heat tolerance in hexaploid wheat. *Euphytica*, 126: 275–282

Zhu, J.K., 2002 Salt and drought stress signal transduction in plants. *Annu. Rev. Plant. Biol.*, 53:  
247–273



**Fig. 1:** Maximum temperature inside and outside of the canopy (recorded at 12 noon) during crop growth months of 2004-2005 and 2005-2006

**Table 1:** Salinity tolerance threshold ( $EC_e$ ) of wheat varieties, change in it after foliar spray with thiourea and net improvement in  $EC_e$  at various growth stages

Growth stage	Varieties	Salinity	Salinity and thiourea spray	Net improvement (%)
Seedling	S-24	8.55±0.29a	9.27±0.82a	7.42±1.42b
	Inqlab-91	7.10±1.03b	7.74±0.84b	8.45±0.91ab
	MH-97	6.85±0.65b	7.52±0.51b	9.09±1.31a
	BK-2002	7.76±0.99ab	8.60±1.11ab	9.60±1.67a
	SH-2002	7.73±0.53ab	8.55±0.12ab	9.67±1.61a
Pre-anthesis	S-24	9.66±0.60a	10.43±0.70a	7.32±1.15c
	Inqlab-91	7.99±0.70b	8.82±0.83b	9.38±0.95b
	MH-97	6.63±1.21c	7.41±1.08c	10.88±1.13ab
	BK-2002	8.18±0.54b	9.21±0.68b	11.16±1.18a
	SH-2002	8.38±0.59b	8.97±0.55b	7.61±1.14c
Maturity	S-24	9.28±0.70a	10.11±0.65a	8.32±0.74b
	Inqlab-91	8.90±0.76a	9.46±0.67ab	5.89±0.72c
	MH-97	7.84±0.83b	8.70±0.93b	9.87±0.94a
	BK-2002	8.92±0.52a	9.71±0.31a	8.20±0.72b
	SH-2002	9.18±0.32a	10.33±0.68a	10.82±0.99a

Mean ± standard deviation. Values sharing same letters differ non-significantly ( $P>0.05$ )

**Table 2:** Changes in high temperature tolerance of wheat varieties, changes and net improvement in it after foliar spray with thiourea at various growth stages

Growth stage	Varieties	Salinity	Salinity and thiourea spray	Net improvement (%)
Seedling	S-24	14.95±1.98a	9.03±1.49a	5.93±0.49a
	Inqlab-91	10.06±0.84b	5.72±1.27b	4.33±0.68b
	MH-97	11.01±1.31b	5.79±0.57b	5.22±0.98ab
	BK-2002	14.61±1.67a	8.42±0.66a	6.19±0.79a
	SH-2002	11.94±1.61b	7.48±0.48a	4.47±0.97b
Pre-anthesis	S-24	21.78±1.95a	12.30±2.17a	9.48±1.39a
	Inqlab-91	15.20±1.09b	8.82±1.27c	6.38±0.91bc
	MH-97	11.61±1.78c	5.08±0.56c	6.53±1.04b
	BK-2002	16.27±1.96b	9.77±1.14bc	6.50±1.06b
	SH-2002	16.95±1.61b	11.83±1.18ab	5.12±0.81c
Maturity	S-24	22.86±2.28a	15.12±1.03a	7.74±0.26ab
	Inqlab-91	16.72±1.15b	8.32±0.32d	8.41±0.70a
	MH-97	10.47±0.68c	4.69±0.71e	5.78±0.93d
	BK-2002	17.31±2.40b	11.22±1.31b	6.09±0.60c
	SH-2002	16.24±2.20b	9.63±1.03c	6.61±0.69b

Mean ± standard deviation. Values sharing same letters differ non-significantly (P>0.05)

## Standard measurement units and their abbreviations

### Full name, Symbol

Barrel, bbl	Liter, L
Base, b	Megabyte, MB
Base pair, bp	Meter, m
Byte, B	Mega, M
Centimeter, cm	Metric ton, MT
Degree centigrade, °C	Microgram, µg
Degree Fahrenheit, °F	Microliter, µL
Einstein, <i>E</i>	Micrometer, µm
Giga, G	Milligram, mg
Gram, g	Milliliter, mL
Gravity, <i>g</i>	Millimeter, mm
Hectare, ha	Millimole, mM
Joule, J	Mole, <i>M</i>
Kilogram, kg	Revolutions per minute, rpm
Kilowatt, kW	Square, sq.
Kilowatt-hour, kWh	Volt, V
	Watt, W

### **Standard Countries abbreviations**

<b>Country full name</b>	<b>Abbreviation</b>
<i>Africa, African</i>	<i>Afr.</i>
<i>America, American</i>	<i>Amer.</i>
<i>Australia</i>	<i>Aust.</i>
<i>Austria</i>	<i>Aus.</i>
<i>Bangladesh</i>	<i>Bang.</i>
<i>Belgium</i>	<i>Belg.</i>
<i>Brazil</i>	<i>Braz.</i>
<i>British, Britain</i>	<i>Brit.</i>
<i>Bulgaria</i>	<i>Bulg.</i>
<i>Canadian</i>	<i>Can.</i>
<i>Chili, Chilean</i>	<i>Chil.</i>
<i>China, Chinese</i>	<i>Chin.</i>
<i>Egyptian</i>	<i>Egypt.</i>
<i>Emirates</i>	<i>Emir.</i>
<i>European</i>	<i>Eur.</i>
<i>India, India</i>	<i>Ind.</i>
<i>Indonesia, Indonesian</i>	<i>Indo.</i>
<i>Iranian</i>	<i>Iran.</i>
<i>Israel</i>	<i>Isr.</i>
<i>Italian</i>	<i>Ital.</i>
<i>Japan</i>	<i>Jpn.</i>
<i>Korea, Korean</i>	<i>Kor.</i>
<i>Malaysian</i>	<i>Malays.</i>
<i>New Zealand</i>	<i>N.Z.</i>
<i>Nigeria, Nigerian</i>	<i>Nig.</i>
<i>Pakistan</i>	<i>Pak.</i>
<i>Philippines</i>	<i>Phil.</i>
<i>South African</i>	<i>S. Afr.</i>
<i>Scandinavian</i>	<i>Scand.</i>
<i>Spanish</i>	<i>Span.</i>
<i>Sweden</i>	<i>Swed.</i>
<i>Turkey, Turkish</i>	<i>Turk.</i>

## Some common abbreviations in journals names

### Full name Abbreviation

<i>Academic, Academia; Acad.</i>	<i>Geology; Geol.</i>
<i>Addiction; Addict.</i>	<i>Hazardous; Haz.</i>
<i>Addition; Addit.</i>	<i>Horticulture, Horticultural, Hirticulturae; Hortic.</i>
<i>Advances; Advancement Adv.</i>	<i>Immunology, Immunological; Immun.</i>
<i>Agriculture, Agricultural; Agric.</i>	<i>Information, Informational; Inform.</i>
<i>Agronomy, Agronomic; Agron.</i>	<i>Instrument(s); Instrum.</i>
<i>Alternative, Alternate; Antern.</i>	<i>Integrative; Interg.</i>
<i>Analytical, Analysis; Anal.</i>	<i>Interaction(s), Interactive; Interact.</i>
<i>Anatomy; Anat.</i>	<i>International; Int.</i>
<i>Annal(s); Ann.</i>	<i>Journal(s); J.</i>
<i>Annual; Annu.</i>	<i>Letter(s); Lett.</i>
<i>Aquatculture; Aqualt.</i>	<i>Liquid; Liq.</i>
<i>Aquatic; Aquat.</i>	<i>Magnetic, Magnet; Magn.</i>
<i>Archive(s); Arch.</i>	<i>Management; Manage.</i>
<i>Association(s); Assoc.</i>	<i>Marine, Marina; Mar.</i>
<i>Bacteriological; Bacteriol.</i>	<i>Material(s); Mater.</i>
<i>Bacterium, Bacteria; Bacter.</i>	<i>Mathematics, Mathematical; Math.</i>
<i>Behavior, Behave; Behav.</i>	<i>Mechanic, Mechanical, Mechanism; Mech.</i>
<i>Biology, Biological, Biologia; Biol.</i>	<i>Medical, Medicine, Medica; Med.</i>
<i>Bioremediation; Bioremed.</i>	<i>Metabolism, Metabolic; Metab.</i>
<i>Biotechnology; Biotechnol.</i>	<i>Microbiology; Microbiol.</i>
<i>Botany, Botanical, Botanica; Bot.</i>	<i>Modern; Mod.</i>
<i>Breeding, Breeder; Breed.</i>	<i>Molecule, Molecular; Mol.</i>
<i>Cellular; Cell.</i>	<i>Monitor, Monitoring; Monit.</i>
<i>Chemistry, Chemical; Chem.</i>	<i>Morphology, Morphological; Morph.</i>
<i>Chromatograph(y), Chromatographer; Chromatogr.</i>	<i>Nature, Natural; Nat.</i>
<i>Clinical Clin.</i>	<i>Nutrition, Nutritional, Nutrient(s); Nutr.</i>
<i>Communication(s); Commun.</i>	<i>Oecologia; Oecol.</i>
<i>Computer(s), Computational Comput.</i>	<i>Operation, Operational; Oper.</i>
<i>Contamination, Contaminant; Contam.</i>	<i>Opinion; Opin.</i>
<i>Critical; Crit.</i>	<i>Optic, Optical; Opt.</i>
<i>Culture; Cult.</i>	<i>Organ , Organic; Org.</i>
<i>Current; Curr.</i>	<i>Parasitology, Parasitological; Parasotol.</i>
<i>Cycle, Cycling; Cycl.</i>	<i>Pathology, Pathological; Pathol.</i>
<i>Development, Developmental; Dev.</i>	<i>Pediatrics; Pediatr.</i>
<i>Diagnostic; Diagn.</i>	<i>Pharmacology; Pharmacol.</i>
<i>Disease; Dis.</i>	<i>Pharmacy, Pharmaceutical; Pharm.</i>
<i>Diverse, Diversity; Divers.</i>	<i>Philosophy, Philosophical; Phil.</i>
<i>Dynamic; Dynam.</i>	<i>Phycology, Phycological; Phycol.</i>
<i>Ecology , Ecological; Ecol.</i>	<i>Physics, Physical; Phys.</i>
<i>Economic, Economical; Econ.</i>	<i>Physiology, Physiological, Physiologia; Physiol.</i>
<i>Electric, Electrical; Electr.</i>	<i>Phytology, Phytologist(s); Phytol.</i>
<i>Electronic; Electron.</i>	<i>Plantarum; Plant.</i>
<i>Engineering, Engineer; Eng.</i>	<i>Pollution; Pollut.</i>
<i>Entomology, Entomological; Entomol.</i>	<i>Polymer; Polym.</i>
<i>Environment, Environmental; Environ.</i>	<i>Practice, Practical, Practitioner; Prac.</i>
<i>Epidemiology; Epidemiol.</i>	<i>Production, productive; Prod.</i>
<i>Experimental; Exp.</i>	<i>Program, Progress, Progressive; Prog.</i>
<i>Faculty; Fac.</i>	<i>Quality, Qualitative; Qual.</i>
<i>Forestry, Forest; For.</i>	<i>Quantity, Quantitative; Quant.</i>
<i>Function(s), Functional; Funct.</i>	<i>Radiation; Radiat.</i>
<i>General; Gen.</i>	<i>Registration; Reg.</i>
<i>Genetics, Genetica; Genet.</i>	<i>Regulation, Regulator(s); Regul.</i>

*Rehabilitation; Rehabil.*  
*Reproduction, Reproductive; Reprod.*  
*Resource(s); Resour.*  
*Revegetation; Reveg.*  
*Review(s); Rev.*  
*Safe, Safety; Saf.*  
*Science, Scientia, Scientific; Sci.*  
*Sensing, Sensor(s); Sens.*  
*Service(s); Serv.*  
*Signaling; Signal.*  
*Sinica; Sin.*  
*Society; Soc.*  
*Software; Softw.*  
*Statistical, Statistics, Statistician; Stat.*  
*Structure, Structural; Struct.*  
*Studies; Stud.*  
*Surface; Surf.*  
*Surgery, Surgical; Surg.*  
*Survey(s); Surv.*

*Sustainable; Sustain.*  
*System(s), Systematic(s); Syst.*  
*Technique, Technical; Tech.*  
*Technological; Technol.*  
*Theoretical, Theory, Theorem; Theor.*  
*Therapy, Therapeutical; Ther.*  
*Tillage; Till.*  
*Toxic; Tox.*  
*Toxicology, Toxicological; Toxicol.*  
*Transaction(s); Trans.*  
*Tropical, Tropics; Trop.*  
*Vegatable(s), Vegatale; Veg.*  
*Veterinary, Veterinus; Vet.*  
*Virological; Virol.*  
*Virus, Viral; Vir.*  
*Weekly; Wkly.*  
*Wetland; Wetl.*  
*Zoology, Zoological, Zoologica; Zool.*

**Journal names not to be abbreviated when given as SINGLE WORD**

<i>Addiction</i>	<i>Biorheology</i>	<i>Genetica</i>
<i>Aerobiologia</i>	<i>Bioscience</i>	<i>Genetics</i>
<i>Age</i>	<i>Biostatistics</i>	<i>Genome</i>
<i>Agribusiness</i>	<i>Biosystems</i>	<i>Genomics</i>
<i>Agrochimica</i>	<i>Biotechniques</i>	<i>Geobiology</i>
<i>Aids</i>	<i>Biotropica</i>	<i>Glycobiology</i>
<i>Alcohol</i>	<i>Blood</i>	<i>Grana</i>
<i>Allergologie</i>	<i>Bone</i>	<i>Heart</i>
<i>Allergy</i>	<i>Brain</i>	<i>Helicobacter</i>
<i>Ambio</i>	<i>Breast</i>	<i>Helminthologia</i>
<i>Animal</i>	<i>Bryologist</i>	<i>Hematology</i>
<i>Anthropologist</i>	<i>Burns</i>	<i>Hemoglobin</i>
<i>Antibiotiques</i>	<i>Caryologia</i>	<i>Hereditas</i>
<i>Apoptosis</i>	<i>Cell</i>	<i>Heredity</i>
<i>Appetite</i>	<i>Chemosphere</i>	<i>Hortscience</i>
<i>Aquaculture</i>	<i>Chest</i>	<i>Hydrobiologia</i>
<i>Atherosclerosis</i>	<i>Chimia</i>	<i>Hypertension</i>
<i>Atmosfera</i>	<i>Chromatographia</i>	<i>Iforest</i>
<i>Autophagy</i>	<i>Chromosoma</i>	<i>Immunity</i>
<i>Bioanalysis</i>	<i>Circulation</i>	<i>Immunobiology</i>
<i>Biochemistry</i>	<i>Clinics</i>	<i>Immunogenetics</i>
<i>Biochimie</i>	<i>Computer</i>	<i>Immunology</i>
<i>Biocontrol</i>	<i>Computing</i>	<i>Infection</i>
<i>Biodegradatiom</i>	<i>Cornea</i>	<i>Injury</i>
<i>Bioelectrochemistry</i>	<i>Cortex</i>	<i>Interciencia</i>
<i>Bioelectromagnetics</i>	<i>Cryobiology</i>	<i>Intervirology</i>
<i>Bioessays</i>	<i>Dendrobiology</i>	<i>Kardiologiya</i>
<i>Bioethics</i>	<i>Dendrochronologia</i>	<i>Labmedicine</i>
<i>Biofabrication</i>	<i>Dermatitis</i>	<i>Maydica</i>
<i>Biofactors</i>	<i>Development</i>	<i>Mealthmed</i>
<i>Biofouling</i>	<i>Diabetes</i>	<i>Medicine</i>
<i>Biofutur</i>	<i>Diabetologe</i>	<i>Metabolism</i>
<i>Biogeochemistry</i>	<i>Diabetologia</i>	<i>Metabolomics</i>
<i>Biogeosciences</i>	<i>Digestion</i>	<i>Molecules</i>
<i>Biogerontology</i>	<i>Drugs</i>	<i>Mutagenesis</i>
<i>Bioinformatics</i>	<i>Ecotoxicology</i>	<i>Mycologia</i>
<i>Biointerphases</i>	<i>Ecotropica</i>	<i>Mycopathologia</i>
<i>Biologia</i>	<i>Environment</i>	<i>Mycorrhiza</i>
<i>Biologicals</i>	<i>Environmetrics</i>	<i>Mycoscience</i>
<i>Biomacromolecules</i>	<i>Epidemiology</i>	<i>Mycoses</i>
<i>Biomarkers</i>	<i>Epigenetics-us</i>	<i>Mycotaxon</i>
<i>Biomaterials</i>	<i>Epigenomics-uk</i>	<i>Nature</i>
<i>Biomedica</i>	<i>Epilepsia</i>	<i>Nephrology</i>
<i>Biometals</i>	<i>Euphytica</i>	<i>Neuroinformatics</i>
<i>Biometrics</i>	<i>Farmacia</i>	<i>Neurologist</i>
<i>Biometrika</i>	<i>Fisheries</i>	<i>Neurology</i>
<i>Biomicrofluidics</i>	<i>Flora</i>	<i>Neuromodulation</i>
<i>Biopolymers</i>	<i>Forestry</i>	<i>Neuropathology</i>
<i>Bioresources</i>	<i>Fruits</i>	<i>Neuropediatrics</i>

*Neuropeptides*  
*Neuropharmacology*  
*Nutrients*  
*nutrition*  
*Oceanologia*  
*Oecologia*  
*Oncogene*  
*Palaeontology*  
*Palynology*  
*Pancreas*  
*Pancreatology*  
*Pathobiology*  
*Pathologie*  
*Pathology*  
*Patient*  
*Pediatrics*  
*Pedologia*  
*Pedosphere*  
*Pharmacogenomics*  
*Pharmacology*  
*Photosynthetica*  
*Phycologia*

*Physiology*  
*Physiotherapy*  
*Phytochemistry*  
*Phytomedicine*  
*Phytoparasitica*  
*Phytopathology*  
*Planta*  
*Platelets*  
*Primates*  
*Prostate*  
*Proteins*  
*Proteomics*  
*Protist*  
*Protoplasma*  
*Protoplasma*  
*Rehabilitation*  
*Reproduction*  
*Respiration*  
*Respirology*  
*Science*  
*Scienceasia*  
*Scientist*

*Statistics*  
*Steroids*  
*Stress*  
*Symbiosis*  
*Theriogenology*  
*Transfusion*  
*Tuberculosis*  
*Ultramicroscopy*  
*Urologie*  
*Urology*  
*Vaccine*  
*Virologie*  
*Virology*  
*Wetlands*  
*Yeast*  
*Zoology*  
*Zoomorphology*  
*Zoosystema*  
*Zootaxa*  
*Zuchtungskunde*  
*Zuckerindustrie*  
*Zygote*