



**Full Length Article**

## Phenotypic Characterization of *Phytophthora infestans* Population in Southwestern China

Honghao Li<sup>1,2</sup>, Jing Shang<sup>1</sup>, Hong Zhang<sup>2</sup>, Bowei Liu<sup>2</sup>, Yunliang Peng<sup>2</sup> and Yun Huang<sup>1\*</sup>

<sup>1</sup>College of Agriculture, Sichuan Agricultural University, Wenjiang 611130, China

<sup>2</sup>Institute of Plant Protection, Sichuan Academy of Agricultural Sciences, Key Laboratory of Integrated Pest Management on Crops in Southwest, Ministry of Agriculture, Chengdu 610066, China

\*For correspondence: 5787huangyun@sina.com.cn

Received 24 August 2019; Accepted 18 January 2020; Published 02 April 2020

### Abstract

Potato late blight, caused by the aggressive pathogen *Phytophthora infestans* (Mont.) de Bary, is one of the most destructive potato diseases worldwide. A total of 688 isolates of *P. infestans* collected from five provinces in southwestern China during 2012 to 2018, were assessed with regard to mating type, response to metalaxyl, virulence race and mitochondrial (mt)DNA haplotype. In terms of mating type, 76% of 679 isolates were of the self-fertile mating type, 16.6% of the isolates were of the A1 mating type and only 7.8% were of the A2 mating type. Among the 299 isolates tested as representative samples, the frequency of resistant to metalaxyl reached 80% in these regions. The race structure of the strains presented variety and complexity and 9.04 virulence factors per isolate and 92 virulence races were found among the 365 tested isolates. The most common pathotypes were 1.2.3.4.5.6.7.8.9.10.11 and 1.2.3.4.5.6.7.8.10.11, representing 46% of the tested strains. Three types of mtDNA haplotypes were found: 78.7% of 362 isolates were Ia, 16.6% were Iib and 4.7% were Iia. The results indicated that the population of *P. infestans* in southwestern China is diverse; and the appearance of the self-fertile isolates with resistance to metalaxyl may contribute to the potential for sexual recombination and disease epidemics. © 2020 Friends Science Publishers

**Keywords:** Mating type; Metalaxyl sensitivity; mtDNA; *Phytophthora infestans*; Virulence phenotype

### Introduction

*Phytophthora infestans* caused the famous Great Famine in Ireland in the 19<sup>th</sup> century, resulting in potato tuber yield loss and approximately one million people starved to death (Fry and Goodwin 1997a). *P. infestans* is known as a heterothallic oomycete. It is widely believed that the first migration of *P. infestans* from the highlands of central Mexico likely occurred in the 1840s (Fry and Goodwin 1997b). Migration plays an important role in the population structure of *P. infestans*. Previous studies have suggested that the population structure of *P. infestans* changed in the past (Hohl and Iselin 1984; Fry *et al.* 1993; Drenth *et al.* 1994; Wharton *et al.* 2015). Oospores may be expected to be reproduced in the populations of *P. infestans* consisting of the A1 and A2 mating types. The evidence that *P. infestans* reproduces sexually on a regular basis is increasing in northern Europe (Yuen and Andersson 2013). Sexual reproduction may occur, which would contribute to increase the genotypic diversity in the *P. infestans* population (Drenth *et al.* 1994; Pipe *et al.* 2000; Śliwka *et al.* 2006). The population characteristics of *P. infestans* are complex

and variable in many countries and many different virulence races have been identified (Kiiker *et al.* 2018; Fukue *et al.* 2018). Potato late blight is considered to be a devastating disease for potato growers and causes more than \$6 billion in losses and management costs every year (Haverkort *et al.* 2008). Due to the widespread use of fungicides, metalaxyl-resistant strains have been widely reported (Aav *et al.* 2015).

Mitochondrial (mt)DNA is uniparentally inherited (Whittaker *et al.* 1994), ideal for tracing lines of descent and easily detected. In addition, mtDNA polymorphisms of *P. infestans* are used to monitor pathogen populations (Griffith and Shaw 1998). Haplotype Ib was identified as the 'old' population of *P. infestans* only in the highlands of central Mexico, and haplotypes Ia, Iib and Iia were classified as the 'new' population (Runno-Paurson *et al.* 2009). Previous studies of phenotypic diversity in China showed that haplotypes Ia, Iia and Iib exist in China. Haplotype Ia is the most common genotype in Fujian (Han *et al.* 2014), Sichuan (Li *et al.* 2013a) and Yunnan (Zhao *et al.* 2002) in China, whereas haplotype Iia is common in Qinghai (Lian *et al.* 2012).

Although China has become the largest potato producer in the world (Alva et al. 2011), the further development of the potato industry has been seriously threatened by potato late blight, especially in southwestern China. Potato production has decreased by 30% every year due to this disease. Unfortunately, very little is known about the population characteristics of this pathogen in southwestern China. To better understand the characteristics of this pathogen in southwestern China, phenotypic and mtDNA haplotypes were analysed in this study, and the sampling area covered the region with the largest potato planting area in China. The results will provide a guideline for future potato breeding and effective prevention and control measures for potato late blight.

## Materials and Methods

### Collection and isolations

Potato leaves with single lesions showing classical typical late blight symptoms were obtained from 33 locations in southwestern China from April to October. The distance between each sampling point is more than 50 kilometres. The elevations ranged from 189 m to 3430 m. In those areas, farmers plant high-quality virus-free seed potato and apply fungicide three to five times to control potato late blight. The infected leaves were cut into small pieces (5 mm<sup>2</sup>) and placed under potato tuber slices from the susceptible cv. Favorita. Small tufts of mycelia grew on the potato slices after being incubated for five to seven days at 18°C in darkness, and the hyphae were transferred to culture medium plates with rye B agar and incubated at 18°C (Li et al. 2013a).

During 2012–18, 688 isolates of *P. infestans* were collected (Table 1). Some isolates from each site were chosen for the analysis of the population diversity of *P. infestans* in terms of mating type, sensitivity to metalaxyl, virulence race and mtDNA haplotype.

### Mating type detection

Isolates of *P. infestans* were evaluated to identify their mating type on rye B agar by pairing them with the *P. infestans* reference isolates A1 (VK98014) and A2 (90128). The confrontational strains were stored at 18°C in the dark for 5 to 7 days and were observed microscopically for oospores in the hyphae interaction area. Mating types were identified according to the method described by Runno-Paurson et al. (2009). The isolates generating oospores with the A1 were classified as A2; the isolates that generating oospores with A2 were classified as A1. The isolates forming oospores with or without A1 and A2 were designated as self-fertile.

### Metalaxyl resistance assessment

The responses of different isolates to metalaxyl were tested on rye B agar plates (90 mm) with the corresponding

**Table 1:** Origin of *P. infestans* isolates collected from southwestern China (2012–2018)

Region	Location	Number of isolates	Latitude	Longitude	Altitude (m)
Guizhou	Weining	28	26°83'	104°24'	2169
	Chishui	8	28°47'	105°76'	299
Hubei	Lichuan	10	30°23'	108°99'	1129
Chongqing	Kaizhou	23	31°20'	108°34'	529
	Shizhu	4	29°99'	108°11'	553
	Wuxi	20	31°41'	109°56'	256
	Yunyang	3	31°36'	108°91'	1280
	Zhongxian	2	30°29'	108°03'	189
Sichuan	Beichuan	6	31°94'	104°41'	996
	Chaotian	11	32°62'	106°10'	1412
	Chongzhou	37	30°54'	103°65'	508
	Danling	6	29°98'	103°36'	485
	Daofu	31	30°48'	101°48'	3430
	Ebian	10	29°24'	103°18'	1197
	Jiangyou	9	31°97'	104°78'	588
	Leibo	7	28°39'	103°77'	1118
	Luding	59	29°64'	102°12'	1562
	Mianning	9	28°74'	102°25'	2076
	Pengzhou	193	31°21'	103°78'	1050
	Puge	1	27°48'	102°48'	1417
	Shunqing	24	31°05'	106°13'	324
	Tongjiang	10	32°47'	107°37'	1200
	Wanyuan	10	32°11'	108°10'	1044
	Xiaojin	43	31°08'	102°30'	3162
	Xide	16	28°30'	102°45'	2410
	Xindu	8	30°77'	104°21'	478
	Xingwen	8	28°27'	105°29'	399
	Xuanhan	27	31°35'	107°72'	297
Yilong	10	30°87'	106°08'	313	
Zhaojue	26	28°01'	102°51'	2600	
Yunnan	Daguan	6	27°74'	103°89'	1135
	Huize	11	26°43'	103°32'	2129
	Ludian	12	27°17'	103°58'	1908
Total		688			

metalaxyl concentration. The fungicide was dissolved in 0.1% acetone and prepared in a stock of 100 gL<sup>-1</sup>. Three different metalaxyl concentrations (0, 5 and 100 mg L<sup>-1</sup>) were tested. Uniformly-sized agar plugs (5 mm) were taken from actively growing colonies of *P. infestans*, and a plug was placed in the middle of each plate. After maintaining the cultures for 7 days at 18°C in the dark, the diameters of the fungal colonies were evaluated in two perpendicular directions through the centre of each plate. Three replicates were used for each isolate. Metalaxyl resistance was determined according to the following scale: those with growth on both the 5 and 100 mg L<sup>-1</sup> plates ≥40% of that on the 0 mg L<sup>-1</sup> plate were regarded as resistant (R), those with growth on the 5 mg L<sup>-1</sup> plate ≥40% of that on the 0 mg L<sup>-1</sup> plate were regarded as intermediate (I), and those with growth on both the 5 and 100 mg L<sup>-1</sup> plates ≤40% of that on the 0 mg L<sup>-1</sup> plate were regarded as sensitive (S) (Forbes 1997).

### Virulence tests

The virulence pathotype was assessed by testing interactions on a subset of different potato genotypes with resistance genes *R1-R11* (Malcolmson and Black 1966). The sterile

differential potato clones were preserved on Murashige and Skoog's medium and were maintained at 23°C with a 16 h light period. Four fully expanded leaves were selected from the potato plants for each isolate and placed abaxial-side up on a wet filter paper in a plastic Petri dish. A 10  $\mu\text{L}$  drop of sporangial suspension ( $2 \times 10^4$  sporangia  $\text{mL}^{-1}$ ), prepared after 5–7 days on tuber slices from susceptible cv. Favorita was placed on each leaflet. After inoculation, the Petri dishes were maintained at 18°C under a 16 h light period. The leaves were checked with a stereomicroscope for sporulation 7 days later. If sporulation was detected, the interaction was evaluated as compatible; if a hypersensitive or no symptom response was detected, the interaction was evaluated as incompatible (Runno-Paurson *et al.* 2009). The results were credible when sporulation occurred on the leaves of impressionable potato clone *r* without any R genes.

### DNA extraction

Isolates of *P. infestans* were incubated at 18°C in the dark for 7 days individually on potato tuber slices of susceptible cv. Favorita, and mycelia were carefully harvested by removing them with an inoculating needle and stored at -20°C in centrifuge tubes for DNA extraction. Genomic DNA from 30 mg of dry mycelium was purified using an E.Z.N.A.® Fungal DNA Kit (OMEGA Bio-Tek, USA). The quantity of the purified DNA samples was measured using a NanoDrop ND-1000 (NanoDrop Technologies, Inc., USA). The DNA samples were diluted to 20 ng  $\mu\text{L}^{-1}$  and stored at -20°C for further analysis.

### mtDNA haplotypes

The mtDNA haplotypes of the *P. infestans* isolates were described by Griffith and Shaw (1998). The mtDNA regions of *P. infestans* were amplified using primer pair 1 (F2 - TTCCCTTTGTCCTCTACCGAT; R2 - TTACGGCGGTTTAGCACATACA) and primer pair 2 (F4 - TGGTCATCCAGAGGTTTATGTT; R4 - CCGATACCGATACCAGCACCAA). Each PCR (25  $\mu\text{L}$ ) mixture consisted of 12.5  $\mu\text{L}$  of 2 $\times$ Taq PCR Master Mix (Sangon Biotech Co., Ltd, China), 10  $\mu\text{M}$  of each primer, 30–50 ng of DNA sample and 8.5  $\mu\text{L}$  of ddH<sub>2</sub>O. The thermal cycling procedure was as follows: initial denaturation for 2 min at 94°C, 35 cycles of denaturation at 94°C for 1 min, annealing at 62°C for 30s and extension at 72°C for 1 min, and a final extension at 72°C for 10 min.

The PCR product P2 was digested with the restriction enzymes *MspI* and the product P4 was digested with the restriction enzymes *EcoRI* at 37°C for 4 h. Digestion mix P2 (20  $\mu\text{L}$ ) consisted of 1  $\mu\text{L}$  of *MspI* (10 U  $\mu\text{L}^{-1}$ ), 2  $\mu\text{L}$  of buffer, 9  $\mu\text{L}$  of ddH<sub>2</sub>O and 8  $\mu\text{L}$  of the PCR product P2. Digestion mix P4 (20  $\mu\text{L}$ ) consisted of 0.5  $\mu\text{L}$  of *EcoRI* (20 U  $\mu\text{L}^{-1}$ ), 2  $\mu\text{L}$  of buffer, 9.5  $\mu\text{L}$  of ddH<sub>2</sub>O and 8  $\mu\text{L}$  of the PCR product P4. The types of mtDNA haplotypes were identified according to the sizes of the digested PCR products (Griffith and Shaw 1998).

### Data analysis

The normalised Shannon index (Sheldon 1969; Runno-Paurson 2009) was used to summarize the pathotype diversity and the formula was as follows:  $H_s = -\sum g_i \ln g_i / \ln N$ , where  $g_i$  is the frequency of races  $i$  and  $N$  is the number of isolates. The index was normalized to a scale of 0 (no diversity) to 1 (a unique pathotype per isolate). SPSS version 22.0 (SPSS Inc., Chicago, IL, USA) was used for statistical analyses. To identify the differences in terms of specific virulence among the different potato R genes or sampling years, ANOVA was performed.

## Results


### Mating type

In total, 679 isolates were randomly chosen for each mating type. The results showed that 16.6% of the isolates were of the A1 mating type, 7.8% of the isolates were of the A2 mating type and 75.6% of the isolates were self-fertile (Fig. 1A). Many self-fertile isolates were detected in the five regions.


A1 mating type isolates were detected every year, and their frequency ranged from 26.7 to 4.2% throughout the survey. However, the A2 mating type was not found in 2015 and 2017, with frequencies ranging from 0 to 33.3%. Conversely, the self-fertile mating type prevailed among the isolates, and the frequency during the sampling years was in the range of 60.0–87.7% (Fig. 1B). Interestingly, most isolates collected in the fall in Sichuan in 2016–2018 were of the A1 mating types. The A1, A2 and self-fertile mating types were sometimes found in the same field in Sichuan. The isolates collected from Hubei and Chongqing were all of the self-fertile mating type. Furthermore, the self-fertile mating type frequencies in Guizhou and Yunnan provinces were 97.2 and 93.1%, respectively (Fig. 1C).

### Metalaxyl resistance

The metalaxyl resistance of 299 isolates was determined on rye B agar plates. Of these isolates, 79.6% were classified as resistant, 17.4% had intermediate sensitivity, and 3.0% were sensitive to metalaxyl (Fig. 2A). Metalaxyl-resistant isolates were detected among the three mating types (A1, A2 and self-fertile mating types), and no sensitive isolates were found in the A2 mating type (Fig 2B). The proportion of metalaxyl-resistant isolates reached from 55.0 to 91.4% in southwestern China during the period of 2012–18. The metalaxyl-sensitive phenotype was not found in 2013, 2014, 2015 or 2017; only 10.0% ( $n = 2$ ), 4.5% ( $n = 3$ ) and 16.0% ( $n = 4$ ) of the isolates were sensitive in 2012, 2016 and 2018, respectively (Fig. 2C). All isolates collected from Guizhou, Hubei and Yunnan were resistant to metalaxyl. Additionally, 76.6% ( $n = 183$ ) of the isolates collected from Sichuan and 80.0% ( $n = 20$ ) of the isolates collected from Chongqing showed resistance to metalaxyl (Fig. 2D).




**Fig. 1:** Frequency of mating types of *P. infestans* isolates from different regions in southwestern China during the 2012–2018 period. (A) all isolates; (B) different sampling years; (C) different sampling regions




**Fig. 2:** Metalaxyl resistance of *P. infestans* isolates from different regions in southwestern China during the 2012–2018 period. (A) For all isolates; (B) with respect to mating type; (C) for different sampling years; (D) for different sampling regions. S: metalaxyl sensitive, I: intermediate metalaxyl sensitivity, R: metalaxyl resistant

### Virulence phenotype

A total of 365 isolates as representative samples purified from the five provinces were tested for pathogenicity. The results showed that virulence factors that could overcome all known resistance genes (R1-R11) were detected. The most common physiological races were 1.2.3.4.5.6.7.8.9.10.11 (26.58%) and 1.2.3.4.5.6.7.8.10.11 (18.90%). The frequency of virulence genes was in the range of 67–89%; the lowest frequency was found for virulence against R9 (67% ± 4.6 SE) and the highest frequency was found for virulence against R6 (89% ± 4.4 SE) (Fig. 3). There were no differences in specific virulence ( $P \geq 0.05$ ), but there were significant differences among the seven years ( $P \leq 0.01$ ). Ninety-nine percent of the tested strains were able to overcome five or more R genes. Moreover, 92 virulence



**Fig. 3:** Frequency of virulence to resistance genes in *P. infestans* isolates from different regions in southwestern China during the 2012–2018 period



**Fig. 4:** Composition and frequency of mitochondrial haplotypes of isolates from different regions in southwestern China during the 2012–18 period. (A) For all isolates (n=362); (B) for different sampling regions; (C) with respect to mating type

genes were detected. The mean number of virulence factors was 9.04 (Table 2) and ranged from 7.2 to 10.4 among the sampling years (Table 3). The normalized Shannon diversity index of the *P. infestans* population in southwest China was 0.55.

### mtDNA haplotypes

Three types of mtDNA haplotypes (Ia, IIa and IIb) were identified in 362 tested isolates. In total, 78.7% (n = 285) of the isolates were Ia. The frequencies of IIa and IIb were 4.7% (n = 17) and 16.6% (n = 60), respectively (Fig. 4A). Haplotype Ia was the common type in all regions, with a frequency of 100% among the tested isolates from Guizhou and Yunnan and 90.0, 72.0 and 95.0% in Hubei, Sichuan and Chongqing, respectively. Haplotype IIa was detected in Hubei and Sichuan. Haplotype IIb was found only in Sichuan and Chongqing (Fig. 4B). Among the five regions, three mtDNA haplotypes were detected in Sichuan. Each

**Table 2:** Number and frequency of isolates of different races among *P. infestans* isolates from southwestern China (2012–2018)

Race	Isolates (n)	Frequency (%)	Race	Isolates (n)	Frequency (%)
1.2.3.4.5.6.7.8.9.10.11	97	26.58	1.2.3.4.6.7.9.10.11	1	0.27
1.2.3.4.5.6.7.8.10.11	69	18.90	1.2.3.4.6.9.10.11	1	0.27
1.3.4.6.7.8.10	16	4.38	1.2.3.4.7.8.10	1	0.27
1.3.4.5.6.7.8.9.11	8	2.19	1.2.3.5.6.8	1	0.27
1.3.4.5.8.9.10	8	2.19	1.2.3.5.6.8.10	1	0.27
2.3.5.6.7.8.9.11	8	2.19	1.2.3.6	1	0.27
1.2.3.4.5.6.7.8.9.11	7	1.92	1.2.4.5.6.7.8.10	1	0.27
1.2.3.4.6.7.8.10	7	1.92	1.2.4.5.6.8.11	1	0.27
1.2.4.5.6.8.9.10	6	1.64	1.2.4.5.6.8.9	1	0.27
1.3.4.5.6.7.8.9.10.11	6	1.64	1.2.4.5.6.9.11	1	0.27
1.2.3.5.6.8.9.11	5	1.37	1.2.4.6.7.9.11	1	0.27
1.2.4.6.7.8.9.10.11	4	1.10	1.2.4.7.8.9.10.11	1	0.27
1.2.4.6.8.9.10.11	4	1.10	1.2.5.6.7.9.10.11	1	0.27
1.2.3.4.5.6.7.9.11	3	0.82	1.3.4.5.6.7.10.11	1	0.27
1.2.3.4.5.6.8.9.11	3	0.82	1.3.4.5.6.7.8.10.11	1	0.27
1.2.3.4.5.8.9.10.11	3	0.82	1.3.4.5.6.7.8.9.10	1	0.27
1.2.3.4.5.8.9.11	3	0.82	1.3.4.5.6.7.9	1	0.27
1.2.3.4.6.10.11	3	0.82	1.3.4.5.6.9.10.11	1	0.27
1.2.3.4.6.8.10.11	3	0.82	1.3.4.5.7.9.11	1	0.27
1.2.4.5.6.7.9.10.11	3	0.82	1.3.4.5.8.10	1	0.27
1.2.5.6.7.9	3	0.82	1.3.4.5.8.9.10.11	1	0.27
1.2.5.6.7.9.11	3	0.82	1.3.4.5.9.10	1	0.27
1.2.5.6.8.9.10.11	3	0.82	1.3.4.6.8.9.10.11	1	0.27
1.3.4.5.6.10	3	0.82	1.3.4.7.8.10.11	1	0.27
3.4.5.6.7.8.9.10.11	3	0.82	1.3.5.6.7.9.11	1	0.27
1.2.3.4.5.6.10.11	2	0.55	1.3.5.6.8.9.11	1	0.27
1.2.3.4.5.6.9.10	2	0.55	1.3.5.8.10	1	0.27
1.2.3.4.5.9.10	2	0.55	1.3.6.8.10	1	0.27
1.2.3.5.6.7.8.9.10.11	2	0.55	1.4.5.6.7.9.10.11	1	0.27
1.2.4.5.6.7.8.10.11	2	0.55	1.4.5.6.7.9.11	1	0.27
1.2.4.5.6.7.8.9.10	2	0.55	1.4.5.6.8.9.10.11	1	0.27
1.2.4.5.6.8.10.11	2	0.55	1.4.5.6.9.10.11	1	0.27
1.2.4.6.7.8.10.11	2	0.55	1.4.6.7.8.10.11	1	0.27
1.2.5.6.7.9.10	2	0.55	1.4.6.9.10.11	1	0.27
1.2.5.6.8.9.10	2	0.55	2.3.4.5.6.8.10	1	0.27
1.3.4.5.6.9.11	2	0.55	2.3.4.7.8.9.10.11	1	0.27
1.3.4.6.7.8.9.10.11	2	0.55	2.3.4.9.11	1	0.27
1.3.4.6.7.9.10	2	0.55	2.3.5.8.9.10	1	0.27
1.3.5.8.9.10	2	0.55	2.3.7	1	0.27
1.4.5.6.7.8.9.11	2	0.55	2.4.5.6.8	1	0.27
2.3.4.5.8.9.10	2	0.55	2.4.5.7.8	1	0.27
2.3.5.6.8.9.11	2	0.55	2.4.6.7.9.10.11	1	0.27
1.2.3.4.5.6.7.10.11	1	0.27	2.5.6.8.9.10	1	0.27
1.2.3.4.5.6.8.10.11	1	0.27	2.6.8.9.10	1	0.27
1.2.3.4.5.6.8.9.10.11	1	0.27	3.4.7.8	1	0.27
1.2.3.4.6.7.8.9.10.11	1	0.27	5.8.10.11	1	0.27
Total number of isolates	365				
Total number of races	92				
Virulence complexity	9.04				

mtDNA haplotype contained A1, A2 and self-fertile isolate clones. Most of haplotype Ia isolates were self-fertile and most of haplotype IIb isolates were A1 (Fig. 4C).

## Discussion

In this study, A1 and A2 mating types of *P. infestans* isolates were found in southwestern China, but most of the isolates were self-fertile. A2 isolates were only detected in Sichuan and Yunnan, and the isolates from Hubei and Chongqing were all self-fertile. The A2 mating type has been subsequently reported in many countries since the discovery of oospores produced by *P. infestans* under natural conditions in Mexico in 1956 (Niederhauser 1956). Since the first report of A2 mating type isolates in northern

**Table 3:** Frequency of virulence to potato R genes among isolates across seven sampling years in southwestern China

Resistance gene	Sampling years						
	2012	2013	2014	2015	2016	2017	2018
R1	87	62	87	81	100	100	100
R2	93	55	57	55	88	91	100
R3	87	86	68	81	82	92	93
R4	53	62	70	84	100	99	100
R5	87	72	70	58	89	92	93
R6	100	72	86	77	88	100	100
R7	67	52	59	68	73	93	93
R8	100	86	63	74	89	96	100
R9	93	69	62	55	66	63	64
R10	60	62	79	71	78	92	100
R11	93	45	48	61	87	96	100
Mean number of virulence isolates	9.2	7.2	7.5	7.6	9.4	10.2	10.4
Number of isolates tested	15	29	63	31	94	119	14

China in 1996 (Zhang et al. 1996), several A2 mating type isolates have been detected in many regions of China. Previous reports documented that the frequency of the A2 mating type was 10% in Yunnan, but it was 91% in Sichuan Province (Li et al. 2013b). The self-fertile isolate was first reported in China in 2002 (Huang 2002). Many self-fertile isolates have been found in China (Li et al. 2009; Li et al. 2013a; Zhu et al. 2015; Tian et al. 2016) and other countries (Aav et al. 2015; Casa-Coila et al. 2017). The probability of the occurrence of self-fertile isolates is related to the survival, environment and genetic structure of *P. infestans*, and there are many cultural practices in which the age of the culture, presence of other organisms, wounding and the addition of fungicide to the culture medium can override heterothallism-inducing oospore production in *P. infestans* (Smart et al. 2000). Self-fertile isolates that replaced A1 and A2 mating type isolates were the dominant clones in southwestern China, indicating that sexual reproduction can occur in all studied potato fields. Early epidemics of potato late blight can be caused by oospores in the field (Hannukkala et al. 2007; Brylińska et al. 2016). In contrast to sporangia, oospores can withstand adverse conditions or circumstances and survive into the next growing season in the soil without a host (Andersson et al. 1998; Fernández-Pavía et al. 2004; Lehtinen and Hannukkala 2004). Furthermore, sexual reproduction not only increases the genotypic variability in *P. infestans* populations, but may also cause increased pathogenicity and/or antimicrobial resistance (Fry et al. 1993; Harutyunyan et al. 2008; Lozoya-Saldaña 2011).

Fungicide-resistance testing indicated that metalaxyl-resistant isolates are widespread in southwestern China. The occurrence of metalaxyl-resistant isolates was not associated with the mating types. Many resistant isolates were detected in Yunnan (Zhao et al. 2007) in China. Resistance to metalaxyl was found in three regions, Guizhou, Hubei and Yunnan, with the highest proportion among the self-fertile isolates in our study. Metalaxyl-based fungicides are recommended for use in many regions because of their low price and high quality. It is possible that extensive use of fungicides caused the increase in the metalaxyl-resistant population. These studies suggest that it is necessary to change the strategy of using metalaxyl to control potato late blight. With the application of potato-resistant varieties and crop rotation (Bimšteine 2008), the amount of fungicides used can be reduced.

In our study, 92 different pathogenic types were identified among 365 isolates and the virulence complexity was 9.04. The most common race was 1.2.3.4.5.6.7.8.9.10.11, which was virulent to 11 known R genes. This means that the *P. infestans* population in southwestern China is highly diverse and complex. The proportion of virulence against R genes was also different in other countries (Chmielarz et al. 2014), and the sexual reproduction in the *P. infestans* populations has been detected in some countries (Küiker et al. 2018). Planting

resistant cultivars is an economical way to control potato late blight, but susceptible cultivars, such as cv. Favorita, are widely grown due to the short growth period, such as cv. Favorita. In some poor areas in southwestern China, cheap fungicides, such as metalaxyl-based fungicides, are applied for potato late blight control, or nothing is done about such infestations. As a result, potato late blight is common in southwestern China. There is a high level of pathotype diversity in southern China. This finding might be related to the specific conditions of year-round potato cultivation.

Overall, mtDNA haplotype tests revealed that all isolates in southwestern China were the 'new' population of *P. infestans*. The most common mtDNA haplotype was Ia, followed by haplotypes Iib and IIa; Ib is not detected in this study. Many researchers have reported that haplotype Ia is dominant in most of the populations of *P. infestans* studied (Shimelash et al. 2016; Fry et al. 1993). The proportion of each mtDNA haplotype differed in the different regions in China. Haplotype IIa dominated the population of *P. infestans* in north-eastern China, growing faster than the other two haplotypes (IIa and Iib) on rye B agar medium (Tian et al. 2018).

## Conclusion

The population of *P. infestans* in southern China is diverse and complex. Seventy-six percent of the tested isolates were self-fertile, 80% of the tested isolates were metalaxyl-resistant, and 26.58% of the tested isolates were members of the most common physiological race 1.2.3.4.5.6.7.8.9.10.11. This suggests that metalaxyl-based fungicides should be carefully used and that resistance to metalaxyl should be monitored in southern China. The control strategies for potato late blight should be organically combined with the different cultivation models, which contribute to maintaining biodiversity and suppressing the occurrence of late blight.

## Acknowledgements

This work was funded by the Sichuan Science and Technology Program (2016NYZ0053), which was also supported by the Institute of Plant Protection, Sichuan Academy of Agricultural Sciences.

## References

- Aav A, I Skrabule, G Bimšteine, T Kaart, IH Williams, E Runno-Paurson (2015). The structure of mating type, metalaxyl resistance and virulence of *Phytophthora infestans* isolates collected from Latvia. *Zemdirbyste-agriculture* 102:335–342
- Alva A, MS Fan, C Qing, C Rosen, HQ Ren (2011). Improving nutrient-use efficiency in Chinese potato production: Experiences from the United States. *J Crop Improv* 25:46–85
- Andersson B, M Sandström, A Strömberg (1998). Indications of soil-borne inoculum of *Phytophthora infestans*. *Potato Res* 41:305–310
- Bimšteine G (2008). *Phytophthora infestans* populations in Latvia. *Proc. Latvian Acad Sci* 62:223–226

- Brylińska M, S Sobkowiak, E Stefańczyk, J Śliwka (2016). Potato cultivation system affects population structure of *Phytophthora infestans*. *Fungal Ecol* 20:132–143
- Casa-Coila VH, MSLehner, BTH Júnior, A Reis, NRX Nazareno, ESG Mizubuti, CB Gomes (2017). First report of *Phytophthora infestans* self-fertile genotypes in southern Brazil. *Plant Dis* 101:1682
- Chmielarz M, S Sobkowiak, K Dębski, DEL Cooke, MB Brurberg, J Śliwka (2014). Diversity of *Phytophthora infestans* from Poland. *Plant Pathol* 63:203–211
- Drenth A, ICQTas, F Govers (1994). DNA fingerprinting uncovers a new sexually reproducing population of *Phytophthora infestans* in the Netherlands. *Eur J Plant Pathol* 100:97–107
- Fernández-Pavía SP, NJ Grünwald, M Díaz-Valasis, M Cadena-Hinojosa, WE Fry (2004). Soil-borne oospores of *Phytophthora infestans* in central Mexico survive winter fallow and infect potato plants in the field. *Plant Dis* 88:29–33
- Forbes GA (1997). *Manual for laboratory work on Phytophthora infestans*. Centro Internacional de la Papa, Quito, Ecuador
- Fry WE, SB Goodwin (1997a). Re-emergence of potato and tomato late blight in the United States. *Plant Dis* 81:1349–1357
- Fry WE, SB Goodwin (1997b). Resurgence of the Irish potato famine fungus. *Bioscience* 47:363–371
- Fry WE, SB Goodwin, AT Dyer, JM Matuszak, A Drenth, PW Tooley, LS Sujkowski, YJ Koh, BA Cohen, LJ Spielman, KL Deahl, DA Inglis, KP Sandlan (1993). Historical and recent migrations of *Phytophthora infestans*: Chronology, pathways and implications. *Plant Dis* 77:653–661
- Fukue Y, S Akino, H Osawa, N Kondo (2018). Races of *Phytophthora infestans* isolated from potato in Hokkaido, Japan. *J Gen Plant Pathol* 84:276–278
- Griffith GW, DS Shaw (1998). Polymorphisms in *Phytophthora infestans*: Four mitochondrial haplotypes are detected after PCR amplification of DNA from pure cultures or from host lesions. *Appl Environ Microb* 64:4007–4014
- Han LL, C Yang, X Pan, JS Zhan, W Zhu (2014). Identity of the mtDNA haplotype(s) of *Phytophthora infestans* in Fujian province. *Acta Laser Biol Sin* 23:165–169
- Hannukkala AO, T Kaukoranta, A Lehtinen, A Rahkonen (2007). Late-blight epidemics on potato in Finland, 1933–2002; increased and earlier occurrence of epidemics associated with climate change and lack of rotation. *Plant Pathol* 56:167–176
- Harutyunyan SR, ZJ Zhao, TD Hartog, K Bouwmeester, AJ Minnaard, BL Feringa, F Govers (2008). Biologically active *Phytophthora* mating hormone prepared by catalytic asymmetric total synthesis. *Proc Natl Acad Sci USA* 105:8507–8512
- Haverkort AJ, PM Boonekamp, R Hutten, E Jacobsen, LAP Lotz, GJT Kessel, RGF Visser, EAG van der Vossen (2008). Societal costs of late blight in potato and prospects of durable resistance through cisgenic modification. *Potato Res* 51:47–57
- Hohl HR, K Iselin (1984). Strains of *Phytophthora infestans* from Switzerland with A2 mating type behaviour. *Trans Br Mycol Soc* 83:529–530
- Huang H (2002). Homothallic *Phytophthora infestans* was found in China. *Acta Phytopath Sin* 32:347–350
- Kiiker R, M Hansen, IH Williams, DEL Cooke, E Runno-Paurson (2018). Outcome of sexual reproduction in the *Phytophthora infestans* population in Estonian potato fields. *Eur J Plant Pathol* 152:395–407
- Lehtinen A, A Hannukkala (2004). Oospores of *Phytophthora infestans* in soil provide an important new source of primary inoculum in Finland. *Agric Food Sci* 13:399–410
- Li BJ, QH Chen, X Lv, CZ Lan, J Zhao, RZ Qiu, QY Weng (2009). Phenotypic and genotypic characterization of *Phytophthora infestans* isolates from China. *J Phytopathol* 157:558–567
- Li HH, HX Peng, YD Xi, XL Wang, BW Liu (2013a). Characteristics of mating types, physiological races, metalaxyl sensitivity and mtDNA haplotypes of *Phytophthora infestans* in Sichuan Province. *Sci Agric Sin* 46:728–736
- Li Y, T Lee, JH Zhu, GH Jin, CZ Lan, SX Zhu, RF Zhang, BW Liu, ZJ Zhao, G Kessel, SW Huang, E Jacobsen (2013b). Population structure of *Phytophthora infestans* in China – geographic clusters and presence of the EU genotype Blue\_13. *Plant Pathol* 62:932–942
- Lian YH, GJ Ye, J Wang (2012). Identification and analysis of mitochondrial haplotypes on potato *Phytophthora infestans* in Qinghai area. *Acta Phytopath Sin* 42:164–168
- Lozoya-Saldaña H (2011). Evolution of vertical and horizontal resistance and its application in breeding resistance to potato late blight. *Potato J* 38:1–8
- Malcolmson JF, W Black (1966). New R genes in *Solanum demissum* Lindl. A and their complementary races of *Phytophthora infestans* (Mont.) de Bary. *Euphytica* 15:199–203
- Niederhauser JS (1956). Division of mycology: The blight, the blighter and the blighted. *Trans NY Acad Sci* 19:55–63
- Pipe ND, VA Zcoitia, DSShaw (2000). Self-fertility in *Phytophthora infestans*. Heterokaryons segregate several phenotypes. *Mycol Res* 104:676–680
- Runno-Paurson E, WE Fry, KL Myers, M Koppel, M Mänd (2009). Characterisation of *Phytophthora infestans* isolates collected from potato in Estonia during 2002–2003. *Eur J Plant Pathol* 124:565–575
- Sheldon AL (1969). Equitability indices: Dependence on the species count. *Ecology* 50:466–467
- Shimelash D, T Hussien, C Fininsa, G Forbes, J Yuen (2016). Mitochondrial DNA assessment of *Phytophthora infestans* isolates from potato and tomato in Ethiopia reveals unexpected diversity. *Curr Genet* 62:657–667
- Śliwka J, S Sobkowiak, R Lebecka, J Avendaño-Córcoles, E Zimnoch-Guzowska (2006). Mating type, virulence, aggressiveness and metalaxyl resistance of isolates of *Phytophthora infestans* in Poland. *Potato Res* 49:155–166
- Smart CD, H Mayton, ESG Mizubuti, MR Willmann, WE Fry (2000). Environmental and genetic factors influencing self-fertility in *Phytophthora infestans*. *Phytopathology* 90:987–994
- Tian HY, JZ Jiang, CB Li, N Hou, F Shen, Y Li (2018). Identity of the mtDNA haplotype and biological characteristics of *Phytophthora infestans* in Northeast China. *J Agric Univ Hebei* 41:25–31
- Tian YE, JL Yin, JP Sun, YF Ma, QH Wang, JL Quan, WX Shan (2016). Population genetic analysis of *Phytophthora infestans* in northwestern China. *Plant Pathol* 65:17–25
- Wharton PS, P Nolte, WW Kirk, SDangi, A Gevens (2015). First Report of Late Blight Caused by *Phytophthora infestans* Clonal Lineage US-23 on Potato in Idaho. *Plant Dis* 99:417–417
- Whittaker SL, SJ Assinder, DS Shaw (1994). Inheritance of mitochondrial DNA in *Phytophthora infestans*. *Mycol Res* 98:569–575
- Yuen JE, B Andersson (2013). What is the evidence for sexual reproduction of *Phytophthora infestans* in Europe? *Plant Pathol* 62:485–491
- Zhang ZM, YQ Li, SM Tian, JH Zhu, J Wang, BF Song (1996). The occurrence of potato late blight pathogen (*Phytophthora infestans*) A2 mating type in China. *J Agric Univ Hebei* 19:62–65
- Zhao ZJ, JF Cao, CH Li, DW Sun, MY Yang, J Wang (2007). Characteristics of mating type, metalaxyl sensitivity, mtDNA haplotype and succession of *Phytophthora infestans* populations in Yunnan, 2000 to 2003. *Sci Agric Sin* 40:727–734
- Zhao ZJ, XP Li, CY Li (2002). Molecular determination on mitochondrial DNA haplotypes of *Phytophthora infestans* in Yunnan of China. *J Yunnan Agric Univ* 17:17–20
- Zhu W, LN Yang, EJ Wu, CF Qin, LP Shang, ZH Wang, JS Zhan (2015). Limited sexual reproduction and quick turnover in the population genetic structure of *Phytophthora infestans* in Fujian, China. *Sci Rep* 5; Article 10094