

Effects of Ewe Oil in Media for Mass Production of *Tricephalobus* (Nematoda: Rhabditida)

SHOKOOHI, E¹, A. KHEIRI, J. ZAD, M. FARZANEH AND M. JAVAN NIKKHAH

Plant Protection Department Faculty of Horticultural Science and Plant Protection, University College of Agriculture, University of Tehran, Karaj, Iran

¹Corresponding author's e-mail: eshokoohi@gmail.com

ABSTRACT

The Media WA 2%, WA 2% + Ewe oil, WA 2% + K₂HPO₄, SDA (Sabouraud 4% Dextrose Agar), SDA + K₂HPO₄, SDA + Ewe oil, NA, NA + K₂HPO₄ and NA + Ewe oil were used for mass production of *Tricephalobus*. One mL Ewe oil extracted from milk was added to the media 2% WA + Ewe oil and NA + Ewe oil and incubated at 25°C for one week. The media were monitored every 24 h. Juvenile and mature nematodes in WA 2% amended with Ewe oil and the other media were counted after one week. These results showed that the reproduction of nematodes in WA 2% + Ewe oil was much more in comparison with other media. The experiment was repeated three times. These results indicated that Ewe oil was a better medium than that of cholesterol for mass production. This is the first report of using Ewe oil in mass production of this nematode and this method is advisable for its mass production.

Key Words: *Tricephalobus*; Ewe oil; K₂HPO₄

INTRODUCTION

The ability to culture organisms is always of great benefit to their scientific study and sometimes simply essential. Culture ensure that a steady supply of live material is available for teaching and research, eliminating the need for repeated sampling of inaccessible or temporally fluctuating habitats in search of living specimen. *Caenorhabditis elegans* is by far the most widely cultured nematode species being maintained in systems ranging from tiny petri dishes to huge bio-reactors (Gbewonyo *et al.*, 1994). Many methods have been developed for axenizing nematodes (Krusberg & Sardanelli, 1984). The free living nematodes can be cultivated axenically in a chemically defined medium (Nicholas, 1962; Sayre *et al.*, 1963). The nematode belong to Pnagrolaimidae is a use-full bait for studying nematophagous fungi and has therefore been used for their isolation from soil (Dackman *et al.*, 1987). Freelifing nematode (*Panagrellus redivivus*) was axenically grown in neutralized soya peptone as in jansson and nordbring-hertz (1979). In the present study, Ewe oil was extracted from milk and its in reproduction and population growth of *Tricephalobus* was investigated.

MATERIALS AND METHODS

Sampling was carried out in the Faculty of Agriculture in Tehran Province, Iran. Nematodes were extracted using the funnel technique according to Beier and Traunspurger (2003) from leaf dust. The samples were sieved through a 38 micrometer sieve. In this study females of *Tricephalobus* sp were maintained on 2% WA in 100 mm diam plastic petri plates. Distilled water (5 - 10 mL) was added to the plates. Six media were prepared for mass production of nematodes as following: WA 2%, WA 2% + Ewe oil, WA

2% + K₂HPO₄, SDA (Sabouraud 4% Dextrose Agar), SDA + K₂HPO₄, SDA + Ewe oil, NA, NA + K₂HPO₄ and NA + Ewe oil. One mL of Ewe oil extracted from milk was added to WA 2% and NA. Three gravid females were added to each plate. Plates were viewed under a microscope one week after inoculation and the number of larvae and females were counted. To ensure that the nematodes in the cultures were *Tricephalobus* three or five adults were randomly picked and placed in a water mount during the experiment and identified. Experiments were arranged in a factorial design with four replicates. The data were analyzed with SAS software.

RESULTS AND DISCUSSION

Effect of these media on production of females and juveniles were significant ($P \leq 0.01$) (Table I & Table II). Results showed that the populations of nematodes in the media WA 2% + K₂HPO₄ and WA 2% + Ewe oil significantly increased over time, when starting with three nematodes. The population of nematode in media amended with Ewe oil in comparison with the media without Ewe oil was different at 1% level. The numbers of juvenile in media WA 2%, WA 2% + k₂HPO₄ and WA 2% + Ewe oil were more than the others. In the media NA, NA + Ewe oil and SDA the population was very low (Table III).

Results showed that WA 2% + k₂HPO₄ and WA 2% + Ewe oil with 289.25 and 609.50 larvae and 70.25 and 89.25 female were the best media for production of *Tricephalobus* and showed significantly differences at 1% level comparison with other media (Table III). The number of larvae and females in the media SDA, SDA + K₂HPO₄, SDA + Ewe oil, NA, NA + K₂HPO₄ and NA + Ewe oil were very low and could not show any significantly differences at 1% level with the others (Table III).

Table I. Analysis of variance of some media and amended additives on production of larvae of *Tricephalobus* after one week under dark condition at 25°C

Treatment	DF	Mean Square
Medium	2	813.200 ^{**}
Amended	2	106.018 ^{**}
Medium × Amended	4	125.090 ^{**}
Error	27	1.765

** Significantly different (1%)

Table II. Analysis of variance of some media and amended additives on production of females of *Tricephalobus* after one week under dark condition at 25°C

Treatment	DF	Mean Square
Medium	2	157.071 ^{**}
Amended	2	7.984 ^{**}
Medium × Amended	4	12.164 ^{**}
Error	27	0.454

** Significantly different (1%)

Table III. Effect of some media and amended additives on production rate of Larvae and Female of *Tricephalobus* after 1 week under dark condition at 25°C

Treatment ^a	Larvae	Female
WA	35.00 c ^b	16.00 b
WA+ K ₂ HPO ₄	289.25 b	70.25 a
WA+ Ewe oil	609.50 a	89.25 a
NA	1.50 d	1.00 c
NA+ K ₂ HPO ₄	1.00 d	0.00 c
NA+ Ewe oil	0.00 d	0.00 c
SDA	5.00 d	1.25 c
SDA+ K ₂ HPO ₄	2.25 d	0.75 c
SDA+ Ewe oil	3.75 d	0.75 c

^a WA, Water Agar 2%; Ewe oil, extracted from milk; NA, Nutrient Agar 2%; SDA, Sabouraud 4% Dextrose Agar 6.5%.^bMean values followed by the same letter are not significantly difference according to Duncan test at 1% significance level.

In this study adults and juveniles of *Tricephalobus* proved relatively easy to be maintained *in vitro* with media WA 2% + K₂HPO₄ and WA 2% + Ewe oil. Our results showed that *Tricephalobus* needs moisture as observed in preliminary culturing plates. The availability of a new source of nematode growth factor from Ewe oil extracted from milk provides a useful supplement for culture of free living nematodes and may be of value in culture of other organisms. This material (Ewe oil) can be used either as a supplement or as a complete medium. De ley (1998) used cholesterol in medium for *in vitro* culturing of free living nematodes. Cholesterol starvation on free living development at both embryonic and post-embryonic stages by examining brood size, embryonic lethality, growth rate and worm size (Yhong-Hee Shim *et al.*, 2002). Nematodes, including free-living *Caenorhabditis elegans*, require sterol for their growth as a nutritional source (Hieb & Rothstein, 1968). It is well known that some free living nematodes such as *C. elegans* is un-able to bio-synthesize sterol *de novo* (Rothstein, 1968). However, apparently *C. elegans* is capable of obtaining cholesterol or cholesterol-like sterol (e.g. 7-DHC) for their growth by metabolizing natural sterols

such as 24-alkyl sterol present in many plants (Chitwood *et al.*, 1995) or sterols from the animal body in soil. Our results showed that Ewe oil is easily available and can be a useful substitution for cholesterol as a lipid source for nematode. It is well established that exogenous factors such as temperature play an important role in altering the rate of development in nematodes (Tahseen *et al.*, 1994). The results showed temperature (25°C) had effective role in reproduction and population of *Tricephalobus*. Some sterol that already have been used for studying of Panagrolaimids nematodes played important role in growth and reproduction of this group, because of they are not able to synthesis the sterol. This demonstration also confirms a vital function in these nematodes (Cole & Dutky, 1969). As Ewe oil is the great source of cholesterol, it can be used for preparation of media for mass production of these nematodes. The ability to culture *Tricephalobus* allows further examination of the ecological, biological and molecular investigation free living and plant parasitic nematodes.

REFERENCES

- Beier, S. and W. Traunspurger, 2003. Seasonal distribution of free living nematodes in Krahenbach, a fine grained submountain carbonate stream in southwest Germany. *Nematology*, 5: 113–36
- Chitwood, D.J., W.R. Lusby, M.J. Thompson, J.P. Kochansky and O.W. Howar, 1995. The glycosylceramides of the nematode *Caenorhabditis elegans* contain an un-usual, branched-chain sphingoid base. *Lipids*, 30: 567–73
- Cole, R.J. and S.R. Dutky, 1969. A sterol requirement in *Turbatrix acetii* and *Panagrellus redivivus*. *J. Nematol.*, 1: 72–5
- Dackman, C., S.Olsson, H.B. Jansson, B. Lundgren and B. Nordbring-Hertz, 1987. Quantitative of predatory and endoparasitic nematofaphus fungi in soil. *Microbiol. Ecol.*, 13: 89–93
- De Ley, P., I.T. De Ley and M. Mundo-Ocampo, 1998. *Identification of Free-living Secernentea*. Nematode workshop, Department of Nematology, University of California, Riverside
- Gbewonyo, K., S.P. Rohre, L. Lister, B. Burgess, D. Cully and B. Buckland, 1994. Large scale cultivation of the free living nematode *Caenorhabditis elegans*. *Biotechnology*, 12: 51–4
- Hieb, W.F. and M. Rothstein, 1968. Sterol requirement for reproduction of a Free-living nematode. *Science*, 160: 778–80
- Jansson, H.B. and B. Nordbring-Hertz, 1979. Attraction of nematodes to living mycelium of nematophagus fungi. *J. Gen. Microbiol.*, 112: 89–93
- Krusberg, L.R. and S. Sardanelli, 1984. Technique for axenizing nematodes. *J. Nematol.*, 16: 384
- Nicholas, W.L., 1962. A study of species of Acrobeloides (Cephalobidae) in laboratory culture. *Nematologica*, 8: 99–109
- Rothstein, M., 1968. Nematode biochemistry. IX Lack of sterol biosynthesis free-living nematodes. *Comp. Biochem. Physiol.*, 27: 309–17
- Sayre, F.W., E.L. Hansen and E.A. Yarwood, 1963. Biochemical aspects of the nutrition of *Caenorhabditis briggsae*. *Exp. Parasitol.*, 13: 98–107
- Tahseen, Q., M.S. Jairajpouri and I. Ahmad, 1994. Factors affecting hatching of the soil nematode *Diploscapter orientalis*. *Nematol. Medit.*, 22: 221–3
- Yhong-Hee Shim, J.U. Hee Chun, L. Eun-Young and Young-Kipaik, 2002. Role of Cholesterol in Germ-line Development of *Caenorhabditis elegans*. *Molecular Reproduction and Development*, 61: 358–66

(Received 16 June 2006; Accepted 20 October 2006)