



Full Length Article

Effects of Ozone Stress on Rice Growth and Yield Formation under Different Planting Densities - A Face Study

Bin Peng^{1,2}, Yunxia Wang³, Jianguo Zhu⁴, Yulong Wang² and Lianxin Yang^{2*}

¹Jiangsu Key Laboratory of Biochemistry and Biotechnology of Marine Wetland, Yancheng Institute of Technology, Yancheng 224051, PR China

²Jiangsu Key Laboratory of Crop Genetics and Physiology/Co-Innovation Center for Modern Production Technology of Grain Crops, Yangzhou University, Yangzhou 225009, PR China

³College of Environmental Science and Engineering, Yangzhou University, Yangzhou 225009, PR China

⁴State Key Laboratory of Soil and Sustainable Agriculture, Institute of Soil Science, Chinese Academy of Sciences, Nanjing 210008, PR China

*For correspondence: hl1343@163.com; ycitbin@163.com; chao.jiang@advancesin.org

Abstract

The rising tropospheric ozone concentration inhibits rice growth and causes yield loss, but whether ozone effects can be ameliorated by cultivation measures such as planting density regulation was unclear, especially under fully open-air field conditions. An experiment was conducted with rice to address this issue by using free-air gas concentration enrichment (FACE) facility stored in paddy field of China. A conventional indica cultivar Yangdao 6 (2011) and a super hybrid indica cultivar IY084 (2012) were grown at ambient or elevated ozone concentration (target at 50% above ambient, E-O₃) under low (LD, 16 hills m⁻²), medium (MD, 24 hills m⁻²) or high planting density (HD, 32 hills m⁻²) from tillering until maturity. E-O₃ greatly accelerated phenological development and reduced plant height of rice LY084. The results from the final harvest showed that E-O₃ reduced grain yield of Yangdao 6 and IY084 by 16 and 27%, respectively. The yield loss was attributed to a great reduction in sink size (the number of spikelets per panicle or unit area, -22%), secondly the decline in grain-filling capacity shown by decreases in filled grain fraction (-13%) and filled grain weight (-5%) but increases in empty and lightened grain fractions. Although yield components were regulated by changing planting density in general, no significant ozone effect by planting density interaction was found for any of the measured parameters, which indicated that planting measures could not ameliorate ozone damage, at least under the current experimental conditions. It was concluded that yield loss of hybrids were greater than conventional rice cultivars, the future projection must include hybrid rice or/and super rice not to underestimate the effects of ozone stress on rice productivity. © 2018 Friends Science Publishers

Keywords: Rice; Ozone; FACE (Free Air ozone Concentration Enrichment); Planting density; Yield

Introduction

In many Asian countries including China and India, the rapid economic development and population growth in recent years has led to increased emissions of air pollutants (NO_x) and volatile organic compounds (VOC), these precursors are converted to ozone in photochemical reactions, which resulted in a rapid rise in tropospheric ozone concentrations (Ohara *et al.*, 2007; Fowler *et al.*, 2008; Emberson *et al.*, 2009; Lei *et al.*, 2012).

Ozone is a strong oxidant and enters plant through leaf stomata and decomposed into active oxygen in the apoplast (Kobayashi *et al.*, 1995; León-Chan *et al.*, 2017). These reactive oxygen damage protein, lipids and DNA, or trigger the programmed cell death, cause leaf damage and lead to reduced photosynthesis, growth inhibition and yield loss (Kobayashi *et al.*, 1995; Fiscus *et al.*, 2005; Ainsworth *et al.*,

2012). Rice is one of the major staple in South and East Asia and rice cultivation regions often experience high ozone concentration (Ohara *et al.*, 2007; Alexandratos and Bruinsma, 2012; Frei, 2015). Furthermore, meta-analysis has indeed demonstrated that rice is not less sensitive to O₃ than other sensitive species such as soybean and wheat (Ainsworth, 2008; Feng and Kobayashi, 2009). Therefore, it is important to study the effects of ozone stress on rice growth, yield and its mechanism, and to develop appropriate response measures to ensure regional food security in the context of global climate change. The responses of rice growth and yield formation under ozone stress have been reported extensively, but most of these studies are single factor tests focusing on ozone effects only (Ainsworth, 2008; Yang *et al.*, 2008). The limited number of multi-factorial experiments investing the interactions between ozone and other factors has demonstrated that the ozone effects on rice

yield varied with fumigation methods, cultivars and environmental conditions (Olszyk and Wise, 1997; Ainsworth, 2008; Wang *et al.*, 2014; Frei, 2015), which justify the fact that understanding ozone effect in field requires more multifactor tests. Rice production management measures including planting density, fertilizer application and water management are essential for achieving high yield. Whether proper management measures can modify ozone stress was unclear, because such research requires larger test space, and better conducted in natural field. But most of the previous studies were conducted in closed or partial closed chambers with relatively small space (diameter less than 2 m). In contrast to chamber, Free Air gas Concentration Enrichment (FACE) platform has a relatively larger operation space (diameter more than 14 m), is closer to the real-world situation than chamber, provides better opportunity for studying interactions between ozone and other factors (Long *et al.*, 2005; Ainsworth *et al.*, 2012). In 2007, for the first time with rice in the world, the full-size (14-m diameter) FACE system was set up in farmer fields in Jiangsu Province, China (Tang *et al.*, 2011).

Because of its great yield potential and strong stress resistance in general, hybrid rice has become the rice type with the largest planting area in China (>60%), plays an important role in ensuring food security (Yuan, 1999). However, in our previous studies, we found the yield loss of hybrid rice under ozone stress that was significantly higher than conventional rice (Shi *et al.*, 2009; Wang *et al.*, 2012). This brings pessimism for the future rice production if ozone level can't be controlled during rice growth period. Therefore, finding proper management measures targeting ozone damage is an urgent issue to secure rice production. Optimizing planting density is an important field management technology, which regulates crops yield and quality. Changing planting density will modify the robustness of individual plant as well as population quality. Meanwhile, changing planting density also changes the free space for individual plant and it may influence stomatal ozone uptake of leaves, especially those at the lower position of a plant. These factors together will determine the final yield response of rice to ozone. The objective of this O₃-FACE experiment is to study the rice yield response to E-O₃ under different planting densities, and its relation with the growth processes. Understanding this interaction, especially under a natural field conditions, will help us to formulate adaptation strategies of rice production systems under high [O₃] in the near future.

Materials and Methods

Research Site and ozone-FACE Platform

The experiment was conducted in the China O₃-FACE platform, located at Xiaoji town, Jiangdu County, Jiangsu province, China (119°45' E, 32°35' N). The site, at 5 m above the sea level in elevation, sits in the subtropical marine

climatic zone. Throughout the period of 113 days growing period from June 21 to October 17 for the experiment in 2012, mean daily temperature was 24.5°C, mean daily integral solar radiation (or PAR) was of 13.9 MJ m⁻², and mean daytime (6:00–18:00) vapor pressure deficit was of 0.67 kPa (Tang-11). This site has been in continuous cultivation for more than 1000 years with rice–wheat or rice–rapeseed rotation. Detailed descriptions of the soil properties for the site can be found in our previous publication (Shi *et al.*, 2009). Details of the design and performance of this O₃-FACE system are provided by Tang-11. In brief, the FACE system has eight 240 m² plots, of which replicate plots were exposed to elevated ozone concentration (hereinafter called E-O₃ plots) and four equal size plots were in ambient ozone concentration (A-O₃ plots). Any one of the E-O₃ plots was separated from other plots by at least 70 m to avoid cross contamination. The quantity and direction of the ozone release for each E-O₃ plot was controlled by a proportional integral derivative algorithm for computer feedback that compares achieved ozone concentration to the target ozone concentration of 1.5 times ambient ozone concentration with an ozone monitor (model 49i, Thermo Environmental Instruments, MA, USA), a data logger-controller, an anemometer and a wind vane. A mixed gas consisting of about 5% ozone and 95% O₂ was produced by an ozone generator (KCF-BT0.2, Jiangsu Koner Ozone Co., Ltd., Yangzhou, China). The mixed gas was released in a stream of compressed air into the plots through the ABS pipes positioned at about 50 above the canopy. In the ambient plots, plants were grown under ambient ozone conditions without the ring structures. The ozone fumigation began on 1st July and continued throughout the rice season until harvest except for occasions such as system malfunction, rain or wet leaves. The fumigation was settled during day-time 7-h (09:00–16:00 Chinese Standard Time) every day. Fig. 1 shows the seasonal change in daily 7-h mean ozone concentration for A-O₃ and E-O₃ plots.

Plant Material and Cultivation

The cultivars were Yangdao 6 (2011) and II You 084 (2012). Rice seeds were sown on 21 May; on which the seedlings grown under ambient air, and manually transplanted into the A-O₃ and E-O₃ plots with one seedling per hill on 21 June 2012. Planting density was 16, 24 and 32 hills m⁻² for low, medium and high density, equivalent to the spacing of hills 25×25, 16.7×25 and 12.5×25, respectively. Nitrogen (N) was supplied as urea (N = 46%) and compound chemical fertilizer (N: P₂O₅: K₂O = 15:15:15) at a rate of 15 g N m⁻². Of the total N, 50% was applied as the basal dressing on 20 June, 10% was top-dressed on 7 July (tillering stage) and 40% top-dressed on 7 August (panicle initiation stage). Both phosphorus (P) and potassium (K) were applied as compound chemical fertilizer at equal rates of 7 g m⁻² as the basal dressing at 1 d before transplanting. Standard cultivation practices as commonly performed in the area

were followed in all experimental plots. In brief, the paddy fields were submerged with water of about 3 in depth from 21 June to 4 July, and then the fields were subjected to wet–dry cycles through natural drainage and intermittent irrigation. At 10 d before harvest, irrigation was terminated to allow paddy fields to dry for final harvesting. Pesticides and fungicides were applied when necessary throughout the experiment.

Plant Phenology, Height and Tillering Dynamics

The date for heading when 50% of plant headed and grain maturity was recorded in each subplot. Plant height and tiller numbers were measured for fixed 10 hills in each treatment at 5–10 days intervals until heading. The productive tiller ratio (%) was expressed as the panicle number per square meter/ the maximum tiller number per square meter x 100.

Grain Yield and Yield Components

Actual grain yield was determined of 60 hills from a 4, 3, and 2 m² patch in each LD, MD and HD subplot, respectively. The grains were air dried and weighted to obtain actual grain yield.

Grain yield components, *i.e.*, the number of panicles per square meter, the number of spikelets per panicle, filled spikelet fraction, and individual grain mass, were determined at maturity with six hills per plants in each subplot. Grains were soaked in tap water (specific gravity=1.0) and the number of sunken and floated grains was counted to determine the filled spikelet fraction. The floated grains were dried and then separated into empty and lightened grains, each of which was counted and the empty grain percentage and lightened grain percentage were calculated. The sunken (filled) grains were oven-dried at 80°C until constant weight achieved. The theoretical grain yield was obtained by adjusting dry weight of filled grains to a moisture content of 0.14 g H₂O g⁻¹ fresh weight. Average grain mass = Total grain mass × 1000 / Total grain number.

Statistical Analysis

A split-plot design was employed with [O₃] as main-plot treatment and planting density as the split-plot treatment. Analysis of variance (ANOVA) was performed using the software Statistical Product and Service Solutions (SPSS Inc., Chicago, IL, USA) to determine the main effects of [O₃], planting density and the effects of their interaction. Treatments were compared by Tukey's test and differences were declared statistically significant ($P < 0.05$). Pearson's correlations were calculated to determine the relationships between the different parameters of rice quality.




Fig. 1: Seasonal change in daily 7-h (9:00–16:00 Chinese Standard Time) mean [O₃] for A-O₃ and E-O₃




Fig. 2: Ozone stress obviously accelerated plant growth period of LY084 under FACE condition. (a: fructificative period of LY084 to ambient O₃ concentration; b: fructificative period of LY084 to elevated O₃ concentration)

Results

Effect of E-O₃ on Phenology and Plant Height

E-O₃ greatly accelerated phenological development of rice LY084. The period from transplanting to heading stage and from heading to maturity stage were reduced by 3 and 4 days, respectively, resulting in one-week earlier maturity by E-O₃ (Fig. 2). The reduction in plant height by E-O₃ increased with plant growth process: with no effect on plant height at tillering stage, but reduced it by 3.1% (2.4, $P < 0.1$), 6.7% (7.0, $P < 0.01$) and 7.8% (8.6, $P < 0.01$), at jointing, heading and maturity stages, respectively (Table 1). No effect of planting density or O₃ by planting density interaction was detected on phenology and plant height.

Effect of E-O₃ on Grain Yield

Final theoretical (TGY) and actual grain yield (AGY) of hybrid LY084 showed that increasing planting density from 16 to 24 hill m⁻², TGY and AGY increased by 14.5% and 5.9%, respectively but higher density level of 32 hill m⁻² did not result in further increases in grain yield (Fig. 3). E-O₃ greatly reduced the TGY and AGY of LY084.

Table 1: Plant height (cm) at different growth stages of hybrid LY084 exposed to ambient (A-O₃) or elevated O₃ concentration (E-O₃, ambient×1.5) under low (LD, 16 hills m⁻²), medium (MD, 24 hills m⁻²) and high (HD, 36 hills m⁻²) levels of planting density

Density	Ozone	Tillering stage	Jointing stage	Heading stage	Maturity Stage
LD	A-O ₃	67.3 ± 1.3	93.6 ± 2.0	105.4 ± 1.8	109.7 ± 1.6
	E-O ₃	68.8 ± 1.6	90.4 ± 2.0	95.4 ± 1.7	100.0 ± 0.5
MD	A-O ₃	70.0 ± 1.4	92.5 ± 1.3	105.7 ± 2.5	111.1 ± 1.8
	E-O ₃	69.1 ± 1.0	91.5 ± 1.4	99.5 ± 0.8	103.4 ± 2.6
HD	A-O ₃	69.6 ± 1.1	99.0 ± 1.8	104.4 ± 3.1	109.3 ± 2.4
	E-O ₃	68.1 ± 1.1	95.9 ± 2.0	99.4 ± 1.6	100.9 ± 2.9
ANVOA	O ₃	0.747	0.139	0.001	<0.001
	Density	0.487	0.017	0.572	0.360
	O ₃ *Density	0.472	0.815	0.458	0.869

A-O₃: Ambient O₃ concentration; E-O₃: Elevated O₃ concentration




Fig. 3: Actual grain yield (A) and theoretical grain yield (B) at final harvest of hybrid LY084 exposed to ambient (A-O₃) or elevated O₃ concentration under low (LD), medium (MD) and high (HD) levels of planting density. These data are from harvests of ca. 60 plants (ca. 2-4 m²) within each replicate plot at maturation of the crop. Bars indicate ± SE (n=4). Bars not sharing the same letter differ significantly at *p* < 0.05

Averaged across 3 densities, E-O₃ decreased TGY from 1111 g m⁻² in the controls to 733 g m⁻² (34.0%) and AGR from 1018 g m⁻² in the controls to 739 g m⁻² (27.4%). Similar patterns of yield response to O₃ were observed for rice grown in different planting density. TGY was lower by 32.4%, 36.5% and 32.9%, and AGY by 26.9%, 25.0% and 30.0% for LM-, MM- and HM-crops, respectively showing no interaction of ozone with planting density for the two parameters. O₃ treatment decreased the actual grain yield of Yangdao 6 by 142.0 g m⁻², with 15.9% (*p* < 0.01) compared to the control, with the decrease of 23.5% (*p* < 0.01), 14% (*p* < 0.05) and 9.9% (*p* < 0.05) under low, medium and high density (Fig. 4).

Effect of E-O₃ on Yield Components

Significant effect of E-O₃ was detected for most yield parameters, however, no interaction between O₃ and planting density was observed for any of the yield parameters.

The number of panicles per unit land area increased with the increase of plant density, but showed no response

to E-O₃ (Fig. 5a). Panicle number is the product of maximum tiller number per square meter (MTN) and productive tiller ratio (PTR). The latter was unaffected by planting density while the former increased as planting density increased (Table 2). E-O₃ influenced the two components in opposite directions: MTN was significantly reduced by 5-11% while PTR increased by a similar amount under ozone stress for all density levels (Table 2).

Maximum tiller number, productive tiller ratio, dry weight per stem at heading and ratio of spikelet number to stem dry weight of hybrid LY084 exposed to ambient (A-O₃) or elevated O₃ concentration (E-O₃, ambient×1.5) under low (LD, 16 hills m⁻²), medium (MD, 24 hills m⁻²) and high (HD, 36 hills m⁻²) levels of planting density.

The number of spikelets per panicle was not changed by density, but responded significantly and negatively to ozone stress (Fig. 5b). E-O₃ significantly decreased the spikelet number per panicle by 22% on average, with the decrease of 16%, 28% and 22% under LD, MD and HD treatments, respectively. The spikelet number per panicle can be factorized into dry weight per stem and the ratio of spikelet number to stem dry weight (Table 2). Averaged across all planting density, E-O₃ significantly decreased dry weight per stem and the ratio of spikelet number to stem dry weight by 13% and 11%, with the range of decrease being 9-17% and 5-14%, respectively (Table 2). Although planting density affected the spikelet number per panicle, and it had no impact on its two components.

Similar to the number of panicles, the spikelet number per unit area increased with the increase of planting density, 13% and 20% increases were found for MD and HD compared with LD (Fig. 5c). E-O₃ decreased the spikelet number per unit land area by 22% (*P* < 0.01) on average, with 20% (*P* < 0.01), 26% (*P* = 0.01) and 19% (*P* = 0.08) decrease at LD, MD and HD, respectively.

There was a trend of decrease in filled grain fraction, filled grain mass and average grain mass with the increase of planting density, and the effects were statistically significant for the latter two parameters (Fig. 5d, e and f). Averaged across all planting densities, E-O₃ decreased filled grain fraction, filled grain mass and average grain mass by 12, 5 and 10% (*P* < 0.01), respectively. Ozone-induced reduction among different planting densities was very close for each parameter, in the range of 11-13%, 4-5% and 8-11% for filled grain fraction, filled grain mass and average grain mass, respectively. With respect to lightened grain fraction (Fig. 5g) and empty grain fraction (Fig. 5h), planting density had no significant influence, but E-O₃ substantially increased these two fractions with 74% and 253% for lightened and empty grain fraction, respectively, in the range of 53-88% and 180-477% at different planting densities.

Discussion

In previous chamber studies, shortened or unchanged rice growth period by ozone stress was reported (Yang *et al.*,

Table 2: Maximum tiller number per square meter, productive tiller ratio, dry weight per stem at heading and ratio of spikelet number to stem dry weight of hybrid LY084 exposed to ambient (A-O₃) or elevated O₃ concentration (E-O₃, ambient×1.5) under low (LD, 16 hills m⁻²), medium (MD, 24 hills m⁻²) and high (HD, 36 hills m⁻²) levels of planting density

Density	Ozone	Maximum tiller number per m ²	Productive tiller ratio (%)	Dry weight per stem at heading stage (g)	Ratio of spikelet number to stem dry weight
LD	A-O ₃	372.8 ± 4.0	84.1 ± 0.9	3.1 ± 0.1	39.6 ± 1.1
	E-O ₃	344.8 ± 2.9	87.4 ± 1.2	2.8 ± 0.2	37.7 ± 1.9
MD	A-O ₃	431.5 ± 5.0	76.5 ± 1.0	3.3 ± 0.2	42.1 ± 2.3
	E-O ₃	383.5 ± 4.1	87.4 ± 1.0	2.7 ± 0.1	36.6 ± 1.9
HD	A-O ₃	460.5 ± 5.2	75.7 ± 1.3	3.0 ± 0.3	44.4 ± 2.5
	E-O ₃	438.0 ± 5.2	83.8 ± 0.9	2.7 ± 0.2	38.3 ± 2.2
ANOVA (p value)	O ₃	<0.001	<0.001	<0.001	<0.001
	Density (D)	<0.001	<0.001	0.199	0.068
	O ₃ ×D	0.026	0.007	0.278	0.167

Table 3: Summary of FACE results Summary of experimental background, absolute response in phenology and relative responses (%) in plant height, grain yield and its components of rice grown at E-O₃ relative to A-O₃ in the Chinese Ozone-FACE experiments

Items	2007	2008	2012
Test cultivars	WJ15 (or WYJ3), YD6, SY63, LYPJ	, SY63	LY084
Genotypes	Inbred japonica, Inbred indica, Hybrid indica, Hybrid japonica	Inbred indica, Hybrid indica	Hybrid indica (Super rice)
N application levels (g m ⁻²)	15	20	15
Planting density (hills m ⁻²)	24	24	16, 24 or 32
Phenology (d)			
Heading	ns, ns, -4 d , ns	ns, -4 d **	-3 d
Maturity	-4 (-2), -1, -8, -1 d	ns, -7 d **	-7 d
Plant height at maturity	ns, ns, 3-5 cm	ns, -10 cm **	-9 cm
Grain yield	ns, ns, -18-21%* , -15%*	ns, -27%* **	-28%* **
Grain yield components			
Panicle number per m ²	ns, ns, ns, ns	ns, ns	ns
Spikelet number per panicle	ns, ns, -12%* , -18%* **	-6%+ , 24%* **	-22%* **
Spikelet number per m ²	ns, ns, -16%* ***, -16%* **	-7%+ , 29%* **	-22%* **
Filled grain fraction	ns, ns, ns, ns	ns, ns	-12%* **
Filled grain mass	-1.3 mg , ns, ns, ns	ns, ns	-5%* **
Data sources	Pang <i>et al.</i> (2009); Shi <i>et al.</i> (2009)	Wang <i>et al.</i> (2012)	Tables or Figures in this paper

ns, no significance; +, $P < 0.1$; *, $P < 0.05$; **, $P < 0.01$

2008). In the present FACE study, heading and maturity was 3 and 7 days earlier for super rice LY084 under elevated O₃ conditions (25% higher than the ambient) (Fig. 2). This change in plant phenology was reported for hybrid indica SY63, but significantly greater than the other c cultivars (Shi *et al.*, 2009; Table 3).

The number of panicles had been reported to increase, decrease or unchanged under E-O₃ (Yang *et al.*, 2008). When averaged across all enclosure studies, the number of panicles was hardly affected by ozone (Ainsworth, 2008). Current free-air experiment indicated no change in panicle number. Little effect on panicle number was due to the offset of its two determinants: MTN (refer to plant capacity in forming tillers) decreased but PTR (refer to plant capacity in forming panicles) increased with E-O₃. Similar increases in PTR with E-O₃ have also been reported elsewhere (Wang *et al.*, 2012), though decrease or unchanged was reported too (Yang *et al.*, 2008; Shi *et al.*, 2009).

In contrast with panicles, the spikelet number per panicle showed a clear trend of decrease in response to ozone stress, and the magnitude of decrease varied with

varieties in the previous chamber studies (Yang *et al.*, 2008). The present FACE study found that E-O₃ decreased the spikelet number per panicle of LY084 independent of planting density (-22%, Fig. 5b), and this response was close to hybrids but substantially greater than conventional rice cultivars in previous FACE studies (Table 3). This large reduction in panicle size was supported by the decreases in the dry weight per stem and the ratio of spikelet number to stem dry weight (Table 2), as well as plant height at heading (Table 1). Our previous FACE study indicated that O₃-induced reduction in number of spikelets resulted from both the inhibition of spikelet differentiation and promotion of spikelet degeneration (Wang *et al.*, 2012). Spikelets per unit land area are the product of panicle number per unit area and spikelets per panicle, representing sink size of rice. Since, E-O₃ did not change panicle number; therefore, O₃-induced reduction in total sink size was driven almost totally by the decrease in panicle size rather than panicle number.

Meta-analysis on rice showed a trend of decrease in grain mass by E-O₃ (Ainsworth, 2008; Feng and Kobayashi, 2009). In the present study, filled grain fraction, filled grain




Fig. 4: Actual grain yield at final harvest of conventional variety yangdao 6 exposed to ambient (A-O₃) or elevated O₃ concentration under low (LD), medium (MD) and high (HD) levels of planting density (the experiment in 2011)




Fig. 5: Yield components at final harvest of hybrid LY084 exposed to ambient (A-O₃) or elevated O₃ concentration under low (LD), medium (MD) and high (HD) levels of planting density. Yield attributes included panicle number per m² (A); spikelet number per panicle (B), spikelet number per m² (C), fulfilled grain fraction (D), and fulfilled grain mass (E), average grain mass (F), lightened grain fraction (G) and empty grain fraction (H). Bars indicate ± SD. Bars not sharing the same letter differ significantly ($p < 0.05$)

mass and mean grain mass was found to decrease with the ozone stress, with the negative effect being more prominent for the filled-grain fraction, which confirmed the more growth suppression by E-O₃ after heading. The reduction in filled-grain fraction was much greater in the present study compared with the results from the previous rice-FACE

studies (Pang *et al.*, 2009; Shi *et al.*, 2009; Wang *et al.*, 2012; Young-Mei *et al.*, 2017), which reflected the high ozone sensitivity of LY084, especially at late growth stages. Low filled-grain fraction results from more empty or lightened grains in a panicle. The increase in empty grain percentage, especially lightened grain percentage (Fig. 5) was consistent with the down-regulation of leaf photosynthesis and plant growth in the grain-filling stage (Wang *et al.*, 2012). Ozone-induced increase in the percentage of grain sterility of rice was also reported elsewhere (Ishii *et al.*, 2004). The above results indicate that E-O₃ not only reduced the sink size of LY084, it also decreased grain-filling capacity of individual spikelet. These negative effects of ozone could not be eliminated by regulating planting density. Therefore, how ozone stress affects the processes of rice fertilization and grain filling requires further research, and special attention should be given to super rice.

Elevated O₃ decreases rice grain yield was found in most cases of previous studies. A meta-analysis of chamber studies by Ainsworth (2008) suggested a 14% decrease in grain yield, when exposed to 62 ppb ozone compared with plants grown in clean air. In this FACE study, a 25% increase from the ambient O₃ ca. 38 ppb to ca. 47 ppb caused a 27% yield loss of hybrid LY084, which substantially greater than the expectations from chamber studies (Ainsworth, 2008; Feng and Kobayashi, 2009) and also greater than the response of conventional rice cultivars yangdao 6, but very close to hybrids (Table 3). The significant decreases in yield components of spikelets per unit area, filled-grain fraction, grain mass, significant increase of empty and lightened grain percentage (Fig. 5), indicates the substantial yield loss of LY084 at E-O₃ was related with ozone-induced inhibition in spikelets formation, fertilization and grain filling processes, and its inhibition was greater than conventional rice cultivars (Table 3).

We didn't find any regulation effect of planting density on the response of rice to ozone stress, and ozone by density interaction was not detected for any parameters measured. Based on growth dynamics of this cultivar and the previous ozone studies on rice, we give the following explanation that higher planting density reduced the air contact of leaves at the lower position of plant population which led to lower ozone contact and uptake of these leaves, thereby amelioration of ozone damage. On the other hand, higher planting density affected the growth and development of the stem, especially in grain filling stage, which made the population quality worse, especially for the rice varieties with strong tillering ability and thin stems, and enhanced exacerbating the effects of E-O₃. These opposite effects might counteract each other, resulted in no regulation effect on ozone damage by changing planting density in this experiment.

Conclusion

To our knowledge, it was the first time that the interactive

effects between ozone and planting density on rice growth and yield were studied under fully open-air field conditions. The results showed that E-O₃ decreased the final productivity of super rice LY084, and the reduction was unaffected by planting density, as reflected in no interaction of ozone with planting density. The O₃-induced yield loss was mainly associated with the depression in reproductive growth, as shown by the negative effects on spikelets formation, fertilization and grain filling processes. The results from the present study provide important information for model predication and adaptation strategies formulation of future rice production. Firstly, because the yield loss of hybrids was much greater than conventional rice cultivars, the future prediction must include hybrids and/or super rice to avoid the underestimation of ozone stress on rice production. Secondly, the important adaptation strategy for hybrid rice to grown under ozone stress is to avoid photosynthesis depression at late growth stages in order to counteract the O₃-induced inhibition of spikelet and grain formation, and thus minimize the yield loss under E-O₃. The present study showing no clear regulation of changing planting density on ozone effects in rice converted an alarm that little potential for crop management practices to reduce the ozone damage; therefore, other adaptation strategies like developing tolerant cultivar need be reinforced.

Acknowledgements

This work was funded jointly by the National Natural Science Foundation of China (Grants Nos. 31371563, 31471437 and 31071359), the Open Fund of the State Key Laboratory of Soil and Sustainable Agriculture (Grants Nos. 0812201233), the Knowledge Innovation Program of Chinese Academy of Sciences (Grants Nos. KZCX2-EW-414), the Natural Science Foundation of Jiangsu Province (Grants Nos. BK20151298) and the Priority Academic Program Development of Jiangsu Higher Education Institutions. The FACE system instruments were supplied by the National Institute of Agro-Environmental Sciences and the Agricultural Research Center of Tohoku Region (Japan). We are grateful to Dr. G. Liu and H.Y. Tang for their technical support in the free-air ozone release system.

References

- Ainsworth, E.A., 2008. Rice production in a changing climate: a meta-analysis of responses to elevated carbon dioxide and elevated ozone concentration. *Glob. Change Biol.*, 14: 1–9
- Ainsworth, E.A., C.R. Yendrek, S. Sitch, W.J. Collins and L.D. Emberson, 2012. The effects of tropospheric ozone on net primary productivity and implications for climate change. *Annu. Rev. Plant Biol.*, 63: 637–661
- Alexandratos, N. and J. Bruinsma, 2012. *World Agriculture Towards 2030/2050: the 2012 Revision*. ESA Working Paper No., 12–03
- Emberson, L.D., P. Büker, M.R. Ashmore, G. Mills, L.S. Jackson, M. Agrawal, M.D. Atikuzzaman, S. Cinderby, M. Engardt, C. Jamir, K. Kobayashi, N.T.K. Oanh, Q.F. Quadir and A. Wahid, 2009. A comparison of North American and Asian exposure-response data for ozone effects on crop yields. *Atmos. Environ.*, 43: 1945–1953
- Feng, Z. and K. Kobayashi, 2009. Assessing the impacts of current and future concentrations of surface ozone on crop yield with meta-analysis. *Atmos. Environ.*, 43: 1510–1519
- Fiscus, E.L., F.L. Booker and K.O. Burkley, 2005. Crop responses to ozone: uptake, modes of action, carbon assimilation and partitioning. *Plant Cell Environ.*, 28: 997–1011
- Fowler, D., M. Amann, R. Anderson, M. Ashmore, P. Cox, M. Depledge, D. Derwent, P. Grennfelt, N. Hewitt, O. Hov, M. Jenkin, F. Kelly, P. Liss, M. Pilling, J. Pyle, J. Slingo and D. Stevenson, 2008. *Ground-level ozone in the 21st century: future trends, impacts and policy implications*. Royal Society Policy Document 15/08, vol. 15/08, RS1276 edition, The Royal Society, London
- Frei, M., 2015. Breeding of ozone resistant rice: relevance, approaches and challenges. *Environ. Pollut.*, 197: 144–155
- Ishii, S., F.M. Marshall and J.N.B. Bell, 2004. Physiological and morphological responses of locally grown Malaysian rice cultivars (*Oryza sativa* L.) to different ozone concentrations. *Water Air Soil Pollut.*, 155: 205–221
- Kobayashi, K., M. Okada and I. Nouchi, 1995. Effects of ozone on dry matter partition and yield of Japanese cultivar of rice (*Oryza sativa* L.). *Agric. Ecosyst. Environ.*, 53: 109–122
- Lei, H., D.J. Wuebbles and X.Z. Liang, 2012. Projected risk of high ozone episodes in 2050. *Atmos. Environ.*, 59: 567–577
- León-Chan, R.G., J.B. Heredia, T. Osuna-Enciso, J.A. Sañudo-Barajas, M. López-Meyer, L.A. Lightbourn-Rojas, J. León-Félix, 2017. Alterations in photosynthetic processes due to the effect of oxidative stress induced by cold and uv-b radiation. *Int. J. Agric. Biol.*, 19: 1363–1371
- Long, S.P., E.A. Ainsworth, A.D. Leakey and P.B. Morgan, 2005. Global food insecurity: treatment of major food crops with elevated carbon dioxide or ozone under large scale fully open air conditions suggests recent models may have overestimated future yields. *Phil. Trans. Royal Soc. B: Biol. Sci.*, 360: 2011–2020
- Ohara, T., H. Akimoto, J. Kurokawa, N. Horii, K. Yamaji, X. Yan and T. Hayasaka, 2007. An Asian emission inventory of anthropogenic emission sources for the period 1980–2020. *Atmos. Chem. Phys.*, 7: 6843–6902
- Olszyk, D.M. and C. Wise, 1997. Interactive effects of elevated CO₂ and O₃ on rice and flacca tomato. *Agric. Ecosyst. Environ.*, 66: 1–10
- Pang, J., K. Kobayashi and J. Zhu, 2009. Yield and photosynthetic characteristics of flag leaves in Chinese rice (*Oryza sativa* L.) varieties subjected to free-air release of ozone. *Agric. Ecosyst. Environ.*, 132: 203–211
- Shi, G.Y., L. Yang, Y. Wang, K. Kobayashi, J.G. Zhu, H. Tang, S. Pan, T. Chen, G. Liu and Y. Wang, 2009. Impact of elevated ozone concentration on yield of four Chinese rice cultivars under fully open-air field conditions. *Agric. Ecosyst. Environ.*, 131: 178–184
- Tang, H.Y., G. Liu, Y. Han, J.G. Zhu and K. Kobayashi, 2011. A system for free-air ozone concentration elevation with rice and wheat: control performance and ozone exposure regime. *Atmos. Environ.*, 45: 6276–6282
- Wang, Y.X., L. Yang, K. Kobayashi, J. Zhu, C.P. Chen, K. Yang, H. Tang and Y. Wang, 2012. Investigations on spikelet formation in hybrid rice as affected by elevated tropospheric ozone concentration in China. *Agric. Ecosyst. Environ.*, 150: 63–71
- Yang, L.X., Y.L. Wang, G.Y. Shi, Y.X. Wang and J.G. Zhu, 2008. Responses of rice growth and development to elevated near-surface layer ozone (O₃) concentration; a review. *J. Appl. Ecol.*, 19: 901–910
- Yong-Mei, H., Z. Fang-Dong, W. Jiong, Z. Yan-Qun, G. Xian-Hua and L. Yuan, 2017. Enhanced UV-B radiation inhibits photosynthesis, growth and yield of two rice landraces at Yuanyang terraces *in situ*. *Int. J. Agric. Biol.*, 19: 1379–1386
- Yuan, L.P., 1999. Hybrid rice development and use: innovative approach and challenges. In: *Proceedings 19th Session of the International Rice Commission*, pp: 77–85. Cairo, Egypt

(Received 30 March 2018; Accepted 06 August 2018)