



Full Length Article

Evaluating Boron Efficiency in Heat Tolerant Wheat Germplasm

Benjavan Rerkasem^{1*}, Sittichai Lordkaew², Narit Yimyam³ and Sansanee Jamjod⁴

¹Plant Genetic Resource and Nutrition Laboratory, Chiang Mai University, Chiang Mai, 50200, Thailand

²Center for Agricultural Resource Systems Research, Faculty of Agriculture, Chiang Mai University, Thailand 50200

³Department of Highland Agriculture and Natural Resources, Faculty of Agriculture, Chiang Mai University, Chiang Mai, 50200, Thailand

⁴Department of Plant and Soil Science, Faculty of Agriculture, Chiang Mai University, Chiang Mai, 50200, Thailand

*For correspondence: benjavan.r@cmu.ac.th

Abstract

Heat tolerance is a desirable trait for wheat in the sub-tropics and in traditional wheat areas under threat from global warming, where the crop may also be exposed to boron (B) deficiency. The 13th High Temperature Wheat Yield Trial (13HTWYT) from CIMMYT was evaluated with Fang 60, a 1977-bred variety as a B-efficient check, in a sand culture with (+B) and without (-B) added B in the nutrient solution in Chiang Mai, Thailand where temperature for the coolest month and during the time of heading both averaged 22.3°C. The wheat genotypes were affected differently in their grain set and grain yield by B deficiency (G x B, $p < 0.001$). Responses to B of the 13HTWYT expressed as grain yield in -B relative to +B correlated positively with the grain set index and the relative number of grains/spike, but not with the relative number spikes, while profuse tillering in -B was associated with lower yield. In -B grain yield also increased with the number of grains/spike. When B was not limiting, the grain yield increased with days to heading, number of tillers/plant, spikes/plant, spikelets/spike, grains/spike and grains/spikelet. Based on their grain set in -B, 39% of the 13HTWYT were considered B-inefficient, with grain set in fewer than half of competent florets, while 26% were B-efficient in the same order as Fang 60, 16% moderately efficient and 18% moderately inefficient. Boron efficiency is available in the 13HTWYT, suggesting that joint tolerance for B efficiency and heat stress can be selected if required. © 2019 Friends Science Publishers

Keywords: Boron; Grain set; Heat tolerance; Wheat; Yield formation

Introduction

The world's wheat is grown largely outside the tropics, with temperature for optimum growth at about 25°C (Briggle, 1980) and in the sub-tropics the crop is grown during the cool months, from October to February in the northern hemisphere. In Asia, where farmers face rising wage and water scarcity, wheat provides a promising dry season alternative to water-hungry rice and labour-intensive vegetables. However, tolerance to high temperature is essential for avoiding yield loss due to heat stress in much of the area that is subjected to average temperature for the coolest month of $> 17.5^\circ\text{C}$ (Fischer *et al.*, 2014), designated the wheat mega-environment 5 (ME5) by CIMMYT (the International Maize and Wheat Improvement Centre). Interest in heat tolerant wheat in India, which accounts for most of the global wheat in ME5, has been stimulated by observation of 34% yield loss due to heat stress in the country's major wheat cultivars and the concern about climate change (Mishra *et al.*, 2014). In addition to ME5, several other wheat mega-environments are also coming under threat of rising temperature (Braun and Payne, 2012).

Some of the first heat tolerant wheat genotypes were identified in the Hot Climate Wheat Screening Nursery, an exchange of wheat germplasm among breeders in hot environment as well as materials generated at CIMMYT for seven seasons in the 1980's (Mann, 1994). Since then wheat breeding for stress-prone environments has been more precisely targeted by the application of physiological understanding of tolerance to high temperature combined with economical and easy to use phenotyping methods such as the measurement of canopy temperature, notably for tolerance to heat and drought (Reynolds *et al.*, 2012).

In addition to the problem of heat stress, however, wheat yield in high temperature environments has also been reported to be depressed by boron (B) deficiency, including the Brazilian cerrados (Silva and de Andrade, 1983), south-western China (Yang, 1992), north-western states of India and along the Indo-Nepal border (Singh *et al.*, 1976; Tandon and Naqvi, 1992), to Nepal (Misra *et al.*, 1992; Subedi *et al.*, 1997) and Bangladesh (Reuter, 1987; Rerkasem, 1996). However, genotypic variation in responses to B is especially broad among crop species, including wheat (Rerkasem and Jamjod, 1997a). Boron-

efficient wheat genotypes may yield normally in the same soil in which B-inefficient genotypes suffer severe yield losses due to B deficiency, as has been found in CIMMYT's international wheat germplasm so far evaluated, although B-inefficient genotypes were in the majority (Rerkasem and Jamjod, 1997b; Rerkasem *et al.*, 2004). This paper reports on an evaluation of a relatively recent heat tolerant wheat germplasm from CIMMYT for B response and yield potential in a high temperature environment in comparison with Fang 60, a local B-efficient check.

Materials and Methods

The 13th High Temperature Wheat Yield Trial (13HTWYT) distributed by CIMMYT in 2014 with 49 entries and Fang 60 as local check was grown in a sand culture experiment at Chiang Mai University (18.81°N; 98.95°E). Each of the 50 entries was grown with 2 B treatments in duplicated drainable pots (Ø 30 cm, 30 cm deep) containing quartz sand with no detectible B, with 5 plants/pot. The pots were watered twice daily with 1 litre/pot of complete nutrient solution containing 10 µM H₃BO₃, 10 µM Fe-EDTA, 1 µM MnSO₄, 0.5 µM ZnSO₄, 0.2 µM CuSO₄, 0.1 µM CoSO₄, 0.1 µM Na₂MoO₄, 250 µM K₂SO₄, 1000 µM CaCl₂, 250 µM MgSO₄, 500 µM KH₂PO₄, and 5 mM KNO₃ (+B), or the same but without B added to nutrient solution (-B) (Rerkasem *et al.*, 2004). Days to heading were determined when 50% of the spikes had emerged. At maturity plants were harvested to determine seed yield, number of tillers/plant, spikes/plant and weight of individual seed for the whole pot (5 plants); number of spikelets/spike, number of grains/spike and grain set index (percentage of grain in the first two basal florets of 10 central spikelets) were determined on 10 spikes from each pot (main stem plus first tiller). Measures of performance of a genotype in B deficiency relative to B sufficiency were expressed as the -B value divided by the corresponding +B value.

Results

January was the coolest month during the experimental period, with monthly average temperature 22.3°C, mean minimum 16.8°C and maximum 29.1°C; 19 mm of rain fall in early November 2014 and 76 mm in early January 2015 (Fig. 1). Temperature from sowing to heading averaged 24.8°C, mean minimum 19.9°C and maximum 31.2°C, with temperatures during heading (21 December to 28 January) coincided exactly with the temperatures during the coolest month. Genotypic variation within the 13HTWYT was indicated by significant effects of genotypes and genotype by boron interaction on grain yield and some yield components (Fig. 2 and Table 2). Highly significant genotypic variation in the response to B was observed in grain yield, grain set index, and the number of tillers/plant, grains/spikelet, and grains/spike ($p < 0.001$), but not in the




Fig. 1: Temperature and rainfall in Chiang Mai during the experimental period, with sowing and heading time for the sand culture experiment

Source: Chiang Mai Meteorological Centre




Fig. 2: Grain yield (a), grain set index (b), yield components (c-h), and days to heading of the heat tolerant wheat germplasm 13HTWYT and boron efficient check Fang 60 in a sand culture with (+B, ■) and without (-B, □) added B (with standard deviation bars)

number of spikes/plant and weight of individual seed. Days to heading differed significantly among the 13HTWYT genotypes, which generally took longer (averaging 68±6 days) than the 58±1 days for Fang 60, but with no apparent effect of B. In +B, the number of grains/spikelets and grains/spike of the 13HTWYT genotypes averaged higher than Fang 60. The weight of individual seeds was generally higher in -B, although in +B it was not significantly different among the genotypes. The variation in the B response in the 13HTWYT was indicated by exceptionally large standard deviations in -B, with coefficient of variation for grain yield, number of grains/spike, grains/spikelet and grain set index ranging

Table 1: Relationship between grain yield and days to heading, yield components and grain set index of a heat tolerant wheat germplasm (13HTWYT) in sand culture with (+B) and without (-B) added boron in the nutrient solution

Boron treatment	+B	-B	+B	-B
	Correlation coefficient (r) ^a			
	With grain yield		With grain set index	
Grain yield (g pot ⁻¹)	-	-	0.05 (NS _{0.05})	0.90***
Days to heading	0.35 (<i>p</i> < 0.05)	0.10 (NS _{0.05})	-0.28 (NS _{0.05})	0.11 (NS _{0.05})
Tillers plant ⁻¹	0.55 (<i>p</i> < 0.001)	-0.52 (<i>p</i> < 0.001)	-0.00 (NS _{0.05})	-0.63***
Spikes plant ⁻¹	0.56 (<i>p</i> < 0.001)	0.03 (NS _{0.05})	0.00 (NS _{0.05})	-0.02 (NS _{0.05})
Spikelets spike ⁻¹	0.34 (<i>p</i> < 0.05)	0.15 (NS _{0.05})	-0.09 (NS _{0.05})	0.07 (NS _{0.05})
Grains spike ⁻¹	0.41 (<i>p</i> < 0.01)	0.93 (<i>p</i> < 0.001)	0.04 (NS _{0.05})	0.96 (<i>p</i> < 0.001)
Grains spikelet ⁻¹	0.34 (<i>p</i> < 0.05)	0.93 (<i>p</i> < 0.001)	0.18 (NS _{0.05})	0.96 (<i>p</i> < 0.001)
Seed weight (mg seed ⁻¹)	-0.12 (NS _{0.05})	-0.05 (NS _{0.05})	-0.06 (NS _{0.05})	-0.16 (NS _{0.05})
Grain set index (%)	0.05 (NS _{0.05})	0.90 (<i>p</i> < 0.001)	-	-

^aSignificance of linear association by analysis of variance in brackets

Table 2: Significant effects of genotype (G), boron (B), and G×B, by analysis of variance


Effect	Grain yield	Tillers/plant	Spikes/plant	Spikelets/spike	Grains/spike
G	***(3.6)	***(3.4)	***(2.3)	***(1.1)	***(6.3)
B	***(0.7)	***(0.7)	***(0.5)	***(0.2)	***(1.3)
G×B	***(5.1)	*(4.8)	NS _{0.05}	*(1.6)	***(9.0)
Effect	Grain set index	Grains/spikelet	Seed weight	Days to heading	
G	***(10.2)	***(0.33)	NS _{0.05}	***(7.2)	
B	***(2.0)	***(0.07)	***(4.9)	*(1.4)	
G×B	***(14.0)	***(0.47)	NS _{0.05}	NS _{0.05}	

(* *p* < 0.05; ** *p* < 0.01; *** *p* < 0.001), with LSD_{0.05} in brackets

from 53 to 65%.

Grain yield response to B of the 13HYWYT wheat genotypes, expressed as the yield in -B relative to +B ranged from 0.05 to 1.21 (the relative yield in the B-efficient check Fang 60 was 0.90), were closely and positively associated with their grain set index ($r=0.92$, $p < 0.001$) and the relative (-B/+B) number of grains/spike ($r=0.93$, $p < 0.001$), negatively with the relative number of tillers/plant ($r=-0.64$, $p < 0.001$), but not with the number of spikes/plant ($r=-0.03$, NS_{*p*<0.05}) (Fig. 3). The B status had different effects on the relationship between grain yield and days to heading and yield components (Table 1). When B was not limiting grain yield increased with days to heading and most of the yield components. In +B grain yield was positively correlated with days to heading ($r=0.35$, $p < 0.05$), spikes/plant ($r=0.56$, $p < 0.001$), spikelets/spike ($r=0.34$, $p < 0.05$), grains/spike ($r=0.41$, $p < 0.01$) and grains/spikelet ($r=0.34$, $p < 0.05$), but not with grain set index ($r=0.05$, NS_{0.05}) and weight of individual seeds ($r=-0.12$, NS_{0.05}). When B was limiting in -B grain yield correlated strongly and positively with the grain set index ($r=0.90$, $p < 0.001$), number of grains/spike ($r=0.93$, $p < 0.001$), and grains/spikelet ($r=0.93$, $p < 0.001$).

The 13HTWYT genotypes were classed into 4 B efficiency groups based on their grain set index in -B, in which the B-efficient check Fang 60 set grain normally (Fig. 4). The B-inefficient group, comprising 39% of the germplasm, set grain in less than half of competent florets in -B (grain set index < 50%), and produced grain yield averaging 23% of the yield in +B. The B-efficient group, accounted for 26% of the germplasm, had >85% grain set index in -B and indistinguishable grain yield in -B and +B,


Fig. 3: Association between grain yield in -B relative to +B and grain set index in -B (upper left), the number of grains/spike (upper right), tillers/plant (lower left) and spikes/plant (lower right), in -B relative to +B of a heat tolerant wheat germplasm (13HTWYT) with Fang 60 (●) as the B-efficient check. (Each data point is mean from 2 replications)

as in the B-efficient check Fang 60. The moderately B-efficient group (grain set index 71-85% in -B) accounted for 16% of the germplasm and in -B produced grain yield approaching that in +B. Moderately B-inefficient group (grain set index 51-70% in -B) accounted for 18% of the germplasm and in -B produced 55% of the grain yield in +B. Pedigrees of the B-efficient wheat genotypes from the 13HTWYT are listed in Table 3.

Table 3: Boron efficient genotypes in the 13HTWYT, with grain set index >85% in sand culture without added boron (-B)

Genotype		Grain set index in -B ^a	
1	PRL/2*PASTOR//WHEAR/SOKOLL	97.0	(107 ± 12)
2	QUAIU #1/SUP152	95.8	(111 ± 1)
3	FRANCOLIN #1	95.5	(121 ± 37)
4	KIRITATI//HUW234+LR34/PRINIA/3/CHONTE/5/PRL/2*PASTOR/4/CHOIX/STAR/3/HE1/3*CNO79//2*SERI	93.3	(128 ± 48)
5	FRET2*2/BRAMBLING//KIRITATI/2*TRCH/3/FRET2/TUKURU//FRET2	92.0	(84 ± 0)
6	KIRITATI//HUW234+LR34/PRINIA/3/FRANCOLIN #1/4/BAJ #1	90.8	(98 ± 18)
7	BAJ #1	90.8	(98 ± 18)
8	CROC_1/AE.SQUARROSA (213)//PGO/10/ATTILA*2/9/KT/BAGE//FN/U/3/BZA/4/TRM/5/ALDAN/6/SERI/7/VEE#10/8/OPATA/11/ATTILA*2/PBW65	90.0	(102 ± 11)
9	WHEAR/KIRITATI/3/C80.1/3*BATAVIA/2*WBLL1/4/BECARD	89.8	(107 ± 9)
10	KIRITATI//HUW234+LR34/PRINIA/3/CHONTE/5/PRL/2*PASTOR/4/CHOIX/STAR/3/HE1/3*CNO79//2*SERI	88.3	(99 ± 33)
11	WHEAR/KUKUNA/3/C80.1/3*BATAVIA/2*WBLL1*2/4/KIRITATI/2*TRCH	87.0	(112 ± 1)
12	FRANCOLIN #1*2//ND643/2*WBLL1	86.3	(97 ± 14)
13	FRANCOLIN #1/CHONTE//FRNCLN	85.3	(74 ± 8)
Boron efficient check, Fang 60 (PIIFDIPI/MZ/3/MXP)		97.0	(90 ± 7)

^a Grain yield in -B as % of the yield in +B ± standard deviation in brackets




Fig. 4: Average grain yield (\pm standard deviation) of wheat genotypes from the 13HTWYT classed by grain set index in -B (grain set index <50%, inefficient; 51-70%, moderately inefficient; 71-85%, moderately efficient; and efficient, >85%), compared with B-efficient check Fang 60

Discussion

The strong genotype by B interaction effects on grain yield, grain set index, number of tillers/plant, grains/spike and grains/spikelet, and the lack the interaction for days to heading, number spikes and spikelets and weight of individual grain mean that the relevance of these parameters to yield in the 13HTWYT need to be considered separately, when B was not limiting (+B) and when it did (-B). Significant correlation between grain yield and days to heading, number of tillers, spikes, spikelets and grains indicated that these parameters contributed positively to grain yield when B was not limiting. The 13HTWYT genotypes were clearly differentiated from Fang 60 by heading much later and having greater number of grains/spike and grains/spikelet (as the result of more competent florets in each spikelet) in +B. The positive correlation between grain yield and days to heading that

ranged from 59 to 79 days, found here contrasts with the wheat genotypes of the 1980's to 1990's that tended to have negative correlation between yield and days to heading when grown under high temperature. For example, grain yield of 84 wheat genotypes grown under heat stress was shown to correlate negatively ($r=-0.55$, $p < 0.01$) with days to heading ranged from 29 to 64 days, with an average of 54 days (Ortiz Ferrara *et al.*, 1994).

Accelerated development towards reproduction before the plant had developed sufficient leaf area and source capacity has been suggested as one of the ways in which wheat yield is limited by high temperature (Rawson, 1988). The advantages of longer time for vegetative development before flowering in the 13HTWYT, however, did not result in higher grain yield than Fang 60 (PIIFDIPI/MZ/3/MXP) which was bred in Pakistan in 1977, selected in Thailand and released in 1987 (Rerkasem, 1996). Although the number of tillers/plant and spikes/plant were strongly correlated with grain yield in +B, the 13HTWYT however generally had fewer spikes than Fang 60. The 13HTWYT was superior to Fang 60 by having more spikelets/spike, grains/spikelet and grains/spike, when B was not limiting. All these suggested some improvement in heat tolerance, as previously high temperature had been noted to reduce tillering, size of the spike, grain size and test weight, and thus adversely affecting biomass and grain yield (Rajaram, 1988).

The adverse effect of high temperature on wheat yield is likely to depend on the seasonal temperature pattern. For example, wheat in the Indo-Gangetic Plain is subjected to high temperature from flowering to maturity, whereas in Southeast Asia the wheat crop is exposed to high temperatures from sowing to maturity (Rajaram, 1988). Some heat tolerance studies of wheat have focused on the heat stress on reproduction and grain filling (Farooq *et al.*, 2011). This study has shed light on wheat adaptation to high temperature for the entire growing season, with grain yield of the heat tolerant wheat germplasm of the 13HTWYT responding positively to longer period of development to heading, tillering ability and number and size of the spikes

(more spikelets with more competent florets), when water and nutrients are not limiting.

Boron efficiency is an agronomic term for describing ability, without any mechanism being inferred, of plant genotypes to grow and yield well in soil in which B is limiting for other genotypes (Rerkasem and Jamjod, 1997b), which has been extended from the idea proposed and applied to zinc and nutrients in general (Graham, 1984). By this definition, wheat genotypes have been identified as B-inefficient (e.g., SW41, BL1022) and B-efficient (e.g., Sonora 64, NL460) in the field in Nepal, south-western China and Campinas in Brazil, as well as several locations in Thailand (Rerkasem, 1992; Subedi *et al.*, 1997). The mechanism for B efficiency in Fang 60, confirmed in field tests in Nepal and Thailand and in several sand culture trials (Anantawiroon *et al.*, 1997; Rerkasem and Jamjod, 1997b; Rerkasem *et al.*, 2004) has been demonstrated by tracing its ability to deliver limited ¹⁰B supply B from the roots to the developing anthers and pollen in the growing spike, while grain set in B-inefficient SW41 with the same limited external source of B failed because it was unable to do the same (Nachiangmai *et al.*, 2004). Thus Fang 60 was able to maintain sufficient level of B in its anthers in low B soil in which anther B concentration in B-inefficient genotypes was depressed by B deficiency (Rerkasem and Jamjod, 2004).

The 13HTWYT was predominantly B-inefficient, with grain yield in -B being determined by the adverse effect of B deficiency on grain set. Tillering was enhanced by B deficiency, just as lateral branching was stimulated by B deficiency that depressed apical growth in mung bean (Rerkasem *et al.*, 1990) and soybean (Rerkasem *et al.*, 1997). The degree to which tillering was increased by B deficiency in the 13HTWYT genotypes was inversely related to the level of their B efficiency, as indicated by the negative correlation between number of tillers and grain set index in -B. However, profuse tillering in -B was in fact detrimental to grain yield which declined with increasing number of tillers, even while grain yield in +B increased with increasing number of tillers. The international wheat germplasm from CIMMYT previously evaluated (the 4HTWYT, 2nd and 3rd International Breeding Nursery, 5th and 6th Hot Climate Wheat Screening Nursery, 17th Elite Selection Wheat Yield Trial, 18th Semi-arid Areas Wheat Screening Nursery and the 33rd International Bread Wheat Screening Nursery) were largely B-inefficient, but some B-efficient genotypes were also identified. Sixty genotypes from a total of 759 (8%) set grain normally along with Fang 60 under the condition in which 60% of the genotypes set grain in less than half of their competent florets (Rerkasem and Jamjod, 1997b; Rerkasem *et al.*, 2004). With grain set index in -B >85% in 13 of 49 genotypes, the 13HTWYT (distributed in 2014) was more B-efficient than the 4HTWYT (distributed in 1999) in which only 2 genotypes having grain set in -B in the same range as Fang 60.

Conclusion

Boron efficiency is a trait already available in the heat tolerant international wheat germplasm and can be easily selected for, if so desired. Where B is not limiting, the yield potential of the heat tolerant 13HTWYT might be increased simply by increasing the stand density by increasing seed rate to compensate for their limited tillering and number of spikes/plant.

Acknowledgments

The authors thank to CIMMYT for providing the 13HTWYT seed. Financial support from the National Research University project of Thailand's Higher Education Commission is acknowledged.

References

- Anantawiroon, P., K. Subedi and B. Rerkasem, 1997. Screening wheat for boron efficiency. *Dev. Plant Soil Sci.*, 76: 101–104
- Braun, H. and T. Payne, 2012. Mega-environment breeding. *In: Physiological Breeding I: Interdisciplinary Approaches to Improve Crop Adaptation*, pp: 5–17. Reynolds, M.P., A.J.D. Pask and D.M. Mullan (eds.). CIMMYT, Mexico DF
- Briggle, L.W., 1980. Origin and Botany of Wheat. *In: Wheat: Documenta Ciba-Geigy*, pp: 6–13. Häfliger, E., (eds.). Basle, Switzerland
- Farooq, M., H. Bramley, J.A. Palta and K.H.M. Siddique, 2011. Heat stress in wheat during reproductive and grain-filling phases. *Crit. Rev. Plant Sci.*, 30: 1–17
- Fischer, R.A., D. Byerlee and G.O. Edmeades, 2014. *Crop Yields and Global Food Security: Will Yield Increase Continue to Feed the World?* Australian Centre for International Agricultural Research, Canberra, Australia
- Graham, R.D., 1984. Breeding for nutritional characteristics in cereals. *Adv. Plant Nutr.*, 1: 57–102
- Mann, C.E., 1994. Seven years of hot climate wheat screening nurseries: 1985–91. *In: Wheat in Heat-Stressed Environments: Irrigated, Dry Areas and Rice-Wheat Farming Systems*, pp: 40–47. Saunders, D.A. and G.P. Hettel (eds.). CIMMYT, Mexico DF
- Mishra, S.C., S.K. Singh, R. Patil, N. Bhusal, A. Malik and S. Sareen, 2014. Breeding for heat tolerance in wheat. *In: Recent Trends on Production Strategies of Wheat in India*, pp: 15–29. Shukla, R.S., P.C. Mishra, R. Chatrath, R.K. Gupta, S.S. Tomar and I. Sharma (eds.). Jawaharlal Nehru Krishi Vishwa Vidyalaya, Madhya Pradesh, India and Indian Institute of Wheat and Barley Research: Haryana, India
- Misra, R., R.C. Munankarmi, S.P. Pandey and P.R. Hobbs, 1992. Sterility work in wheat at Tarahara in the eastern Terai of Nepal. *In: CIMMYT Wheat Special Report No. 11, Boron Deficiency in Wheat*, pp: 67–71. Mann, C.E. and B. Rerkasem (eds.). CIMMYT, Mexico DF
- Nachiangmai, D., B. Dell, R. Bell, L. Huang and B. Rerkasem, 2004. Enhanced boron transport into the ear of wheat as a mechanism for boron efficiency. *Plant Soil*, 264: 141–147
- Ortiz Ferrara, G., S. Rajaram and M.G. Mossad, 1994. Breeding strategies for improving wheat in heat-stressed environments. *In: Wheat in Heat-Stressed Environments: Irrigated, Dry Areas and Rice-Wheat Farming Systems*, pp: 24–32. Saunders, D.A. and G.P. Hettel (eds.). CIMMYT, Mexico DF
- Rajaram, S., 1988. Breeding and testing strategies to develop wheats for rice-wheat rotation areas. *In: Wheat Production Constraints in Tropical Environments*, pp: 187–196. Klatt, A.R. (ed.). CIMMYT, Mexico DF
- Rawson, H.M., 1988. Effects of high temperatures on the development and yield of wheat and practices to reduce deleterious effects. *In: Wheat Production Constraints in Tropical Environments*, pp: 44–62. Klatt, A.R. (eds.). CIMMYT, Mexico DF

- Rerkasem, B., 1996. A general survey of the incidence of wheat sterility. In: *ACIAR Proceedings No. 72, Sterility in Wheat in Subtropical Asia: Extent, Causes and Solution*, pp: 8–12. Rawson, H.M. and K.D. Subedi (eds.). Australian Centre for International Agricultural Research, Canberra, ACT
- Rerkasem, B., 1992. Boron probe nursery. In: *CIMMYT Wheat Special Report No. 11, Boron Deficiency in Wheat*, pp: 21–31. Mann, C.E. and B. Rerkasem (eds.). CIMMYT, Mexico DF
- Rerkasem, B. and S. Jamjod, 1997a. Genotypic variation in plant response to low boron and implications for plant breeding. *Plant Soil*, 193: 169–180
- Rerkasem, B. and S. Jamjod, 1997b. Boron deficiency induced male sterility in wheat (*Triticum aestivum* L.) and implications for plant breeding. *Euphytica*, 96: 257–262
- Rerkasem, B., S. Jamjod and S. Niruntrayagul, 2004. Increasing boron efficiency in many international bread wheat, durum wheat, triticale and barley germplasm will boost production on soils low in boron. *Field Crops Res.*, 86: 175–184
- Rerkasem, B., R.W. Bell, S. Lordkaew and J.F. Loneragan, 1997. Relationship of seed boron concentration to germination and growth of soybean (*Glycine max*). *Nutr. Cycl. Agroecosyst.*, 48: 217–223
- Rerkasem, B., R.W. Bell and J.F. Loneragan, 1990. Effects of seed and soil boron on early seedling growth of black and green gram (*Vigna mungo* and *Vigna radiata*). In: *Plant Nutrition-Physiology and Applications*, pp: 281-285. Springer, Dordrecht, The Netherlands
- Reuter, D.J., 1987. An assessment of wheat nutrition research in Bangladesh. *A Consultancy Report for CIMMYT and CIDA*. SAGRIC International, Adelaide, SA, Australia
- Reynolds, M., Y. Manes and G. Rebetzke, 2012. Application of physiology in breeding for heat and drought stress. In: *Physiological Breeding I: Interdisciplinary Approaches to Improve Crop Adaptation*, pp: 18–31. Reynolds, M.P., A.J.D. Pask and D.M. Mullan (eds.). CIMMYT, Mexico DF
- Silva, A.R.D. and J.M.V. de Andrade, 1983. Influence of micronutrients on the male sterility, on upland wheat and on rice and soybean yields, in red-yellow latosol. *Pesqui. Agropec. Bras.*, 18: 593–601
- Singh, H.M., S.D. Sinha and R.B. Prasad, 1976. Effect of boron on seed setting in wheat under North Bihar conditions. *Ind. J. Agron.*, 21: 100–101
- Subedi, K.D., C.B. Budhathoki and M. Subedi, 1997. Variation in sterility among wheat (*Triticum aestivum* L.) genotypes in response to boron deficiency in Nepal. *Euphytica*, 95: 21–26
- Tandon, J.P. and S.M.A. Naqvi, 1992. Wheat varietal screening for boron deficiency in India. In: *CIMMYT Wheat Special Report No. 11, Boron Deficiency in Wheat*, pp: 76–78. Mann, C.E. and B. Rerkasem (eds.). CIMMYT, Mexico DF
- Yang, Y.H., 1992. Wheat boron deficiency in Yunnan, China. In: *CIMMYT Wheat Special Report No. 11, Boron Deficiency in Wheat*, pp: 72–75. Mann, C.E. and B. Rerkasem (eds.). CIMMYT, Mexico DF

(Received 20 March 2018; Accepted 13 September 2018)