



**Full Length Article**

## Sucrose Metabolism and Accumulation is Disturbed during another Development in the T-Type Cytoplasmic Male Sterility of Wheat (*Triticum aestivum*)

Qingsong Ba<sup>1,2</sup>, Guiping Li<sup>1</sup>, Yulong Song<sup>2\*</sup>, Hongzhan Liu<sup>3\*</sup>, Gensheng Zhang<sup>1</sup>, Chu Chen<sup>1</sup>, Yunxian Song<sup>1</sup> and Zhaolin Fu<sup>1</sup>

<sup>1</sup>School of Life Science, Huaibei Normal University, Huaibei, 235000, Anhui, P.R. China

<sup>2</sup>Key Laboratory of Crop Heterosis of Shaanxi Province, Northwest A&F University, Yangling, 712100, Shaanxi, P.R. China

<sup>3</sup>College of Life Science and Agronomy, Zhoukou Normal University, Zhoukou, 466001, Henan, P.R. China

\*For correspondence: baqs1234@163.com

### Abstract

T-type cytoplasmic male sterility (T-CMS) plays a significant role in hybrid seed production of wheat. An obstruction of starch accumulation in the T-CMS pollen grain insinuated that an anomaly in sugar metabolism might be associated with pollen abortion. To elucidate the relationship between sugar metabolism and T-CMS, levels of sucrose levels and its decomposition products, activities of key enzymes and gene expression level involved in its sucrose metabolism at mononucleus stage were investigated. The content of sucrose in T-CMS anthers remained higher than that in the fertile line, but the contents of glucose and fructose in T-CMS anthers were dramatically decreased in contrast with the fertile line. The activities of cell wall bound invertase (CWIN), vacuolar invertase (VIN) and neutral invertase (NIN) during sucrose accumulation in anthers of T-CMS were also decreased to 17.4, 57.8 and 55.9%, respectively. The results of Real-time PCR also manifested that the expression levels of CWIN gene (*IVR1*), VIN gene (*IVR5*) and sucrose transporter gene (*TaSUT1*) were significantly down regulated in T-CMS anther. It showed that the accumulation of sucrose in T-CMS anthers might involve a decrease in activity and a reduction in content of invertase. It can be concluded that an inability to metabolize incoming sucrose to hexoses resulted in T-CMS pollen developmental lesion. © 2019 Friends Science Publishers

**Keywords:** Male sterility; Sucrose metabolism; Invertase; Wheat

### Introduction

Breeding of hybrid wheat is a promising pathway for increasing wheat production. Utilization of hybrid wheat could achieve nearly 15% increase in yield (Singh *et al.*, 2015). However, the nature of self-fertilization of wheat makes it difficult to hybrid seed production. Several pathways have been used to create male sterile systems which include engineered male sterility, male sterility induced by chemical killing agents, nuclear cytoplasmic male sterility, nuclear male sterility, and photo/thermo/photo-thermo/thermo-photosensitive male sterility system (Singh *et al.*, 2015). The discovery of cytoplasmic male sterility in *Aegilops-Triticum* crosses by Kihara established the initial interest in hybrid wheat breeding (Kihara, 1951). In subsequent studies, to develop commercial hybrid wheat, dependable male sterility systems were identified in the genetic background of *Triticum timopheevi* Zhuk. cytoplasm with the substitution of the nuclear genome of bread wheat (*T. aestivum*) by Wilson and Ross (1961).

Wheat T-type cytoplasmic male sterility (the cytoplasm came from *T. timopheevi* cytoplasm, T-CMS) (Mukai and Tsunewaki, 1979), which lacks negative cytoplasmic effects and possess more restoration resources, has been used commercially to a large extent (Singh *et al.*, 2010). Despite some research on mechanism of T-CMS, a lot of achievements have been scored (Hernould *et al.*, 1993; Hedgcoth *et al.*, 2002; Xu *et al.*, 2008; Sinha *et al.*, 2013). However, the underlying mechanism involved in abnormal carbohydrate metabolism in wheat T-CMS is not fully explored.

Sucrose could be unloaded from the phloem in anthers, which is capable of exporting to microspores (Lalonde *et al.*, 2003). Transport of sucrose during pollen development is the most important step for starch biosynthesis, and the downstream process of sucrose metabolism and starch synthesis depends on sucrose supply (Kong *et al.*, 2007). Anther-specific expression of sucrose transporter genes (SUTs) have been reported in some plant species such as tobacco (Lemoine *et al.*, 1999), *Arabidopsis* (Sivitz *et al.*, 2008), potato (KÜHN *et al.*, 2010) and rice

(Hirose, 2010; Eom *et al.*, 2016). In wheat, the homologous SUT genes may have crucial functions in transport of sucrose molecules (Aoki *et al.*, 2002; Deol *et al.*, 2013). Sucrose transport is accompanied by sucrose hydrolysis into hexose sugars by sucrose synthases (SUS: EC 2.4.1.13) and invertases (INV: EC 3.2.1.26), and the resulting hexoses serve as the major substrate for starch synthesis (Emes *et al.*, 2003). The *INV* catalyzed conversion of sucrose to glucose and fructose, and participate in sugar partitioning (Sturm, 1999). Invertases can be categorized as subcellular location, that is, neutral invertase (NIN), cell wall bound invertases (CWIN), and vacuolar invertase (VIN). In wheat, the cell wall bound invertases (*IVR1*) and the vacuolar invertase (*IVR5*) were cloned respectively, which were also abundantly present in the tapetum (Koonjul *et al.*, 2005). Moreover, inhibition of the anther specific *Nin88* gene (a cell wall bound invertase gene) impeded pollen development (Goetz *et al.*, 2001).

Carbohydrates have a variety of functions in the development of pollen (Goetz *et al.*, 2001; Hirsche *et al.*, 2009; Kunz *et al.*, 2014; Hirsche *et al.*, 2017). However, little or no starch was observed in the mature pollen in T-CMS (Joppa *et al.*, 1966). To better understand the relationship between sucrose metabolism and T-CMS, sucrose metabolism in wheat T-CMS anther was drilled down. Experiments have shown that the pollen abortion of the T-CMS lines mainly occurred at the mononucleate stage (Yao *et al.*, 2003), so the expression of *TaSUT1A*, *IVR1* and *IVR5* genes from mononucleus stage of anther were analyzed via quantitative real-time PCR (qRT-PCR). Moreover, invertase activity and sucrose content from the same stage of anthers were also determined in the T-CMS anthers.

## Materials and Methods

### Plant Materials

A wheat T-type CMS line (90-110, TA) and a maintainer line (90-110, TB) (Ba *et al.*, 2014), were sown in experimental plot in the Binhu campus of Huaibei Normal University, China. Fertility appraisal of the CMS line and its maintainer line were carried out for microspore observations with KI-I<sub>2</sub> solution staining. To identify the developmental stages, we collected some spikes using conventional acetic red dyeing techniques and 4',6'-diamidino-2-phenylindole staining method (Fig. 1A and B). We collected the anthers in the middle florets of spikes at mononucleus stage of pollen development from the main stems of TA and TB. The harvested anthers were placed immediately into liquid nitrogen and stored at -80°C.

### Microscopic Observation

Scanning electron microscope (JSM-6360LV, Japan) was used to observe pollen morphology. Fluorescence microscope tested pollen fertility using 0.1% iodine-potassium iodide staining method.

### Measurement of Sugar and Determination of Invertase Activity

The methods of testing sucrose and hexose were determined according to Oliver *et al.* (2005). The invertase activities were determined according to the optimization method (Koonjul *et al.*, 2005).

### Gene expression Analysis

The qRT-PCRs were performed for gene expression analyses according to the protocol. The primers of invertase (*IVR1* and *IVR5*) and one sucrose transporter gene (*TaSUTI*) were designed on the basis of Primer-Premier 5.0. Gene-specific primers (Zhu *et al.*, 2015) were used for gene expression analyses (Table 1). Furthermore, *18S* rRNA gene was as control for normalization, the relative transcript amount of each sample was expressed according to the optimization method (Yuan *et al.*, 2006). Data analysis was carried out through SPSS18.0 software.

## Results

### Pollen Grains in T-CMS are Depleted in Starch


Pollens from TB displayed starch staining (Fig. 1C), meanwhile, pollen is round (Fig. 1E). However, pollens from plants in TA did not exhibit any starch staining (Fig. 1D), and the absence of starch resulted in pollen wall depression (Fig. 1F). These results indicated that an obstruction of starch accumulation in TA.

### Anthers in Wheat T-CMS Accumulate Sucrose

To deepen understanding of mechanism of starch deficiency in T-CMS, we detected the levels of sucrose, glucose and fructose (Fig. 2). Sucrose level in TA was increased upto 3.2-fold ( $P < 0.01$ ; Fig. 2a), compared with TB. This suggests that sucrose accumulated in TA anthers is metabolized or relocated in the process of pollen development. Compared with TA, the levels of glucose and fructose in TB anthers fortified 1.2 times and 2.1 times, respectively. However, ratio of sucrose to hexoses (glucose and fructose) was more in TA anthers (1.18), than in TB anthers (0.19), which indicated that sucrose massively accumulated in wheat TA anthers. One possibility is that the process of converting sucrose into glucose is blocked.

### Changes in Anther Invertase Activity in Wheat T-CMS

In T-CMS anther, CWIN activity was significantly reduced compared with TB ( $\pm 49.2\%$ ;  $P < 0.01$ ; Fig. 3a). VIN activity in TA anther was nearly 5-fold lower than those in TB. Meanwhile the activity of NIN in TB was more than double those in TA. The significant reduction in invertase activity may result in disruption of sucrose conversion.



**Fig. 1:** Microspore development and pollen fertility in wheat  
A, B were stained with acetocarmine and 4',6-diamidino-2-phenylindole at uninucleate stage. Pollen grains of C and D stained with 1% I<sub>2</sub>-KI solution. Scanning electron microscopy analysis of the pollen surface from E and F. The pollens of C, E and D, F were from TB (maintainer line) and TA (T-type cytoplasmic male sterility line), respectively

### Alteration of Expression of Anther Invertase Genes in Wheat T-CMS

To explore the relationship between gene expression in sucrose metabolism and the observed changes of the sucrose levels, we used the *IVR1* and *IVR5* genes and *TaSUT1* gene to analyze the expression level (Fig. 4).

Compared with the maintainer TB, the transcription of *IVR1* in T-CMS was severely suppressed (56.2%,  $P < 0.01$ ), and *IVR5* transcription in T-CMS was nearly 4-fold decreased, simultaneously the *TaSUT1* in T-CMS was prominent in anthers ( $P < 0.01$ ).

### Discussion

Male sterility is often related to abnormal accumulation of starch, brought about by abiotic stresses. There are some clues demonstrating carbohydrate metabolism participates in the formation of microspores in adverse environmental conditions. Low temperatures during pollen development lead to depletion of starch in pollens (Oliver *et al.*, 2005). High temperature treatment reduced barley anther length, and impeded pollen starch formation (Sakata *et al.*, 2010). Drought-induced pollen abortion in rice is related to inhibition of starch accumulation in pollen (Sheoran and Saini, 1996), and drought-induced wheat male sterility is related to a drop of invertase activity in pollen (Dorion *et al.*, 1996).

Previous researches have demonstrated that the emergence of male sterile line is closely related to abnormal carbohydrate metabolism. Transcriptome analysis reveals starch metabolism had blocked male sterility in rapeseed. (Li *et al.*, 2015). Furthermore, sucrose metabolism was disturbed by chemical hybridizing agent SQ-1 with little starch in wheat pollen (Zhu *et al.*, 2015). Compared with the maintainer line, generous genes in cotton (*Gossypium*

*hirsutum* L.) genetic male sterility (GMS) lines were involved in starch and sucrose metabolism, and lack of starch accumulation was in male sterile pollen (Zhang *et al.*, 2015). Our observations also indicated that pollen grains of T-CMS fail to develop normally at maturity (Fig. 1F). Majority of pollens in T-CMS displayed little starch staining by I<sub>2</sub>-KI staining (Fig. 1D). Thus, an impediment in starch accumulation was in the T-CMS pollen grains. Thus, a decrease of starch granules is a universal phenomenon of multiple male sterility types.

An important finding of present study was the amount of glucose in anthers which was approximately 2–4 times greater than that of fructose throughout anther development (Fig. 2). Fructose resulting from sucrose cleavage was probably phosphorylated very rapidly, since pollen grains have very high fructokinase activity (Nikolov and Nikolov, 2010). The process of converting sucrose into glucose was blocked, and sucrose accumulated in T-CMS anthers (Fig. 2). This interruption of converting sucrose into glucose in pollen is probably a key contributor to T-CMS, because the supply of glucose to starch synthesis in pollen plays an important role in pollen fertility (Lee *et al.*, 2016).


Sucrose is generally converted to hexoses by invertase (INVs). In anthers, INVs are the crucial enzymes of sucrose conversion in pollen development (Saini and Westgate, 2000). As the experiment confirmed, CWINs play a dominant role in regulating the sucrose to hexose ratio (Kocal *et al.*, 2008). CWINs have been also shown to participate in different aspects of physiological actions (Albacete *et al.*, 2011, 2014). Tissue-specific antisense repression of cwINVs resulted in an early arrest of pollen development (Zanor *et al.*, 2009; Engelke *et al.*, 2010). Thus, the two genes could be deemed to regulate sucrose transport from the tapetum to wheat pollen grains. An inhibition of any of the steps in sucrose metabolism could limit starch accumulation in T-CMS pollen. Our data also revealed that sucrose vastly accumulated in T-CMS anthers (Fig. 2). Simultaneously, expression of the wheat anther *IVR1* and *IVR5* in T-CMS were inhibited (Fig. 4). Furthermore, expression of *IVR1* and *IVR5* were suppressed in T-CMS anthers, and the tendency of this suppression was similar to the reduction in invertase activities (Fig. 3). The reduced invertase activity affects the generation of a metabolic signal as it is apparently the case during pollen development (Lee *et al.*, 2016).

Sucrose transporter genes (SUTs) are prerequisite for conveying sucrose unloaded from the phloem to tissue sinks (Stadler *et al.*, 2005; Milne *et al.*, 2017). The homologous gene wheat SUT gene *TaSUT1* is expressed in large quantities in anthers (Sauer, 2007; Nguyen *et al.*, 2010). A significant decline in *TaSUT1* gene transcription manifested that sucrose inversion of sucrose was smothered (Fig. 4A). The results supported the former hypothesis that sucrose conversion in T-CMS was suffocated in T-CMS pollen, and supply of glucose to the developing microspores was restricted.


**Table 1:** Target genes for analysis of expression profiles

Gene name	Gene bank ID	Primer sequence (5'-3')	cDNA (bp)	annealing temperature
<i>18S</i>	AY049040	[F]AGTAAGCGCGAGTCATCAGCT [R]CATTCAATCGGTAGGAGCGAC	80	57
<i>TaSUT1</i>	AF408842	[F]GTGCTCTGATGGCTGATT [R]GAGGAACGGAAACCACTT	140	60
<i>IVR1</i>	AF030420	[F]TCGCCCTCAGGACATTG [R]CCAGACGCTTGTTCATCG	204	58
<i>IVR5</i>	AF069309	[F]TTCCTGTGCCTGTGCTCG [R]TCCGTCGGATACACCCCTC	123	58


F, forward primer; R, reverse primer. Primer reference



**Fig. 2:** Amount of sucrose (A), glucose (B) and fructose (C) of anthers from TA (T-type cytoplasmic male sterility line) and TB (maintainer line). Different lowercase letters mean significant differences



**Fig. 3:** Activities of cell wall invertase (A), vacuolar invertase (B) and neutral invertase (C) of anthers from TA (T-type cytoplasmic male sterility line) and TB (maintainer line). Different lowercase letters mean significant differences



**Fig. 4:** Expression of *TaSUT1A* (A), *IVR1* (B), and *IVR5* (C) of anthers from TA (T-type cytoplasmic male sterility line) and TB (maintainer line). Different lowercase letters mean significant differences

**Conclusion**

Invertase activity and its gene expression are inhibited in

T-CMS, as a consequence, inversion of sucrose is hindered in wheat anthers, accounting for absence of starch in T-CMS pollen.

## Acknowledgments

The research was supported by Anhui Provincial Natural Science Foundation (1708085QC62), Scientific Research Foundation of the Higher Education Institutions of Anhui Province, China (KJ2016A624), the Department of Science and Technology Planning Project of Henan Province (172102110165) and the Doctoral Scientific Research Foundation of Huaibei Normal University (15600971).

## References

- Albacete, A., E. Cantero-Navarro, M.E. Balibrea, D.K. Grobkinsky, M.D.L.C. Gonzalez, C. Martínez-Andújar, A.C. Smigocki, T. Roitsch and F. Pérez-alfocea, 2014. Hormonal and metabolic regulation of tomato fruit sink activity and yield under salinity. *J. Exp. Bot.*, 65: 6081–6095
- Albacete, A., D.K. Grosskinsky and T. Roitsch, 2011. Trick and treat: A review on the function and regulation of plant invertases in the abiotic stress response. *Phyton*, 50: 181–204
- Aoki, N., P. Whitfield, F. Hoeren, G. Scofield, K. Newell, J. Patrick, C. Offler, B. Clarke, S. Rahman and R.T. Furbank, 2002. Three sucrose transporter genes are expressed in the developing grain of hexaploid wheat. *Plant Mol. Biol.*, 50: 453–462
- Ba, Q.S., G.S. Zhang, N. Niu, S.C. Ma and J.W. Wang, 2014. Cytoplasmic effects on DNA methylation between male sterile lines and the maintainer in wheat (*Triticum aestivum* L.). *Gene*, 549: 192–197
- Deol, K.K., S. Mukherjee, F. Gao, A. Brülé-Babel, C. Stasolla and B.T. Ayele, 2013. Identification and characterization of the three homologues of a new sucrose transporter in hexaploid wheat (*Triticum aestivum* L.). *BMC Plant Biol.*, 13: 1–15
- Dorion, S., S. Lalonde and H.S. Saini, 1996. Induction of male sterility in wheat by meiotic-stage water deficit is preceded by a decline in invertase activity and changes in carbohydrate metabolism in anthers. *Plant Physiol.*, 111: 137–145
- Emes, M.J., C.G. Bowsher, C. Hedley, M.M. Burrell, E.S. Scrasefield and I.J. Tetlow, 2003. Starch synthesis and carbon partitioning in developing endosperm. *J. Exp. Bot.*, 54: 569–575
- Engelke, T., J. Hirsche and T. Roitsch, 2010. Anther-specific carbohydrate supply and restoration of metabolically engineered male sterility. *J. Exp. Bot.*, 61: 2693–2706
- Eom, J.S., D.N. Cong, D.W. Lee, S.K. Lee and J.S. Jeon, 2016. Genetic complementation analysis of rice sucrose transporter genes in *Arabidopsis* SUC2 mutant atsuc2. *J. Plant Biol.*, 59: 231–237
- Goetz, M., D.E. Godt, A. Guivarc'h, U. Kahmann, D. Chriqui and T. Roitsch, 2001. Induction of male sterility in plants by metabolic engineering of the carbohydrate supply. *Proc. Natl. Acad. Sci. U.S.A.*, 98: 6522–6527
- Hedgcock, C., A.M. El-Shehawi, P. Wei, M. Clarkson and D. Tamalis, 2002. A chimeric open reading frame associated with cytoplasmic male sterility in alloplasmic wheat with *Triticum timopheevi* mitochondria is present in several *Triticum* and *Aegilops* species, barley and rye. *Curr. Genet.*, 41: 357–365
- Hernould, M., S. Suharsono, S. Litvak, A. Araya and A. Mouras, 1993. Male-sterility induction in transgenic tobacco plants with an unedited atp9 mitochondrial gene from wheat. *Proc. Natl. Acad. Sci. U.S.A.*, 90: 2370–2374
- Hirose, T., 2010. Disruption of a gene for rice sucrose transporter, OsSUT1, impairs pollen function but pollen maturation is unaffected. *J. Exp. Bot.*, 61: 3639–3646
- Hirsche, J., J.M.G. Fernández, E. Stabentheiner, D.K. Grobkinsky and T. Roitsch, 2017. Differential effects of carbohydrates on arabidopsis pollen germination. *Plant Cell Physiol.*, 58: 691–701
- Hirsche, J., T. Engelke, D. Völler, M. Götz and T. Roitsch, 2009. Interspecies compatibility of the anther specific cell wall invertase promoters from *Arabidopsis* and tobacco for generating male sterile plants. *Theor. Appl. Genet.*, 118: 235–245
- Joppa, L.R., F.H. Mcneal and J.R. Welsh, 1966. Pollen and anther development in cytoplasmic male sterile wheat (*Triticum aestivum* L.). *Crop Sci.*, 6: 296–297
- Kihara, H., 1951. Substitution of nucleus and its effects on genome manifestations. *Cytologia*, 16: 177–193
- Kocal, N., U. Sonnewald and S. Sonnewald, 2008. Cell wall-bound invertase limits sucrose export and is involved in symptom development and inhibition of photosynthesis during compatible interaction between tomato and *Xanthomonas campestris* pv *vesicatoria*. *Plant Physiol.*, 148: 1523–1536
- Kong, J., Z.Y. Li, C. Wan, S. Li and Y. Zhu, 2007. Different gene expression patterns of sucrose-starch metabolism during pollen maturation in cytoplasmic male-sterile and male-fertile lines of rice. *Physiol. Plantarum*, 130: 136–147
- Koonjul, P.K., J.S. Minhas, C. Nunes, I.S. Sheoran and H.S. Saini, 2005. Selective transcriptional down-regulation of anther invertases precedes the failure of pollen development in water-stressed wheat. *J. Exp. Bot.*, 56: 179–186
- Kühn, C., W.P. Quick, A. Schulz, J.W. Riesmeier, U. Sonnewald and W.B. Frommer, 2010. Companion cell - specific inhibition of the potato sucrose transporter SUT1. *Plant Cell Environ.*, 19: 1115–1123
- Kunz, S., E. Pesquet and L.A. Kleczkowski, 2014. Functional dissection of sugar signals affecting gene expression in *Arabidopsis thaliana*. *PLoS One*, 9: 1–9
- Lalonde, S., M. Tegeder, M. Throne-Holst, W.B. Frommer and J.W. Patrick, 2003. Phloem loading and unloading of sugars and amino acids. *Plant Cell Environ.*, 26: 37–56
- Lee, S.K., J.S. Eom, S.K. Hwang, D. Shin, G. An, T.W. Okita and J.S. Jeon, 2016. Plastidic phosphoglucomutase and ADP-glucose pyrophosphorylase mutants impair starch synthesis in rice pollen grains and cause male sterility. *J. Exp. Bot.*, 67: 5557–5569
- Lemoine, R., L. Bürkle, L. Barker, S. Sakr, C. Kühn, M. Regnacq, C. Gaillard, S. Delrot and W.B. Frommer, 1999. Identification of a pollen - specific sucrose transporter - like protein NtSUT3 from tobacco. *FEBS Lett.*, 454: 325–330
- Li, Z.J., Y.F. Cheng, J.M. Cui, P.P. Zhang, H.X. Zhao and S.W. Hu, 2015. Comparative transcriptome analysis reveals carbohydrate and lipid metabolism blocks in *Brassica napus* L. male sterility induced by the chemical hybridization agent monosulfuron ester sodium. *BMC Genom.*, 16: 206–225
- Milne, R.J., J.M. Perroux, A.L. Rae, A. Reinders, J.M. Ward, C.E. Offler, J.W. Patrick and C.P. Grof, 2017. Sucrose transporter localization and function in phloem unloading in developing stems. *Plant Physiol.*, 173: 1330–1340
- Mukai, Y. and K. Tsunewaki, 1979. Basic studies on hybrid wheat breeding. *Theor. Appl. Genet.*, 54: 153–160
- Nguyen, G.N., D.L. Hailstones, M. Wilkes and B.G. Sutton, 2010. Role of carbohydrate metabolism in drought-induced male sterility in rice anthers. *J. Agron. Crop Sci.*, 196: 346–357
- Nikolov, N. and Z. Nikolov, 2010. Characterization of hexose kinases from camellia and lily pollen grains. *Physiol. Plantarum*, 81: 215–220
- Oliver, S.N., J.T.V. Dongen, S.C. Alfred, E.A. Mamun, X. Zhao, H.S. Saini, S.F. Fernandes, C.L. Blanchard, B.G. Sutton and P. Geigenberger, 2005. Cold-induced repression of the rice anther-specific cell wall invertase gene OSINV4 is correlated with sucrose accumulation and pollen sterility. *Plant Cell Environ.*, 28: 1534–1551
- Saini, H.S. and M.E. Westgate, 2000. Reproductive development in grain crops during drought. *Adv. Agron.*, 68: 59–96
- Sakata, T., T. Oshino, S. Miura, M. Tomabechei, Y. Tsunaga, N. Higashitani, Y. Miyazawa, H. Takahashi, M. Watanabe and A. Higashitani, 2010. Auxins reverse plant male sterility caused by high temperatures. *Proc. Natl. Acad. Sci. USA*, 107: 8569–8574
- Sauer, N., 2007. Molecular physiology of higher plant sucrose transporters. *FEBS Lett.*, 581: 2309–2317
- Sheoran, I.S. and H.S. Saini, 1996. Drought-induced male sterility in rice: Changes in carbohydrate levels and enzyme activities associated with the inhibition of starch accumulation in pollen. *Sex. Plant Reprod.*, 9: 161–169
- Singh, S.K., R. Chatrath and B. Mishra, 2010. Perspective of hybrid wheat research: a review. *Ind. J. Agric. Sci.*, 80: 1013–1027

- Singh, S.P., R. Srivastava and J. Kumar, 2015. Male sterility systems in wheat and opportunities for hybrid wheat development. *Acta Physiol. Plantarum*, 37: 1701–1713
- Sinha, P., S.M. Tomar, V.K. Singh and H.S. Balyan, 2013. Genetic analysis and molecular mapping of a new fertility restorer gene Rf8 for *Triticum timopheevi* cytoplasm in wheat (*Triticum aestivum* L.) using SSR markers. *Genetica*, 141: 431–441
- Sivitz, A.B., A. Reinders and J.M. Ward, 2008. *Arabidopsis* sucrose transporter AtSUC1 is important for pollen germination and sucrose-induced anthocyanin accumulation. *Plant Physiol.*, 147: 92–100
- Stadler, R., C. Lauterbach and N. Sauer, 2005. Cell-to-cell movement of green fluorescent protein reveals post. *Plant Physiol.*, 139: 701–712
- Sturm, A., 1999. Invertases. Primary structures, functions, and roles in plant development and sucrose partitioning. *Plant Physiol.*, 121: 1–8
- Wilson, J.A. and W.M. Ross, 1961. Cross-breeding in wheat, *Triticum aestivum*. i. frequency of the pollen-restoring character in hybrid wheats having *Aegilops ovata* cytoplasm. *Crop Sci.*, 3: 191–193
- Xu, P., Y. Yang, Z. Zhang, W. Chen, C. Zhang, L. Zhang, S. Zou and Z. Ma, 2008. Expression of the nuclear gene TaFAD is under mitochondrial retrograde regulation in anthers of male sterile wheat plants with timopheevii cytoplasm. *J. Exp. Bot.*, 59: 1375–1381
- Yao, Y.Q., G.S. Zhang, H.W. Liu and J.W. Wang, 2003. Cytomorphology and cytochemical localization of K-type and T-type cytoplasmic male sterile pollens in wheat. *J. Integr. Agric.*, 2: 591–596
- Yuan, J.S., A. Reed, F. Chen and C.N. Stewart, 2006. Statistical analysis of real-time PCR data. *BMC Bioinform.*, 7: 73–85
- Zanor, M.I., S. Osorio, A. Nunes-Nesi, F. Carrari, M. Lohse, B. Usadel, C. Kühn, W. Bleiss, P. Giavalisco and L. Willmitzer, 2009. RNA interference of LIN5 in tomato confirms its role in controlling Brix content, uncovers the influence of sugars on the levels of fruit hormones, and demonstrates the importance of sucrose cleavage for normal fruit development and fertility. *Plant Physiol.*, 150: 1204–1217
- Zhang, Y., J. Chen, J. Liu, M. Xia, W. Wang and F. Shen, 2015. Transcriptome analysis of early anther development of cotton revealed male sterility genes for major metabolic pathways. *J. Plant Growth Regul.*, 34: 223–232
- Zhu, W., S. Ma, G. Zhang, H. Liu, Q. Ba, Z. Li, Y. Song, P. Zhang, N. Niu and J. Wang, 2015. Carbohydrate metabolism and gene regulation during anther development disturbed by chemical hybridizing agent in wheat. *Crop Sci.*, 55: 868–876

(Received 06 June 2018; Accepted 19 June 2018)