



Full Length Article

Evaluation of Anti-Angiogenic Activity of Latex and Extracts of *Euphorbia helioscopia* using Chorioallontoic Membrane (CAM) Assay

Uzma Saleem^{1,2}, Bashir Ahmad^{*}, Mobasher Ahmad¹, Khalid Hussain¹, Nadeem Irfan Bukhari¹ and Muhammad Ashraf³

¹University College of Pharmacy, University of the Punjab, Lahore-Pakistan

²College of Pharmacy, Government College University, Faisalabad-Pakistan

³Department of Pharmacology, University of Veterinary and Animal Sciences, Lahore-Pakistan

*For correspondence: ahmadbprof@gmail.com

Abstract

The aim of this study was to investigate anti-angiogenic activity of *Euphorbia helioscopia* using CAM assay. Fertilized white leghorn chicken eggs were purchased from a local hatchery and placed in humidified incubator. On 5th day of incubation, a 2 cm wide window was cut at blunt ends of eggs and 4 mL of albumin was aspirated from each egg then each egg was sealed with parafilm tape and put back in incubator. On 6th day of incubation, 200 µL, containing 100 µg concentration of each sample / standard was applied over the growing embryo and again sealed and incubated. On 7th incubation day all the eggs were opened and images were taken with camera for quantification of CAM vessels using SPIP software. Sterilization was strictly maintained during the experiment. There were statistically significant variations in primary, secondary and tertiary blood vessels diameters and CAM areas of all the treated groups. Of all the extracts and latex, methanol leaves extract showed highest anti-angiogenic activity. From the data obtained in this study, it can be concluded that *Euphorbia helioscopia* possessed significant potential to cause inhibition of angiogenesis and this property could be useful in controlling oxidative stress induced angiogenesis. © 2015 Friends Science Publishers

Keywords: Anti-angiogenesis; *Euphorbia helioscopia* extracts; CAM assay

Introduction

Tumor growth can be controlled by inhibiting the angiogenesis. Judah Folkman was the first scientist who proposed anti-angiogenesis as a target site for treating tumors and decreasing metastasis (Folkman, 1971). Various *in vitro* (endothelial cells proliferation assay, endothelial cells migration assay, tube formation assay and aortic ring assay) and *in vivo* (matrigel plug assay, corneal angiogenesis assay and CAM assay) methods are available to evaluate pro-angiogenic and anti-angiogenic potentials of molecules (Auerbach *et al.*, 2003).

Angiogenic factors IL 8 and vascular endothelial growth factor (VEGF) production is stimulated with generation of free radicals in the cells (Brown *et al.*, 2000). VEGF, known as vascular permeability factor, is a key mediator in angiogenesis; it enhances the expression of endothelial nitric oxide synthase (eNOS), which may play important role in VEGF-induced angiogenesis (Bouloumie *et al.*, 1999). After binding with specific receptors present on endothelium VEGF promotes cellular events such as endothelial proliferation, migration and degradation of extracellular matrix components (Ferrara and Davis-Smyth, 1997). Oxidative stress increases the secretion of matrix metalloproteinase -1 (MMP-1), a collagenase that helps in

the vessel growth within tumor cells. Thus, oxidative stress can cause angiogenesis within carcinoma cells (Brown *et al.*, 2000). De novo synthesis of blood vessels plays pivotal role in tumour (Pepper, 1997) and warts growth (weblink 1) and also activates embryogenesis in female reproductive system (Fraser and Lunn, 2000).

Euphorbia helioscopia is an annual weed belonging to medicinally rich family *Euphorbiaceae* (Nadkarni, 2002) and is extensively grows in Pakistan. In China, this is a part of traditional Chinese medicine, used to treat various diseases such as tuberculosis, ascites, dysentery, edema, cervical cancer, lung cancer and esophageal cancer (Editorial Committee of the ABTCM, 1998; Pang and Lian, 2007; Yang *et al.*, 2007). Number of research findings have confirmed the anticancer, antioxidant, antibacterial, antiviral, antifungal, vasodepressor, anti-asthmatic, phytotoxicity, and molluscicidal activities of *Euphorbia helioscopia* (Ramezani *et al.*, 2008; Uzair *et al.*, 2009; Nikolova *et al.*, 2011; Ben-Mohamed *et al.*, 2012; Wang *et al.*, 2012). Secondary metabolites of the plant are responsible for its every pharmacological action. Flavonoids (Kawase and Kutani, 1968; Chen *et al.*, 1979), diterpenoids (Yamamura *et al.*, 1981; Shizuri *et al.*, 1983, 1984a,b; Kosemura *et al.*, 1985; Yamamura *et al.*, 1989; Zhang and Guo, 2006; Barile *et al.*, 2008; Tao *et al.*, 2008),

triterpenoids (Nazir *et al.*, 1998), steroids and lipids (Kosemura *et al.*, 1985), polyphenols (Wei-Sheng *et al.*, 2009) have been isolated in number of studies. Quercetin, a flavonoid obtained from plant, has been quantified in the leaves of *Euphorbia helioscopia* (Liu *et al.*, 2011) and its anticancer activity was confirmed by Wang *et al.*, (2012). The antineoplastic activity of *Euphorbia helioscopia* has been analyzed and confirmed by number of researchers (Kang and Liang, 1997; Cai *et al.*, 1999 a,b; Caltagirone *et al.*, 2000).

In Pakistan, the milky juice (latex) obtained from stem of *Euphorbia helioscopia* is being used as traditional remedy for the removal of warts from the fingers and success rate of this therapy is very high but no research data are available. In case of cancer/tumor, angiogenesis plays pivotal role in metastasis of cancer cells within the body. It is hypothesized that the understudied plant possessed compounds that have anti-angiogenic effect that cure the warts and it may also be helpful as an adjunct therapy with other anticancer drugs because it will inhibit vessels growth which is necessary to supply oxygen and nutrients to cancer cells and ultimately cancer will come in remission phase from progressive phase.

The aim of study was to investigate anti-angiogenic effect of latex and different extracts of *Euphorbia helioscopia* using *in vivo* CAM assay.

Materials and Methods

Chemicals

Ethanol (Analytical grade; BDH Laboratory), petroleum ether (analytical grade; Sigma Aldrich), methanol and chloroform (analytical grade; Merck, Germany), quercetin [QTN] (Merck Germany), phosphate buffer saline (PBS) were procured from local market of Lahore-Pakistan.

Plant Collection

The plant "*Euphorbia helioscopia*" was collected from suburbs of Lahore – Pakistan in the months of February and March. After identification and authentication by a Taxonomist of Botany Department, Government College University (GCU) Lahore-Pakistan, a voucher specimen (1501) was deposited to their herbarium. Leaves and stem were separated and dried under shade, then ground to fine powder separately which were later on used in extraction. The latex was collected in dried bottles by cutting the leafy part from the stem.

Preparation of Extract

The pulverized material from both parts of the plant was extracted separately at ambient temperature by maceration in water and ethanol as solvents. Then both the materials were extracted sequentially using solvents (petroleum ether,

chloroform, and methanol) in the order of increasing polarity by soxhlet apparatus. The solvents were removed from the extracts on rotary evaporator at 40°C.

UV and FTIR fingerprints of all the extracts of leaves and stems showed overlapping behavior in our earlier research (Saleem *et al.*, 2014b), keeping in view this result, only leaves extracts were selected for the current study.

Preparation of Chorioallantoic Membranes

Fertile white leghorn chicken eggs purchased from local hatchery (Big Bird-Lahore, Pakistan) were incubated at 37.5°C in humidified incubator (humidity: 55–60%) that was rotated at hly cycle upto 60° angle. Ethanol (70%) was used to sterilize the surface of all the eggs. On 5th day of incubation, a window of 2 cm in diameter was cut at blunt end of eggs and 4 mL of albumin was sucked from each egg with 21 gauge sterile syringe. This step was performed in order to separate the CAM from dry white membrane lying under the egg shell to get better quantification of CAM vasculature. Eggs were sealed with sterile parafilm tape and incubated for 24 h.

Preparation and Administration of Samples Solutions

PBS was used as vehicle in preparation of samples (leaves aqueous extract [L.AQ], leaves ethanol extract [L.ET], leaves petroleum ether extract [L.PE], leaves chloroform extract [L.CH], and leaves methanol extract [L.MT]) solutions of concentrations 10 µg/mL, 30 µg/mL, 50 µg/mL, 80 µg/mL, and 100 µg/mL. Standard (QTN) solution, 100 µg/mL, was prepared similarly. The pH of each solution was maintained between 6.5 and 7.5.

Two hundred microliter of each sample was applied over the growing embryos of all groups on 6th day of their incubation and again incubated for next 24 h.

Image Acquisition and Image Probing System (IPS) for the Quantification of Blood Vessels Growth over CAM

Chick embryos of all the treated, control and standard groups (n=10 per each group), aged seven days and staged 31-32 along with CAM were collected in separate petri dishes and images were recorded with COOLPIX (Nikon-china): wide 10X zoom, Lense-shift VR 16.0 megapixels camera. All images were cropped in Adobe Photoshop 6.0 to highlight the vessels on every image and then imported to SPIP version 6.2.5, an image-processing program that works on specific algorithm for automatic measurement of 3D surface roughness and related parameters to evaluate the anti-angiogenic response. For precise quantification, the x, y and z dimensions of each image were loaded to software.

Statistical Analysis

SPIP software version 6.2.5 was used for quantification of angiogenesis. Results were expressed as mean ± SD. One sample t-test was applied using SPSS version 12.

Graphs were drawn with graphpad prism version 4.00. Significance value was set at P < 0.05.

Results

Blood vessels growth pattern, diameter, 3D surface roughness, CAM areas of control, standard, L.AQ, L.ET, L.PE, L.CH, L.MT, and latex were observed and quantified. All the samples showed prominent results at 100 µg/mL, so remaining concentrations were excluded from the study in subsequent experimentation.

On macroscopic examination of CAMs, normal vascular architecture i.e. main “Y” branch (primary blood vessel) which further divided into secondary and tertiary blood vessels was found in control group. Quercetin (QTN) is a known anti-angiogenic compound, used as standard in the study. There was decrease in the secondary and tertiary blood vessels branching along with reduction in their thickness in all the extracts-treated groups. The branching pattern of blood vessels of latex, L.MT, L.CH, L.AQ and L.ET treated groups showed similarity with that of QTN treated group. L.PE treated group exhibited gross changes in vascular architecture (Fig. 1).

CAM areas of control and standard groups were 1372 mm² and 1228 mm² respectively. There was significant decrease (P < 0.05) in CAM areas of extracts-treated groups as compared to control (Fig. 2). The diameters of blood vessels growing on CAMs were also significantly reduced (P < 0.05) with respect to control group blood vessels (Fig. 3). It strengthens our hypothesis that *Euphorbia helioscopia* possesses anti-angiogenic activity.

Neovascularization on CAMs was quantified by evaluating 3D surface roughness parameters (Table 1). Average surface roughness values of control and standard were 7.534 ± 5.8 mm and 6.150 ± 3.5 mm respectively. There was decrease in average surface roughness of all the test groups, confirming anti-angiogenic effect of *E. helioscopia* (Table 1).

Anti-angiogenic activity of samples was calculated with following formula:

$$\% \text{ Inhibition of angiogenesis} = \frac{\text{CAM area of control} - \text{CAM area of standard or sample}}{\text{CAM area of control}} \times 100$$

QTN exhibited 10.50% anti-angiogenic effect. L.MT showed 11.64% inhibition of angiogenesis. The effect of latex was same as that of standard. L.AQ and L.PE showed minimum anti-angiogenic effect i.e. 5.39% among all the samples while L.CH and L.ET had 10.20% and 7.94% respectively (Fig. 4). There was non-significant difference in anti-angiogenic property of latex and L.CH extract as compared to standard while anti-angiogenic potential of L.MT extract was significantly greater than that of standard. L.AQ, L.ET and L.PE extracts showed significantly lower anti-angiogenic power with respect to that of standard (Fig. 4).




Fig 1: SPIP generated branching pattern and quantification of primary, secondary and tertiary blood vessels
1=Primary blood vessel, 2= Secondary blood vessel, 3= Tertiary blood vessel; A = Control, B = Standard (QTN), C = Methanol extract, D = Petroleum ether extract, E = Chloroform extract, F = Aqueous extract, G = Ethanol extract, H = Latex, a – h = quantification of blood vessels in term of diameter of above mentioned control, standard and test samples




Fig 2: Effect of different extracts and latex of *Euphorbia helioscopia* on CAM areas. *P < 0.05 as compared to control

Discussion

The CAM is a thin and extensively vascularized membrane. Due to its easy use and simplicity, the CAM assay is most commonly used to study normal angiogenesis and screen anti-angiogenic activity of new compounds (Patel *et al.*, 1994; Cao *et al.*, 1995; Friedlander *et al.*, 1995; Maragoudakis *et al.*, 1995; Hatjikondi *et al.*, 1996). The results of *in vitro* techniques, based on visual scoring to evaluate angiogenesis, are valid only when there is marked change in vascular area while minor changes due to physiological processes need accurate quantitative technique (Defouw *et al.*, 1989; Strick *et al.*, 1991; Danesi *et al.*, 1993; Nguyen *et al.*, 1994). Quantitative assessment of angiogenesis requires selection of appropriate parameters (Ejaz *et al.*, 2004). Thus, CAM areas, blood vessels diameter, surface roughness parameters were quantified with SPIP software to assess even minute anti-angiogenic effect.

Since long time, the natural products have key place in healthcare system (Goldman, 2001). The popularity of plant based medicine has also proved statistically by the Consumer Association in the United Kingdom indicating that in 1991 one out of four individuals is consuming such remedies whereas the corresponding estimate in 1985 was one in seven (British Medical Association, 1993). Newman and Cragg (2012) reported that approximately 85% of new drugs introduced from 1981-2010 have resulted from the research on natural products. Furthermore, the global documentation of more than 85000 plant species for medical use indicates the interest of scientists in this area (Balunas and Kinghorn, 2005).

There are few examples of anticancer drugs isolated from plants such as: vincristine and vinblastine, used to treat leukemias, lymphomas, advanced testicular cancer, breast and lung cancers, and Kaposi's sarcoma, topotecan to treat ovarian and small cell lung cancer and irrinotecan, is used in colorectal cancer (Cragg and Newman, 2005), etoposide and teniposide are used in lymphomas, bronchial and testicular cancer (Harvey, 1999; Cragg and Newman, 2005).

Anticancer activity of *Euphorbia helioscopia* has been studied on five cancer cell lines by Wang *et al.* (2012), and found promising effects on apoptosis, inhibitory effect on cell invasion and matrix metalloproteinase (MMP) 9 expression with ethyl acetate fraction (Wang *et al.*, 2012).

In the present study, different extracts of *Euphorbia helioscopia* were studied for anti-angiogenic effect. Average surface roughness and CAM areas of standard and all treated groups were lower than control group values. A direct relationship has been observed between CAM area and angiogenesis; it increases with increase in blood vessels growth and vice versa (Ejaz *et al.*, 2004).

There was statistically ($P < 0.05$) significant reduction in blood vessels diameters of extracts treated groups with respect to control value. Percent inhibition of angiogenesis was calculated and methanolic extract of leaves showed




Fig 3: Comparison of diameters of blood vessels of CAMs treated with different extracts / latex
P.B.V = Primary blood vessels, S.B.V = Secondary blood vessels, T.B.V = Tertiary blood vessels. * $P < 0.05$ with respect to control.




Fig 4: Anti-angiogenic effect of different extracts and latex of *Euphorbia helioscopia*. N.S. = non-significant, * $P < 0.05$ with respect to standard

maximum anti-angiogenic effect among all the treated groups. Leaves methanolic extract showed maximum *in vitro* antioxidant activity in DPPH and FRAP assay and high contents of flavonoids i.e., quercetin and kaempferol were quantified in the methanolic leaves extract in our

Table 1: 3D surface roughness parameters of normal and treated CAM

Groups	Roughness Parameters				
	Sa (mm)	Sq (mm)	Sz (mm)	Sv (mm)	Sp (mm)
Control	7.534 ± 5.8	9.772 ± 4.3	41.254 ± 8.3	21.962 ± 5.2	19.292 ± 4.9
Standard	6.150 ± 3.5	8.220 ± 6.4	41.300 ± 6.6	23.200 ± 7.6	18.100 ± 6.4
Latex	6.150 ± 4.3	7.970 ± 3.6	40.000 ± 9.1	22.500 ± 5.4	17.400 ± 3.9
L.MT	6.400 ± 3.3	8.340 ± 6.4	40.100 ± 4.5	23.100 ± 6.1	17.100 ± 5.3
L.PE	5.784 ± 5.6	7.594 ± 6.8	39.870 ± 6.2	21.530 ± 4.6	18.340 ± 6.3
L.CH	6.890 ± 6.2	8.759 ± 5.2	40.288 ± 4.8	20.448 ± 4.2	19.840 ± 6.2
L.AQ	7.164 ± 8.1	9.348 ± 5.9	41.523 ± 5.5	23.027 ± 3.6	18.495 ± 6.8
L.ET	6.025 ± 4.8	7.738 ± 7.5	39.506 ± 5.6	21.826 ± 4.6	17.68 ± 7.4

Each value represents mean ± SD of 10 observations.

Sa = average roughness, Sq = root mean square deviation, Sz = maximum height of the surface, Sv = reduce valley depth, Sp = reduce summit height

earlier work (Saleem *et al.*, 2014a,b,c). This suggests that highest anti-angiogenic activity of methanol leaves extract is due to its strong antioxidant activity which can be attributed to high flavonoids contents, quercetin and kaempferol.

Approximately 5000 polyphenols had been described till 2002 and categorized into subgroups flavonoids, lignans and isoflavones (Cao and Brakenhielm, 2002). Polyphenols are one of the abundantly found secondary metabolites in plants (Mojzis *et al.*, 2008; Jager *et al.*, 2009). Generally polyphenols with powerful antioxidant activity tend to possess good anti-angiogenic activity (Aisha *et al.*, 2010). For example quercetin (Tan *et al.*, 2003; Pratheeshkumar *et al.*, 2012), resveratrol (Cao *et al.*, 2005), rosmarinic acid (Huang and Zheng, 2006) and genistein (Su *et al.*, 2005) are polyphenols isolated from plant sources showed potent anti-angiogenic activity. These findings are in full agreement with data obtained in present study.

Conclusion

Euphorbia helioscopia has significant anti-angiogenic potential due to its high flavonoid contents. The folklore use of the latex obtained from this plant seems to be justified and attributed to anti-angiogenic activity. The latex / methanol extract can be exploited as adjuvant therapy with anti-neoplastic drugs to inhibit tumor growth/metastasis.

Acknowledgement

The authors gratefully acknowledge the financial grant from HEC-Pakistan and technical support by UVAS, Lahore-Pakistan

Reference

Aisha, A.M., K. Abu-Salah, Z. Ismail and A.A. Majid, 2010. Screening For antiangiogenesis activity in natural products: a review. *Webmed Cent. Pharmacol.*, 1: WMC001315

Auerbach, R., L. Rachel, S. Brenda, K. Louis, and A. Nasim, 2003. Angiogenesis assays: a critical overview. *Clinical Chem.*, 49: 132–140

Balunas, M.J. and A.D. Kinghorn, 2005. Drug discovery from medicinal plants. *Life Sci.*, 78: 431–441

Barile, E., M. Borriello, P.A. Di, D. Anges, F. Caterina, F. Ernesto, and V. Lanzotti, 2008. Discovery of a new series of jatrophone and lathyrane diterpenes as potent and specific P-glycoprotein modulators. *Org. Biomol. Chem.*, 6: 1756–1762

Ben-Mohamed, M.L., A. Jelassi, I. Hassen and M.S.O.B.A. Ould, 2012. Antioxidant proprieties of methanolic and ethanolic extracts of *Euphorbia helioscopia*, (L.) aerial parts. *Int. Food Res. J.*, 19: 1125–1130

Bouloumie, A., B. Valerie, K. Schini and B. Rudi, 1999. Vascular endothelial growth factor upregulates nitric oxide synthase expression in endothelial cells. *Cardiovascular Res.*, 41: 773–780

Brown, N.S., A. Jones, C. Fujiyama, A.L. Harris and R. Bicknell, 2000. Thymidine Phosphorylase Induces carcinoma cell oxidative stress and promotes secretion of angiogenic factors. *Cancer Res.*, 60: 6298–6302

British Medical Association, 1993. *Complementary medicine new approaches to good practice*. Oxford University Press, Oxford, UK

Cai, Y., J. Wang and B. Liang, 1999a. Antitumor activity of the root of *Euphorbia helioscopia* in vitro. *J. Chin. Med. Mater.*, 22: 85–87

Cai, Y., Y. Lu, B. Liang, G. Yu and Y. Xie, 1999b. Experimental study on antitumor activity of the root of *Euphorbia helioscopia* in vivo. *J. Chinese Med. Mater.*, 22: 579–581

Caltagirone, S., C. Rossi, A. Poggi, F.O. Ranelletti, P.G. Natali, M. Brunetti, F.B. Aiello and M. Piantelli, 2000. Flavonoids apigenin and quercetin inhibit melanoma growth and metastatic potential. *Int. J. Cancer*, 87: 595–600

Cao, Y., C. Chen, J.A. Weatherbee, M. Tsang and J. Folkman, 1995. Gro-b, a -C-X-C-chemokine, is an angiogenesis inhibitor that suppresses the growth of Lewis lung carcinoma in mice. *J. Exp. Med.*, 182: 2069–2077

Cao, Y., R. Cao and E. Brakenhielm, 2002. Antiangiogenic mechanisms of diet-derived polyphenols. *J. Nutr. Biochem.*, 13: 380–390

Cao, Y., Z.D. Fu, F. Wang, H.Y. Liu and R. Han, 2005. Anti-angiogenic activity of resveratrol, a natural compound from medicinal plants. *J. Asian Nat. Prod. Res.*, 7: 205–13

Chen, Y., Z. Tang, F. Jiang, X. Zhang and A. Lao, 1979. Studies on the active principles of Ze-Qi (*Euphorbia helioscopia* L.), a drug used for chronic bronchitis. *Acta Pharm. Sinica*, 14: 91–95

Cragg, G.M., and D.J. Newman. 2005. Plants as a source of anticancer agents. *J. Ethnopharmacol.*, 100: 72–79

Danesi, R., S.D. Bianchi, P. Soldani, A. Campagni, R.V. La Rocca, C.E. Myers, A. Paparelli and M.D. Tacca, 1993. Suramin inhibits bFGF induced endothelial cell proliferation and angiogenesis in the chick chorioallantoic membrane. *Br. J. Cancer*, 68: 932–938

Defouw, D.O., V.J. Rizzo, R. Steinfeld and R.N. Feinberg, 1989. Mapping of the microcirculation in the chick chorioallantoic membrane during normal angiogenesis. *Microvasc. Res.*, 38: 136–147

Editorial Committee of the Administration Bureau of Traditional Chinese Medicine, 1998. *Chinese Materia Medica.*, 4: 782–785

Ejaz, S., B.S. Kim and C.W. Lim, 2004. A novel image probing system for precise quantification of angiogenesis. *Tumori*, 90: 611–617

Ferrara, N. and T. Davis-Smyth, 1997. The biology of endothelial growth factor. *Endocrine Rev.*, 18: 4–25

Folkman, J., 1971. Tumor angiogenesis: therapeutic implications. *New England J. Med.*, 285: 1182–1186

Fraser, H.M. and S.F. Lunn, 2000. Angiogenesis and its control in the female reproductive system. *Br. Med. Bull.*, 56: 787–797

Friedlander, M., P.C. Brooks, R.W. Shaffer, C.M. Kincaid, J.A. Varner and D.A. Cheresh, 1995. Definition of two angiogenic pathways by distinct alpha v integrins. *Science*, 270: 1500–1502

Goldman, P., 2000. Herbal medicine today and roots of modern pharmacology. *Anna. Internal Med.*, 135: 594–600

Harvey, A.L., 1999. Medicines from nature: are natural products still relevant to drug discovery. *Trends Pharmacol. Sci.*, 20: 196–98

- Hatjickondi, O., P. Ravazoula, D. Kardamakis, J. Dimopoulos and S. Papaioannou, 1996. In vivo experimental evidence that the nitric oxide pathway is involved in the X-ray-induced antiangiogenicity. *Brit. J. Cancer*, 74: 1916–1923
- Huang, S.S. and R.L. Zheng, 2006. Rosmarinic acid inhibits angiogenesis and its mechanism of action in vitro. *Cancer Lett.*, 239: 271–280
- Jager, S., H. Trojan, T. Kopp, M.N. Laszczyk and A. Scheffler, 2009. Pentacyclic triterpene distribution in various plants - rich sources for a new group of multi-potent plant extracts. *Molecules*, 14: 2016–2031
- Kang, T.B. and N.C. Liang, 1997. Study on the inhibitory effects of quercetin on the growth of HL-60 leukemia cells. *Biochem. Pharmacol.*, 54: 1013–1018.
- Kawase, A. and N. Kutani, 1968. Some properties of a new flavonoid, tithymalin, isolated from the herbs of *Euphorbia Helioscopia* Linnaeus. *Agric. Biol. Chem.*, 32: 121–122
- Kosemura, S., Y. Shizuri and S. Yamamura, 1985. Isolation and structures of euphohelins, new toxic diterpenes from *Euphorbia helioscopia* L. *Bull Chem. Soc. Jpn.*, 58: 3112–3117
- Liu, H.P., X.F. Shi, Y.C. Zhang, Z.X. Li, L. Zhang and Z.Y. Wang, 2011. Quantitative analysis of quercetin in *Euphorbia helioscopia* L by RP/HPLC. *Cell Biochem. Biophys.*, 61: 59–64
- Maragoudakis, M.E., G.C. Haralabopoulos, N.E. Tsopanoglou and E. Pipili-Synetos, 1995. Validation of collagenous protein synthesis as an index for angiogenesis with the use of morphological methods. *Microvasc. Res.*, 50: 215–222
- Mojzis, J., L. Varinska, G. Mojziso, I. Kostova and L. Mirossay, 2008. Antiangiogenic effects of flavonoids and chalcones. *Pharmacol. Res.*, 57: 259–265
- Nadkarni, A.K., 2002. *Indian Materia Medica*. 3rded. Pp: 523. Popular Parkashan, Bombay, India
- Nazir, M., W. Ahmad and W. Kreiser, 1998. Isolation and NMR-assignments of 19alphaH-lupeol from *E. helioscopia* Linn (N.O. *Euphorbiaceae*). *Pak. J. Sci. Ind. Res.*, 41: 6–10
- Newman, D.J. and G.M. Cragg, 2012. Natural products as sources of new drugs over the 30 years from 1981–2010. *J. Nat. Prod.*, 75: 311–335
- Nguyen, M., Y. Shing and J. Folkman, 1994. Quantitation of angiogenesis and antiangiogenesis in the chick embryo Chorioallantoic membrane. *Microvasc. Res.*, 20: 531–539
- Nikolova, M., L. Evstatieva and T.D. Nguyen, 2011. Screening of plant extracts for antioxidant properties. *Bot. Serbica*, 35: 43–48.
- Pang, D.X. and J.W. Lian, 2007. Investigation on therapy of Ze Qi Decoction syndrome in diseases, synopsis of golden chamber. *China J. Trad. Chin. Med. Pharma.*, 22: 829–831
- Patel, P.C., R. Barrie, N. Hill, S. Landeck, D. Kurozawa and E.A. Woltering, 1994. Postreceptor signal transduction mechanisms involved in octreotide-induced inhibition of angiogenesis. *Surgery*, 116: 1148–1152
- Pepper, M.S., 1997. Manipulating angiogenesis. From basic sciences to besides. *Arterioscler Thromb Vasc. Biol.*, 17: 605–619
- Pratheeshkumar, P., A. Budhraj, Y.O. Son, X. Wang and Z. Zhang, 2012. Quercetin inhibits angiogenesis mediated human prostate tumor growth by targeting VEGFR-2 regulated AKT/mTOR/P70S6K signaling pathways. *PLoS ONE*, 7: e47516
- Ramezani, M., J. Benervan, M. Arab and F.S. Amer, 2008. Antiviral activity of *Euphorbia helioscopia* extracts. *J. Biol. Sci.*, 8: 809–813
- Shizuri, Y., S. Kosemura, J. Ohtsuka, Y. Terada and S. Yamamura, 1983. Structural and conformational studies on euphohelioscopins A and B and related diterpenes. *Tetrahedron Lett.*, 24: 2577–2580
- Shizuri, Y., J. Ohtsuka, S. Kosemura, Y. Terada and S. Yamamura, 1984b. Biomimetic reactions of some macrocyclic diterpenes. *Tetrahedron Lett.*, 25: 5547–5550
- Shizuri, Y., S. Kosemura, J. Ohtsuka, Y. Terada, S. Yamamura, S. Ohba, M. Ito and Y. Saito, 1984a. Structural and conformational studies on euphomin and related diterpenes. *Tetrahedron Lett.*, 25: 1155–1158
- Strick, D.M., R.L. Waycaster, J.P. Montani, W.J. Gay and T.H. Adair, 1991. Morphometric measurements of chorioallantoic membrane vascularity: effects of hypoxia and hyperoxia. *Amer. J. Physiol.*, 260: 1385–13891
- Su, S.J., T.M. Yeh, W.J. Chuang, C.L. Ho, K.L. Chang, H.L. Cheng, H.S. Liu, P.Y. Hsu and N.H. Chow, 2005. The novel targets for anti-angiogenesis of genistein on human cancer cells. *Biochem. Pharmacol.*, 69: 307–318
- Tan, W.F., L.P. Lin, M.H. Li, Y.X. Zhang, Y.G. Tong, D. Xiao and J. Ding, 2003. Quercetin, a dietary-derived flavonoid, possesses antiangiogenic potential. *Eur. J. Pharmacol.*, 459: 255–262
- Tao, H.W., X.J. Hao, P.P. Liu, and W.M. Zhu, 2008. Cytotoxic macrocyclic diterpenoids From *Euphorbia helioscopia*. *Arch. Pharm. Res.*, 31: 1547–1551
- Uzair, M., B.A. Loothar and B.A. Choudhary, 2009. Biological screening of *Euphorbia helioscopia* L. *Pak. J. Pharm. Sci.*, 22: 184–186
- Saleem, U., A. Bashir, H. Khalid, A. Mobasher, and I.B. Nadeem, 2014c. Simultaneous quantification of quercetin, myricetin and kaempferol in extracts and latex of *Euphorbia helioscopia* using RP-HPLC. *Asian J. Chem.*, in press
- Saleem, U., A. Bashir, H. Khalid, A. Mobasher, I.B. Nadeem and I. Saiqa, 2014a. Estimation of antioxidant power in various extracts of *Euphorbia helioscopia* L. with five different *in vitro* antioxidant models. *Asian J Chem.*, 26: 1241–1245
- Saleem, U., H. Khalid, A. Mobasher, I.B. Nadeem, M. Arif and A. Bashir, 2014 b. Physicochemical and phytochemical analysis of *Euphorbia helioscopia* (L.). *Pak. J. Pharm. Sci.*, 27: 577–585
- Wang, Z.Y., P.H. Liu, Y.C. Zhang, L.Q. Guo, Z.X. Li, and X.F. Shi, 2012. Anticancer potential of *Euphorbia helioscopia* L extracts against human cancer cells. *The Anat. Records*, 295: 223–233
- Weblink 1. www.webmd.com/skin...and.../warts-and-plantar-warts-topic-overview [Accessed on 16 February 2014]
- Wei-Sheng, F., G. Li, Z. Xiao-Ke, and W. Yan-Zhi, 2009. Polyphenols of *Euphorbia helioscopia*. *Chin. J. Nat. Med. Jan.*, 7: 37–39
- Yamamura, S., S. Kosemura, S. Ohba, M. Ito and Y. Saito, 1981. The isolation and structures of euphoscopins A and B. *Tetrahedron Lett.*, 22: 5315–5318
- Yamamura, S., Y. Shizuri, S. Kosemura, J. Ohtsuka, T. Tayama, S. Ohba, M. Ito, Y. Saito and Y. Terada, 1989. Diterpenes from *Euphorbia helioscopia*. *Phytochemistry*, 28: 3421–3436
- Yang, L., H.X. Chen and W.Y. Gao, 2007. Advances in studies on chemical constituents in *Euphorbia helioscopia* and their biological activities. *Chin. Trad. Herbal Drugs*, 38: 1585–1589
- Zhang, W. and Y.W. Guo, 2006. Chemical studies on the constituents of the Chinese medicinal herb *Euphorbia helioscopia* L. *Chem. Pharma. Bull.*, 54: 1037–1039

(Received 13 March 2014; Accepted 24 April 2014)